

Article (refereed) - postprint

This is the peer reviewed version of the following article:

Van der Ploeg, Martine J.; Baartman, Jantiene E.M.; Robinson, David A.
2018. **Biophysical landscape interactions: bridging disciplines and scale with connectivity**. *Land Degradation & Development*, 29 (4). 1167-1175,
which has been published in final form at <https://doi.org/10.1002/ldr.2820>

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

© 2017 John Wiley & Sons, Ltd.

This version available <http://nora.nerc.ac.uk/518820/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

This document is the author's final manuscript version of the journal article, incorporating any revisions agreed during the peer review process. There may be differences between this and the publisher's version. You are advised to consult the publisher's version if you wish to cite from this article.

The definitive version is available at <http://onlinelibrary.wiley.com/>

Contact CEH NORA team at
noraceh@ceh.ac.uk

BIOPHYSICAL LANDSCAPE INTERACTIONS: BRIDGING DISCIPLINES AND SCALE WITH CONNECTIVITY

Short Title: CONNECTIVITY FOR BIOPHYSICAL LANDSCAPE INTERACTIONS

Martine J. van der Ploeg¹, Jantiene E.M. Baartman¹, David A. Robinson²

¹Soil Physics and Land Management, Wageningen University, the Netherlands

²NERC- Centre for Ecology and Hydrology, Deiniol Road, Bangor, UK LL57 2UW.

Abstract

Landscape composition and land use impact the interactions between soil and vegetation. Differences in micro-behaviour, driven by the interplay of heterogeneous soil and vegetation dynamics, affect emergent characteristics across a landscape. Scaling approaches to understand the drivers of these emergent characteristics have been attempted, but the blueprint of interacting biophysical processes in landscapes is inherently messy and often still unknown. A complicating factor is single disciplinary focus in environmental sciences. Integrated knowledge is vital especially in view of future challenges posed by climate change, population growth and soil threats. In this paper we give examples of biophysical interactions which occur across various temporal and spatial scales and discuss how connectivity can be useful for bridging disciplines and scales to increase our understanding.

1 INTRODUCTION

The combination of climate change, population growth and soil threats including carbon loss, biodiversity decline and erosion increasingly challenge the global community (Schwilch et al., 2016). A major scientific challenge in understanding processes involved in soil threats, landscape resilience, ecosystem stability, sustainable land management and the economic consequences, is that it is an interdisciplinary field (Pelletier et al., 2012), requiring more openness between scientific disciplines (Liu et al., 2007). As a result of single disciplinary focus, ambiguity arises in the understanding of landscape interactions, especially interactions between biological and physical processes in a landscape (Cook & Hauer, 2007).

We think that integrated concepts of biophysical landscape interactions are needed to preserve ecosystem functioning in landscapes, especially in light of soil threats, population growth, climate change, and global water scarcity (Falkenmark, 1990; Schwilch et al., 2016). This requires interdisciplinary collaboration. An integrated concept can only be established by bridging the gap between several disciplines (Schulz et al., 2006; Seppelt et al., 2009), in a way that is appealing to those disciplines at the same time. Unfortunately, as evidence suggests, interdisciplinary work is more challenging to get funded (Bromham et al., 2016). The paper discusses how interdisciplinary challenges in biophysical landscape interactions at several scales can benefit from a connectivity approach.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/ldr.2820

Biophysical landscape interactions are those biotic and abiotic processes in a landscape that have an influence on the developments within and evolution of a landscape. Examples are the impact of soil heterogeneity on promoting coexistence of microbial life in the vadose zone (Long and Or, 2005), interaction between soil structure, hydraulics and climate and related effects of vegetation (Robinson et al., 2016; Reinsch et al., 2017), and cloud cover enhancement over forests (Teuling et al., 2017). An important aspect of biophysical landscape interactions is the different scales at which the various processes occur.

Scaling of environmental processes is possible, as long as the specific processes under consideration can be described by the same set of differential equations (Roth, 2008). Biophysical landscape interactions pose problems in this regard, because the combined physical and biochemical processes at different scales cannot be described by the same set of differential equations. For example, the description of the flow domain is scale dependent (Gelhar, 1986; Sánchez-Vila et al., 1996, Nieman & Rovey, 2009). The flow domain is that part of the system that can be mathematically described with one set of equations, such as the preferential flow domain, the groundwater flow domain, and the river channel flow domain. However, in scaling up, these domains may coincide and therefore no longer be described by the same equations.

There are two other complicating factors in understanding biophysical landscape interactions as well. While vegetation in many soil and hydrological models is approached physically, for example as a sink term in many soil water models, plant biology depends on more than physics alone (e.g. Wassen et al., 2013; Moreno de Las-Heras et al., 2016). And the response of vegetation to changing environmental conditions can include a possible, and often unknown, time-lag (e.g. Metzger et al., 2009). The interplay between the physical landscape and vegetation, which often co-evolve, and the resulting heterogeneity and emerging patterns is the reason it is so challenging to establish a theoretical basis for describing biophysical processes in landscapes.

In view of the considerable complexity of soil, numerical modelling is widely used to understand processes in soils, however the intricacies of biological responses in plants are mostly ignored. An integrative description for modelling biophysical interactions has been a long-standing goal in soil science (Vereecken et al., 2016). Figure 1 shows the scales involved in biophysical landscape interactions and examples of interdisciplinary challenges. In order to capture biophysical landscape interactions in models, it is important to find ways of dealing with feedbacks, the evolving heterogeneity when feedbacks play out differently, and with different scales. Interactions between ecology, hydrology, and geomorphology may be widely recognized, but they present grand challenges in themselves, especially the incorporation of feedbacks to understand system-level characteristics of landscapes.

Many disciplines involved in environmental research acknowledge spatial structure and heterogeneity in environmental systems (e.g. Schröder & Seppelt, 2006). Much research is focused on quantifying spatial structure and heterogeneity, such as climate variability, urban sprawl, deforestation and habitat loss (Ahlqvist & Shorridge, 2010). To be able to better understand the emerging patterns resulting from spatial structure and heterogeneity, connectivity has been acknowledged as a useful theoretical concept.

The concept has already been used in several (sub)disciplines: From an ecologists' perspective connectivity describes the understanding of water-mediated transfers of matter, energy and organisms (Pringle, 2001). Biologists often define connectivity as the degree to which a landscape facilitates or impedes the movement of individuals (Taylor et al., 1993). For hydrologists one definition is linked to how the hillslope's macropore network controls the flow through, replenishment of, and drainage from certain spots within the hillslope. This could lead to a hillslope scale connectedness of areas with relatively high hydraulic conductivity (Gomi et al., 2008). For soil

scientists, connectivity may for example relate pore structure and effective properties for solute transport (Vogel, 2000). For geomorphologists connectivity can be described as the physical (de)coupling of landforms (Baartman et al., 2013). Finally, for sedimentologists in erosion research, connectivity relates to the transfer of sediments through a basin (Bracken & Croke, 2007), either by wind or water movement.

This paper will apply connectivity to the challenges of integrated concepts for biophysical landscape interactions. We do this by briefly reviewing examples of (a) existing studies of biophysical interactions and (b) various scales at which these interactions take place. We then (c) introduce connectivity and outline how we think it can be used to bridge disciplines, (d) assess the connectivity concept for sustainable landscape management, and (e) address challenges across scales, including examples and ideas on how to quantify connectivity.

2 BIOPHYSICAL INTERACTIONS

Habitat manifestation is an expression of its evolutionary history. While the spatial distribution of habitats is largely driven by current climates and management, a soil's depth and water holding capacity will have played a role as the plant-soil system coevolves over time.

Within a landscape, microclimate and soil composition may differ, resulting in species adaptation to local conditions (Schenk & Jackson, 2002). Plants also exhibit adaptivity depending on environmental conditions; this was shown for African savannah grasses (Hartnett et al., 2013) and an alpine perennial herb (von Arx et al., 2012). Plants may even develop habitat specific, symbiotically-conferred stress tolerance (Rodriguez et al., 2008). Differences in drought sensitivity shape tree and shrub distribution in tropical forests at local and regional scales (Engelbrecht et al., 2007). Plants scan their environment biochemically, resulting in a myriad of internal information that specify its ecological niche (Trewavas, 2002).

Correlations between soil water availability and species distribution have been recognized since the last century (Schimper, 1903); and nowadays hydrogeophysical soil mapping enables visualization of above and below ground spatial connectivity patterns (Robinson et al., 2008). It is becoming increasingly clear that root-sourced signals appear to play a key role in regulating stomatal aperture in response to soil water availability (Bacon, 2004). Constant exposure to environmental stresses, biotic or abiotic, influences plant physiology, gene adaptations, and flexibility in gene adaptation (Van der Ploeg & Teuling, 2013). Addressing gene-expression and genotype adaptation is challenging as it may complicate modelling efforts in for example climate change impacts, because the precise response to changing conditions is unknown (e.g Rodriguez et al., 2008). Yet, climate change is expected to lead to more spatiotemporal variability and intensity in the water cycle (e.g. Vereecken et al., 2016). Productivity and survival are therefore not only the result of a plant's genotype, but depend critically on how fast and how severe environmental conditions change (Jones, 2007). Understanding the feedbacks between environmental change, and the subsequent signals and responses in a plant species, is crucial for understanding the effects of environmental stresses on vegetation in landscapes.

Many of these feedbacks come together in the concept of coevolution, which is being used in the context of evolving non-linear trends in the landscape (Pelletier et al. 2013). The coevolution concept includes the change in topography or morphology of the landscape in interaction with the climate, vegetation and hydrology. Related, catchment coevolution has been defined as the process of spatial and temporal interactions between water, energy, bedrock, sediments, carbon, ecosystems and

anthropogenic influences that lead to changes in catchment characteristics and responses (Troch et al., 2015). There is growing recognition of the importance of coevolution and biophysical interactions, which is needed to be able to better understand and sustainably manage our (natural) environment.

Pelletier et al. (2012) assessed coevolution within vegetation dynamics, pedogenesis and topographic development in southern California using a landscape modelling approach. They found strong correlations between Effective Energy and Mass Transfer (EEMT), above-ground biomass, soil thickness, hillslope-scale relief and mean distance-to-valley. Saco & Moreno de las Heras (2013) quantified the coevolution of vegetation and topography in semiarid areas. There, nonlinear interactions between physical and biological factors result in the emergence of remarkable landform related vegetation patterns such as striped and banded patterns. They found that variations in slope and abiotic or biotic factors can affect the vegetation patterns and resulting (micro)topography. Other work on coevolution includes e.g. D'Alpaos et al. (2007), who modelled the interplay of erosion, sedimentation and vegetation dynamics in tidal embayments; Perdigão & Bloeschl (2014) who investigated and quantified landscape-climate coevolution in Austria; and Jefferson et al. (2010) who studied the coevolution of hydrology and topography in a basalt topo-chronosequence landscape in Oregon, USA.

Coevolution is increasingly incorporated into models that simulate biophysical interactions. For example, a few models have started to treat soil formation as coevolution of a large number of soil parameters (Finke & Hutson, 2008). Other approaches include temporal changes of soil structure, a major determinant of water partitioning in the (sub)surface, and driver of biological activity, root growth and soil erosion (Leij et al., 2002; Stamati et al., 2013). A few models fully incorporate interactions between physical and biological processes (e.g. Laudone et al., 2011).

3. SCALES

Environmental and societal problems require an understanding of how processes operate at different scales, and how they can be linked across scales. Processes relevant in biophysical interactions in landscapes play a role at spatial scales ranging from millimetres to kilometres, i.e. from microbiology, such as soil microorganisms, to regional groundwater flow and landscape morphology, and at temporal scales ranging from seconds, e.g. earthquakes, to millennia, e.g. erosion and sculpting of landscapes by glaciers.

“Scaling refers to the transfer of understanding and of quantitative results from one spatial or temporal scale to another.” according to Roth (2008). Most properties in landscapes have some degree of correlation, but that depends on the scale at which observations have been made. Considering the scale at which processes occur and become visible, heterogeneity in biotic and abiotic processes is the most defining property of a landscape. Landscape heterogeneity accounts for markedly different system responses (e.g. Laudon et al., 2016). One striking example in hydrology is the emergence of scale dependency in transmissivity, which leads to an increase in effective transmissivity (or hydraulic conductivity) with an increase in observation scale (Sanchez et al., 1996; Schulze-Makuch et al., 1999; Nieman & Rovey, 2009; Fodor et al., 2011).

For biophysical processes an important factor for scaling is the soil moisture, which is dependent on soil physical properties (Robinson et al., 2016), landscape (Charpentier & Groffman, 1992), vegetation (Mohanty & Skaggs, 2001; Scanlon et al., 2007), and atmospheric conditions (Teuling et al., 2005). Combined, these factors regulate vadose zone processes including infiltration, permeability, water holding capacity and moisture loss rates. Techniques to monitor the resulting variables of heterogeneity become increasingly available, for example remote sensing allows surveys

covering large extents carried out at different scales (Lillesand et al., 2015) and hydrogeophysics allows mapping of the subsurface (Binley et al., 2015).

Scaling in time depends largely on the time-span involved because variables of a landscape system can change status from time independent to dependent or even be irrelevant (Schumm & Lichty, 1965). On a millennial timescale, the independent variables are, for instance, lithology, climate and vegetation, while hillslope and channel morphology are dependent variables and observed discharge and flow characteristics are considered indeterminable. Yet, on a short timescale (< 1 year), these indeterminable variables become the dependent ones.

Additionally, there is often a time-lag between an event (e.g. an extreme rainfall event, flood or landslide) and landscape or biophysical response (Phillips, 2003; Temme & Veldkamp, 2009). An extreme example is an earthquake, which may last only seconds, but may cause millennial-scale landscape responses, including increased erosion and catchment adjustment. In landscape ecology, time-lagged responses of biological variables to landscape modifications are widely recognised (e.g. Metzger et al., 2009). However, these are rarely considered in management plans. Understanding the ecological impacts of time-lagged responses to landscape modifications is critical for interpreting contemporary patterns of biodiversity (Royo et al., 2010).

4 POTENTIAL ROLE OF CONNECTIVITY

Connectivity is increasingly recognised as a major issue facing a hot, flat and crowded digital society (Friedman, 2009). In economics Didier Sornette has developed a new Dragon King (DK) theory for events that are generated by, or correspond to, changes in connectivity related mechanisms such as positive feedbacks, tipping points, bifurcations, and phase transitions. These phenomena are at the heart of connectivity and occur in nonlinear and complex systems, serving to amplify DK events to extreme levels, such as financial bubbles (Sornette and von der Berke, 2011).

What does the concept of connectivity offer for a better understanding of biophysical landscape interactions across various spatial and temporal scales? The key aspect of the connectivity concept is that it can create pathways for feedbacks which are often missing in soil models. Connectivity could thus play an important role in bridging disciplines and scales (Turnbull et al., 2008; Fryirs, 2012; Okin et al., 2015). Connectivity is dynamic over time and may change slowly or quickly, depending on the system and properties that are assessed. For example, connectivity in spatial landscape patterns changes slowly in response to dynamics of vegetation and soil processes, while connectivity changes quickly for example between and during rainfall events (Bracken & Croke, 2007; Wainwright et al., 2011).

The connectivity concept is a spatially explicit approach, by inclusion of neighbourhood effects (Peters et al., 2004), and therefore calls for inclusive information on flows or fluxes that connect different spatial units. Yet, connectivity has an advantage over spatially explicit modelling, where neighbourhood effects yield an increased parameter set and increased prediction uncertainty compared to non-spatial modelling. Instead, connectivity can be used to determine which spatial and temporal processes are likely to have an impact, and therefore the resulting modelling exercise can be simplified (Paola & Leeder, 2011). In modelling studies the connectivity concept is illustrated by using simple models and connecting different components of the system to learn more about the processes and feedbacks, for example for ecology (e.g. Tilman, 1994), hydrology (e.g. Porporato et al., 2003) and geomorphology (e.g. Saco et al., 2007).

Quantifying connectivity is challenging, according to recent discussion with scientists within the EU network project focussed on connectivity (COST1306 Action ‘Connecteur’ <http://connecteur.info/>). Connectivity is not often directly measured, but instead is inferred from other properties, such as soil texture distribution, moisture dynamics or the amount of discharge or sediment. We need to develop models where connectivity becomes a major part of the formulation, and tools that allow us to measure what we need to parameterise.

For biophysical landscape interactions, connectivity can be considered in terms of ‘high’, ‘medium’ and ‘low’ for the various processes at various scales, leading to various possible (sub)systems. Subsequently, for each (sub)system a particular process under consideration can be defined in terms of its connectivity status (i.e. high, medium, low), if it is related to other (sub)systems, and if it is relevant for the objective under investigation.

The classical example in which feedbacks between biotic and abiotic processes as well as (dis)connectivity play an important role is the vegetation patterns in semiarid landscapes, where isolated vegetation bands act as local sinks for water and nutrient flow, thereby disrupting sediment and hydrologic connectivity (e.g. Deblauwe et al., 2012). These banded patterns, consisting of alternating vegetated and bare bands, are formed because of an ecohydrological feedback system (Stewart et al., 2013). Differences in infiltration rates due to presence/absence of roots, macropores and soil aggregation in the vegetated/bare areas (Mora & Lazaro, 2013) lead to a runoff-runon mechanism (Saco et al., 2007). The runoff-runon mechanism is key for productivity, and disturbance of this ecohydrological feedback system can lead to severe land degradation (Okin et al., 2009; Moreno-de las Heras et al., 2011). Within this system, connectivity can be defined at several scales (see also Okin et al., 2015): (1) at the landscape scale, connectivity is low when the system is intact (Fig. 2a), because flow is limited to between two vegetated bands. Flow does not, or hardly, reach the end of the hillslope or outlet of the catchment. If the system is disrupted, connectivity on the landscape scale is significantly increased (Fig. 2b) and could lead to rill or gully erosion. (2) At a local scale, connectivity between individual plants is medium to high: on the one hand they compete for water, on the other hand, they help each other by providing e.g. increased organic matter underneath the plant patches (Verwijmeren et al., 2014). Finally, (3) at a micro scale, connectivity in the bare areas is relatively low because of crusting which inhibits percolation and infiltration, but high in the vegetated areas because of bioturbation (Klass et al., 2012).

Another example where connectivity can be quantified between the various (sub)systems is the agricultural system in NW Ethiopia (Tebebu et al., 2015; Zegeye et al., 2016) where soil erosion and sediment transfer to downstream reservoirs is an acute problem. Extensive soil and water conservation (SWC) measures have been introduced throughout the watershed to mitigate soil erosion. These measures are targeted at preventing erosion by on-field interventions such as soil bunds and grass barriers, and reducing sediment transfer pathways with sediment storage dams in gullies (Mekonnen et al., 2015). These interventions promote infiltration of precipitation into the soil and thereby reduce runoff and erosion. However, the side effect of increased infiltration is an increase in interflow from upstream to downstream parts of the catchment leading to shallower groundwater tables in the bottom of the catchment and faster saturation of the Vertisols during precipitation events (Tebebu et al., 2015). Therefore the upstream SWC measures in the hillslopes, eventually promote gully formation and expansion in the valleys (Zegeye et al., 2016). In this example connectivity can also be defined at various scales: (1) at the local scale, both the on-field (bunds and grass barrier) and between-fields (storage dams) SWC measures aim to reduce connectivity within the catchment as much as possible; however (2) at the landscape scale this resulted in increased connectivity between the upper and lower parts of the catchments through interflow processes, enhanced by the SWC measures.

Several studies have shown the usefulness of describing biophysical landscape interactions in terms of connectivity for more humid conditions as well. For intermontane depressional wetlands, landscape (sub-)surface hydrology, water chemistry and vegetation structure were highly connected to landscape scale processes (Cook & Hauer, 2007). Small but connected wetlands, stored more water longer, had higher productivity and different plant community composition compared to larger but isolated wetlands. At a more local scale, a wetland's ecology is determined by both frequency and duration of saturation and local groundwater quality, and species preference can thus be highly driven by microtopographical features (Van der Ploeg et al., 2012). In these systems the hydrology and vegetation are adjusted to each other. Despite such interplay between species distribution and complex patterning of ephemeral channels and streams, a relatively simple reservoir approach can capture the essence of drought and flood hazards (Van der Ploeg et al., 2012). Drier conditions trigger local connections between hummocks and hollows for hydrological routing, while wet conditions show a connected catchment response (Oosterwoud et al. 2017). Emergent vegetation properties can thus also be an indication of lateral hydrologic connectivity in addition to other controls (Hwang et al., 2012).

5 CONNECTIVITY FOR SUSTAINABLE LANDSCAPE MANAGEMENT

According to Venter et al. (2016), 75% of all terrestrial surface is experiencing human pressure, leading to declines of natural systems and biodiversity. Hydrological connectivity in the landscape itself is often declining due to anthropogenic influences such as dams and water diversions (Pringle, 2001). However, landscapes are also increasingly connected in terms of wind erosion when desertification comes into play (Okin et al., 2009). To manage these pressures, frameworks are being proposed for incorporating ecosystem services into land-management decision making (Schwilch et al., 2016). However, this remains challenging. The growing link between humans and ecosystem services necessitates a bridge between the question-driven, bioecology-centered spatial view and solution-driven society-centered holistic view (Wu, 2006).

In this sense, connectivity can be a very useful tool to assess how resilient a landscape is to change, and directs how it can be sustainably managed. For example, Jackson et al. (2013) applied connectivity in ecosystem service modelling to understand the impact of interventions like tree planting on flows of mass and energy.

6 CONNECTIVITY FOR EARTH-SYSTEM CHALLENGES

While connectivity is an ongoing area of research in ecology it has yet to fully migrate into our earth system science thinking. The problems humanity faces, such as climate change, require insight into the consequences of drivers such as extreme weather events and impacts on earth systems, their management and interventions we employ. Therefore, connectivity can be not only an interesting concept to describe emergent properties, but also a tool to evaluate management and mitigation strategies for sustainable land management (Okin et al., 2009, 2015; Bracken et al., 2013).

Insight into and acknowledgement of connectivity can be helpful, if not essential, for addressing key earth-system challenges, for example:

1. Natural flood management (NFM): To what extent can NFM alleviate flood risk? Research shows that the manipulation of the landscape at the hectare scale using shelter belts for example can reduce local flood risk (Marshall et al., 2009). However, the arrangement of vegetation and infrastructure features across the landscape speed up or slow down water and sediment movement, how this scales, and how this either synchronises, or desynchronises flows downstream remains

unknown. Quantifying connectivity within the arrangement of vegetation and infrastructure features would shed light on the most optimal design and therefore the benefits of natural flood management.

2. Soil carbon and nutrient cycling: What are the feedbacks from soil to climate and how will they impact change over the next century? Soils are estimated to store three times more carbon than plants or the atmosphere (Fischlin et al., 2007) and yet the connection between soils and global circulation models remains poor (Schmidt et al. 2011). Our understanding of the links between physics and biology to impact greenhouse gas emissions from soil is also limited (Reichstein et al., 2011; Blagodosky and Smith, 2012). Here connectivity could play an important role in understanding how carbon cycling is affected when soil hydraulics changes from uniform piston flow to bypass flow induced by soil water repellency.

Other challenges where connectivity concepts have already been successfully used include restoration of rivers (Reckendorfer et al., 2006; Jansson et al., 2007). In addition, dryland communities worldwide have implicitly used connectivity for their production systems for millennia (Okin et al., 2015). The scientific challenge will be to determine when complexity is important in biophysical landscape interactions and when it can essentially be ignored in models. Only as our modelling capability and understanding of complex phenomena increases will we be able to address this challenge. Remote sensing measurement techniques, such as LiDAR, UAVs, radar interferometry, hydrogeophysics and analysis of optical image data facilitate non-interfering observation of biophysical interactions on a landscape scale (e.g. Vogelmann et al., 2016; Lausch et al., 2013; Brake et al., 2013). A joint effort to connect Earth's (sub)surface processes by a combination of innovative big data-assimilation, measurement and modelling techniques will enable the scientific community to accurately address vital issues.

7 OUTLOOK

In this paper we have discussed how connectivity can be used to understand and connect biophysical processes at different spatial and temporal scales and why such a unifying concept is essential. Connectivity can bridge (sub)disciplines, although the differences in definitions and understanding need to be addressed: for example in ecology increased connectivity of ecosystems is seen as positive, while in soil science increased connectivity in erosion is negative. We need the next generation of models to incorporate connectivity and allow for the feedbacks that make earth system infrastructure so dynamic. This calls for development of models that are less focussed on detailed mechanistic understanding, and more focused on networks, connectivity and feedbacks while still incorporating the most important aspects of detailed mechanistic modelling (Paola & Leeder, 2011). Connectivity focused models hold promise for dealing with unprecedented levels of uncertainty in future trends of climate, population dynamics, economic development, and international trade barriers.

ACKNOWLEDGEMENTS

Special thanks to those involved in the discussion on this paper: Selamawit Amare, Coleen Carranza, Bram te Brake, Karrar Mahdi, Rens Masselink, Beatriz Ramirez, Ricardo da Silva. Van der Ploeg thanks the ES1308 COST CLIMMANI for their kind support to visit the CEH in Bangor (UK). We also thank Demie Moore and two anonymous reviewers for their useful suggestions.

REFERENCES

- Ahlqvist O, Shortridge A. 2010. Spatial and semantic dimensions of landscape heterogeneity. *Landscape Ecology* 25:573–590, doi:10.1007/s10980-009-9435-8
- Baartman JEM, Masselink R, Keesstra SD, Temme AJAM. 2013. Linking landscape morphological complexity and sediment connectivity. *Earth Surface Processes and Landforms* 38: 1457-1471, doi:10.1002/esp.3434
- Bacon MA. 2004. Water use efficiency in plant biology. In: Bacon MA (ed.) *Water use efficiency in plant biology*. Blackwell publishing, Oxford, UK
- Binley A, Hubbard SS, Huisman JA, Revil A, Robinson DA, Singha K, Slater LD. 2015. The emergence of hydrogeophysics for improved understanding of subsurface processes over multiple scales. *Water resources research* 51.6: 3837-3866, doi:10.1002/2015WR017016
- Blagodatsky S, Smith P. 2012. Soil physics meets soil biology: towards better mechanistic prediction of greenhouse gas emissions from soil. *Soil Biology and Biochemistry*, 47: 78-92, doi:10.1016/j.soilbio.2011.12.015
- Bracken LJ, Wainwright J, Ali GA, Tetzlaff D, Smith MW, Reaney SM, Roy AG. 2013. Concepts of hydrological connectivity: research approaches, pathways and future agendas. *Earth-Science Reviews* 119: 17-34, doi:10.1016/j.earscirev.2013.02.001
- Bracken LJ, Croke J. 2007. The concept of hydrological connectivity and its contribution to understanding runoff-dominated geomorphic systems. *Hydrological Processes* 21: 1749-1763, doi:10.1002/hyp.6313
- Brake B te, Hanssen RF, van der Ploeg MJ, de Rooij GH. 2013. Satellite-based radar interferometry to estimate large-scale soil water depletion from clay shrinkage: possibilities and limitations. *Vadose Zone Journal* 12.3, doi:10.2136/vzj2012.0098
- Bromham, L, Dinnage R, Hua X. 2016. Interdisciplinary research has consistently lower funding success. *Nature* 534: 684-687, doi:10.1038/nature18315
- Charpentier MA, Groffman PM. 1992. Soil moisture variability within remote sensing pixels. *Journal of Geophysical Research: Atmospheres* 97.D17: 18987-18995, doi:10.1029/92JD00882
- Cook BJ, Hauer FR. 2007. Effects of hydrologic connectivity on water chemistry, soils, and vegetation structure and function in an intermontane depression wetland landscape. *Wetlands* 27.3: 719-738, doi:10.1672/0277-5212(2007)27
- D'Alpaos A, Lanzoni S, Marani M, Rinaldo ACF. 2007. Landscape evolution in tidal embayments: Modeling the interplay of erosion, sedimentation, and vegetation dynamics. *Journal of Geophysical Research: Earth Surface* 112: F1, doi:10.1029/2006JF000537
- Deblauwe V, Couteron P, Bogaert J, Barbier N. 2012. Determinants and dynamics of banded vegetation pattern migration in arid climates. *Ecological Monographs* 82: 3-21, doi:10.1890/11-0362.1
- Engelbrecht BMJ, Comita LS, Condit R, Kursar TA, Tyree MT, Turner BL, Hubbell SP. 2007. Drought sensitivity shapes species distribution patterns in tropical forests. *Nature* 447, doi:10.1038/nature05747
- Falkenmark, M. 1990. Global water issues confronting humanity. *Journal of Peace Research*, 27(2): 177-190
- Finke PA., Hutson JL. 2008. Modelling soil genesis in calcareous loess. *Geoderma* 145.3: 462-479, doi:10.1016/j.geoderma.2008.01.017
- Fischlin A, Midgley GF, Price JT, Leemans R, Gopal B, Turley C, Rounsevell MDA, Dube OP, Tarazona J, Velicko AA. 2007. Ecosystems, their properties, goods and services. In *Climate Change 2007: Impacts, Adaptation and Vulnerability* (eds Parry, M. L., Canziani, O. F., Palutikof, J. P., van der Linden, P. J. & Hanson, C. E.) 211–272 (Cambridge Univ. Press, 2007).
- Fodor N, Sándor R, Orfanus T, Lichner L, Rajkai K. 2011. Evaluation method dependency of measured saturated hydraulic conductivity. *Geoderma* 165.1: 60-68, doi:10.1016/j.geoderma.2011.07.004
- Friedman TL. 2009. *Hot, flat, and crowded 2.0: Why we need a green revolution--and how it can renew America*. Macmillan.
- Fryirs K. 2012. (Dis)Connectivity in catchment sediment cascades: A fresh look at the sediment delivery problem. *Earth Surface Processes and Landforms* 38: 30-46, doi:10.1002/esp.3242View
- Gelhar LW. 1986. Stochastic subsurface hydrology from theory to applications. *Water Resources Research* 22.9S, doi:10.1029/WR022i09Sp0135S
- Gomi T, Sidle RC, Miyata S, Kosugi KI, Onda Y. 2008. Dynamic runoff connectivity of overland flow on steep forested hillslopes: scale effects and runoff transfer. *Water Resources Research* 44(8), doi:10.1029/2007WR005894
- Hartnett DC, Wilson GW, Ott JP, Setshogo M. 2013. Variation in root system traits among African semi-arid savanna grasses: Implications for drought tolerance. *Austral Ecology* 38.4: 383-392, doi:10.1111/j.1442-9993.2012.02422.x
- Hwang T, Band LE, Vose JM, Tague C. 2012. Ecosystem processes at the watershed scale: hydrologic vegetation gradient as an indicator for lateral hydrologic connectivity of headwater catchments. *Water Resources Research* 48.6, doi:10.1029/2011WR011301
- Jackson B, Pagella T, Sinclair F, Orellana B, Henshaw A, Reynolds B, ..., Eycott A. 2013. Polyscape: A GIS mapping framework providing efficient and spatially explicit landscape-scale valuation of multiple ecosystem services. *Landscape and Urban Planning* 112: 74-88, doi:10.1016/j.landurbplan.2012.12.014
- Jansson R, Nilsson C, Malmqvist B. 2007. Restoring freshwater ecosystems in riverine landscapes: the roles of connectivity and recovery processes. *Freshwater Biology* 52: 589-596, doi:10.1111/j.1365-2427.2007.01737.x
- Jefferson A, Grant GE, Lewis SL, Lancaster ST. 2010. Coevolution of hydrology and topography on a basalt landscape in the Oregon Cascade Range, USA. *Earth Surface Processes and Landforms* 35: 803-816, doi:10.1002/esp.1976

- Jones, HG. 2007. Monitoring plant and soil water status: established and novel methods revisited and their relevance to studies of drought tolerance. *Journal of Experimental Botany* 58(2): 119-130, doi: 10.1093/jxb/erl118
- Klass JR, Peters DPC, Trojan JM, Thomas SH. 2012. Nematodes as an indicator of plant–soil interactions associated with desertification. *Applied Soil Ecology* 58: 66-77, doi:10.1016/j.apsoil.2012.03.005
- Laudone GM, Matthews GP, Bird NRA, Whalley WR, Cardenas LM, Gregory AS. 2011. A model to predict the effects of soil structure on denitrification and N₂O emission. *Journal of Hydrology* 409.1: 283-290, doi:10.1016/j.jhydrol.2011.08.026
- Lausch A, Pause M, Merbach I, Zacharias S, Doktor D, Volk M, Seppelt R. 2013. A new multiscale approach for monitoring vegetation using remote sensing-based indicators in laboratory, field and landscape. *Environmental Monitoring and Assessment* 185: 1215–1235, doi:10.1007/s10661-012-2627-8
- Laudon H, Kuglerová L, Sponseller RA, Futter M, Nordin A, Bishop K, Lundmark T, Egnell G, Ågren AM. 2016. The role of biogeochemical hotspots, landscape heterogeneity, and hydrological connectivity for minimizing forestry effects on water quality. *Ambio* 45(Suppl 2): 152, doi:10.1007/s13280-015-0751-8
- Leij, FJ, Ghezzehei TA, Or D. 2002. Modeling the dynamics of the soil pore-size distribution. *Soil and Tillage Research* 64.1: 61-78, doi:10.1016/S0167-1987(01)00257-4
- Lillesand TM, Kiefer RW, Chipman JW. 2015. *Remote Sensing and Image Interpretation*. John Wiley & Sons
- Liu J, Dietz T, Carpenter SR, Alberti M, Folke C, Moran E, Pell AN, Deadman P, Kratz T, Lubchenco J, Ostrom E, Ouyang Z, Provencher W, Redman CL, Schneider SH, Taylor WW. 2007. Complexity of coupled human and natural systems. *Science* 317.5844: 1513-1516, doi:10.1126/science.1144004
- Long T, Or D. 2005. Aquatic habitats and diffusion constraints affecting microbial coexistence in unsaturated porous media. *Water resources research* 41.8, doi:10.1029/2004WR003796
- Marshall MR, Francis OJ, Frogbrook ZL, Jackson BM, McIntyre N, Reynolds B, Solloway I, Wheeler HS, Chell J. 2009. The impact of upland land management on flooding: results from an improved pasture hillslope. *Hydrological Processes*, 23(3), 464-475, doi:10.1002/hyp.7157
- Mekonnen M, Keesstra SD, Stroosnijder L, Baartman JEM, Maroulis J. 2015. Soil Conservation Through Sediment Trapping: A Review. *Land Degradation & Development* 26: 544-556, doi:10.1002/ldr.2308
- Metzger JP, Camargo Martensen A, Dixo M, Bernacci LC, Ribeiro MC, Godoy Teixeira AM, Pardini R. 2009. Time-lag in biological responses to landscape changes in a highly dynamic Atlantic forest region. *Biological Conservation* 142: 1166-1177, doi:10.1016/j.biocon.2009.01.033
- Mohanty BP, Skaggs TH. 2001. Spatio-temporal evolution and time-stable characteristics of soil moisture within remote sensing footprints with varying soil, slope, and vegetation. *Advances in Water Resources* 24.9: 1051-1067, doi:10.1016/S0309-1708(01)00034-3
- Mora JL, Lázaro R. 2013. Evidence of a threshold in soil erodibility generating differences in vegetation development and resilience between two semiarid grasslands. *Journal of Arid Environments* 89: 57-66, doi:10.1016/j.jaridenv.2012.10.005
- Moreno-de las Heras M, Díaz-Sierra R, Nicolau JM, Zavala MA. 2011. Evaluating restoration of man-made slopes: a threshold approach balancing vegetation and rill erosion. *Earth Surface Processes and Landforms* 36: 1367-1377, doi:10.1002/esp.2160
- Moreno-de las Heras M, Turnbull L, Wainwright J. 2016 Seed-bank structure and plant-recruitment conditions regulate the dynamics of a grassland-shrubland Chihuahuan ecotone. *Ecology* 97(9): 2303-2318, doi:10.1002/ecy.1446
- Niemann, WL, and Rovey II CW. 2009. A systematic field-based testing program of hydraulic conductivity and dispersivity over a range in scale. *Hydrogeology Journal* 17.2: 307-320, doi:10.1007/s10040-008-0365-3
- Okin GS, Parsons AJ, Wainwright J, Herrick JE, Bestelmeyer BT, Peters DC, Fredrickson EL. 2009. Do Changes in Connectivity Explain Desertification? *BioScience* 59: 237-244, doi:10.1525/bio.2009.59.3.8
- Okin GS, Heras MM-dl, Saco PM, Throop HL, Vivoni ER, Parsons AJ, Wainwright J, Peters DPC. 2015. Connectivity in dryland landscapes: shifting concepts of spatial interactions. *Frontiers in Ecology and the Environment* 13:20-27, doi:10.1890/140163
- Oosterwoud M, van der Ploeg M, van der Schaaf S, van der Zee S. 2017. Variation in hydrologic connectivity as a result of microtopography explained by discharge to catchment size relationship. *Hydrological Processes* 31:2683–2699. doi:10.1002/hyp.11164
- Paola C, Leeder M. 2011. Environmental dynamics: Simplicity versus complexity. *Nature* 469: 38-39, doi:10.1038/469038a
- Pelletier JD, DeLong SB, Orem CA, Becerra P, Compton K, Gressett K, Lyons-Baral J, McGuire LA, Molaro JL, Spinler JCCF. 2012. How do vegetation bands form in dry lands? Insights from numerical modeling and field studies in southern Nevada, USA. *Journal of Geophysical Research: Earth Surface* 117: F04026, doi:10.1029/2012JF002465
- Pelletier JD, Barron-Gafford GA, Breshears DD, Brooks PD, Chorover J, Durcik M, Harman CJ, Huxman TE, Lohse KA, Lybrand R, Meixner T, McIntosh JC, Papuga SA, Rasmussen C, Schaap M, Swetnam TL, Troch PA. 2013. Coevolution of nonlinear trends in vegetation, soils, and topography with elevation and slope aspect: A case study in the sky islands of southern Arizona. *Journal of Geophysical Research: Earth Surface* 118.2: 741-758, doi: 10.1002/jgrf.20046
- Perdigão RAP, Blöschl G. 2014. Spatiotemporal flood sensitivity to annual precipitation: Evidence for landscape-climate coevolution. *Water Resources Research* 50: 5492-5509, doi:10.1002/2014WR015365
- Peters DPC, Herrick JE. 2004. Strategies for ecological extrapolation. *Oikos* 106.3: 627-636, doi:10.1111/j.0030-1299.2004.12869.x
- Phillips JD. 2003. Sources of nonlinear complexity in geomorphic systems. *Progress in Physical Geography* 26: 339–361

- Pringle CM. 2001. Hydrologic connectivity and the management of biological reserves: a global perspective. *Ecological Applications*, pp.981-998, doi:10.1890/1051-0761(2001)011[0981:HCATMO]2.0.CO;2
- Porporato A, D'odorico P, Laio F, Rodriguez-Iturbe I. 2003. Hydrologic controls on soil carbon and nitrogen cycles. I. Modeling scheme. *Advances in Water Resources* 26.1: 45-58, doi:10.1016/S0309-1708(02)00094-5
- Reckendorfer W, Baranyi C, Funk A, Schiemer F. 2006. Floodplain restoration by reinforcing hydrological connectivity: expected effects on aquatic mollusc communities. *Journal of Applied Ecology* 43: 474-484, doi:10.1111/j.1365-2664.2006.01155.x
- Reichstein M, Bahn M, Ciais P, Frank D, Mahecha MD, Seneviratne SI, Zscheischler J, Beer C, Buchmann N, Frank DC, Papale D. 2013. Climate extremes and the carbon cycle. *Nature*, 500(7462): 287-295, doi:10.1038/nature12350
- Reinsch S, Koller E, Sowerby A, de Dato G, Estiarte M, Guidolotti G, Kovács-Láng E, Kröel-Dulay G, Lellei-Kovács E, Larsen KS, Liberati D, Penuelas J, Ransijn J, Robinson DA, Schmidt IK, Smith AR, Tietema A, Dukes JS, Emmett BA. 2017. Shrubland primary production and soil respiration diverge along European climate gradient. *Scientific Reports* 7: 43952, doi:10.1038/srep43952
- Robinson DA, Abdu H, Jones SB, Seyfried M, Lebron I, Knight R. 2008. Eco-geophysical imaging of watershed-scale soil patterns links with plant community spatial patterns. *Vadose Zone Journal* 7(4): 1132-8, doi:10.2136/vzj2008.0101
- Robinson DA, Jones SB, Lebron I, Reinsch S, Domínguez MT, Smith AR, Jones DL, Marshall MR, Emmett BA. 2016. Experimental evidence for drought induced alternative stable states of soil moisture. *Scientific reports* 6, doi:10.1038/srep20018
- Rodriguez RJ, Henson J, Van Volkenburgh E, Hoy M. 2008. Stress tolerance in plants via habitat-adapted symbiosis. *ISME Journal* 2: 404-416, doi:10.1038/ismej.2007.106
- Royo AA, Stout SL, deCalesta DS, Pierson TG. 2010. Restoring forest herb communities through landscape-level deer herd reductions: Is recovery limited by legacy effects? *Biological Conservation* 143: 2425-2434, doi:10.1016/j.biocon.2010.05.020
- Roth K. 2008. Scaling of water flow through porous media and soils. *European journal of soil science* 59.1: 125-130, doi:10.1111/j.1365-2389.2007.00986.x
- Saco PM, Moreno-de las Heras M. 2013. Ecogeomorphic coevolution of semiarid hillslopes: Emergence of banded and striped vegetation patterns through interaction of biotic and abiotic processes. *Water Resources Research* 49: 115-126, doi:10.1029/2012WR012001
- Saco PM, Willgoose GR, Hancock GR. 2007. Eco-geomorphology of banded vegetation patterns in arid and semi-arid regions. *Hydrol. Earth Syst. Sci.* 11: 1717-1730
- Sánchez-Vila X, Carrera J, Girardi JP. 1996. Scale effects in transmissivity. *Journal of Hydrology* 183.1: 1-22, doi:10.1016/S0022-1694(96)80031-X
- Scanlon TM, Caylor KK, Levin SA, Rodriguez-Iturbe I. 2007. Positive feedbacks promote power-law clustering of Kalahari vegetation. *Nature* 449.7159: 209-212, doi:10.1038/nature06060
- Schenk HJ, Jackson RB. 2002. The global biogeography of roots. *Ecological Monographs* 72: 311-328, doi:10.1890/0012-9615(2002)072[0311:TGBOR]2.0.CO;2
- Schimper AFW. 1903 *Plant geography upon a physiological basis*. Clarendon, Oxford
- Schmidt MW, Torn MS, Abiven S, Dittmar T, Guggenberger G, Janssens IA, Kleber M, Kögel-Knabner I, Lehmann J, Manning DA, Nannipieri P. 2011. Persistence of soil organic matter as an ecosystem property. *Nature*, 478(7367): 49-56, doi:10.1038/nature10386
- Schröder B, Seppelt R. 2006. Analysis of pattern-process interactions based on landscape models—overview, general concepts, and methodological issues. *Ecological modelling* 199.4: 505-516, doi:10.1016/j.ecolmodel.2006.05.036
- Schulz K, Seppelt R, Zehe E, Vogel, HJ, Attinger S. 2006. Importance of spatial structures in advancing hydrological sciences. *Water Resources Research* 42, W03S03, doi:10.1029/2005WR004301
- Schulze-Makuch D, Carlson DA, Cherkauer DS, Malik P. 1999. Scale dependency of hydraulic conductivity in heterogeneous media. *Ground Water* 37.6: 904-919, doi:10.1111/j.1745-6584.1999.tb01190.x
- Schumm SA, Licity RW, 1965. Time, space and causality in geomorphology. *American Journal of Science* 263: 110-119, doi:10.2475/ajs.263.2.110
- Schwilch G, Bernet L, Fleskens L, Giannakis E, Leventon J, Marañón T, Mills J, Short C, Stolte J, van Delden H, Verzandvoort S. 2016. Operationalizing ecosystem services for the mitigation of soil threats: A proposed framework. *Ecological Indicators* 67: 586-597, doi:10.1016/j.ecolind.2016.03.016
- Seppelt R, Müller F, Schröder B, Volk M. 2009. Challenges of simulating complex environmental systems at the landscape scale: A controversial dialogue between two cups of espresso. *Ecological Modelling* 220: 3481-3489, doi:10.1016/j.ecolmodel.2009.09.009
- Sornette D, von der Becke S. 2011. Complexity clouds finance-risk models. *Nature*, 471(7337): 166-166, doi:10.1038/471166a
- Stamati FE, Nikolaidis NP, Banwart S, Blum WE. 2013. A coupled carbon, aggregation, and structure turnover (CAST) model for topsoils. *Geoderma* 211: 51-64, doi:10.1016/j.geoderma.2013.06.014
- Stewart J, Parsons AJ, Wainwright J, Okin GS, Bestelmeyer BT, Fredrickson EL, Schlesinger WH. 2013. Modeling emergent patterns of dynamic desert ecosystems. *Ecological Monographs* 84: 373-410, doi:10.1890/12-1253.1
- Taylor PD, Fahrig L, Henein K, Merriam G. 1993. Connectivity is a vital element of landscape structure. *Oikos*: 571-573, doi:10.2307/3544927

- Tebebu TY, Steenhuis TS, Dagnew DC, Guzman CD, Bayabil HK, Zegeye AD, Collick AS, Langan S, MacAlister C, Langendoen EJ, Yitaferu B, Tilahun SA. 2015. Improving efficacy of landscape interventions in the (sub) humid Ethiopian highlands by improved understanding of runoff processes. *Frontiers in Earth Science* 3: 49, doi:10.3389/feart.2015.00049
- Temme AJAM, Veldkamp A. 2009. Multi-process Late Quaternary landscape evolution modelling reveals lags in climate response over small spatial scales. *ESPL* 34: 573-589, doi:10.1002/esp.1758
- Teuling AJ, Uijlenhoet R, Troch PA. 2005. On bimodality in warm season soil moisture observations. *Geophysical research letters* 32.13, doi:10.1029/2005GL023223
- Teuling AJ, Taylor CM, Meirink JF, Melsen LA, Miralles DG, van Heerwaarden CC, Vautard R, Stegehuis AI, Nabuurs G-J, Vilà-Guerau de Arellano J. 2017. Observational evidence for cloud cover enhancement over western European forests. *Nature Communications* 8: 14065, doi:10.1038/ncomms14065
- Tilman, David. 1994. Competition and biodiversity in spatially structured habitats." *Ecology* 75.1: 2-16, doi:10.2307/1939377
- Trewavas A. 2002. Plant Intelligence: Mindless mastery. *Nature* 415: 841, doi:10.1038/415841a
- Troch PA, Lahmers T, Meira A, Mukherjee R, Pedersen JW, Roy T, Valdés-Pineda R. 2015. Catchment coevolution: A useful framework for improving predictions of hydrological change? *Water Resources Research* 51: 4903-4922, doi:10.1002/2015WR017032
- Turnbull L, Wainwright J, Brazier RE. 2008. A conceptual framework for understanding semi-arid land degradation: ecohydrological interactions across multiple-space and time scales. *Ecohydrology* 1: 23-34, doi:10.1002/eco.4
- Van der Ploeg MJ, Appels WM, Cirkel DG, Oosterwoud MR, Witte JP, Van der Zee SEATM. 2012. Microtopography as a driving mechanism for ecohydrological processes in shallow groundwater systems. *Vadose Zone Journal* 11.3, doi:10.2136/vzj2011.0098
- van der Ploeg MJ, Teuling AJ. 2013. Going back to the roots: The need to link plant functional biology with vadose zone processes. *Procedia Environmental Sciences* 19: 379-383, doi:10.1016/j.proenv.2013.06.043
- Venter O, Sanderson EW, Magrath A, Allan JR, Beher J, Jones KR, Possingham HP, Laurance WF, Wood P, Fekete BM, Levy MA, Watson JEM. 2016. Sixteen years of change in the global terrestrial human footprint and implications for biodiversity conservation. *Nature Communications* 7,doi:10.1038/ncomms12558
- Vereecken H, Schnepf A, Hopmans JW, Javaux M, Or D, Roose T, ..., Young IM. 2016. Modeling Soil Processes: Review, Key Challenges, and New Perspectives. *Vadose Zone Journal* 15.5, doi:10.2136/vzj2015.09.0131
- Verwijmeren M, Rietkerk M, Bautista S, Mayor AG, Wassen MJ, Smit C. 2014. Drought and grazing combined: Contrasting shifts in plant interactions at species pair and community level. *Journal of Arid Environments* 111: 53-60, doi:10.1016/j.jaridenv.2014.08.001
- Vogel HJ. 2000. A numerical experiment on pore size, pore connectivity, water retention, permeability, and solute transport using network models. *European Journal of Soil Science* 51.1: 99-105, doi:10.1046/j.1365-2389.2000.00275.x
- Vogelmann JE, Gallanta AL, Shib H, Zhub Z. 2016. Perspectives on monitoring gradual change across the continuity of Landsat sensors using time-series data, *Remote Sensing of Environment* 185, 258-270, doi: 10.1016/j.rse.2016.02.060
- von Arx G, Archer SR, Hughes MK. 2012. Long-term functional plasticity in plant hydraulic architecture in response to supplemental moisture. *Annals of Botany* doi:10.1093/aob/mcs030
- Wainwright J, Turnbull L, Ibrahim TG, Lexartza-Artza I, Thornton SF, Brazier RE. 2011. Linking environmental regimes, space and time: Interpretations of structural and functional connectivity. *Geomorphology* 126: 387-404, doi:10.1016/j.geomorph.2010.07.027
- Wassen MJ, de Boer HJ, Fleischer K, Rebel KT, Dekker SC. 2013. Vegetation-mediated feedback in water, carbon, nitrogen and phosphorus cycles. *Landscape ecology* 28.4: 599-614, doi:10.1007/s10980-012-9843-z
- Wu J. 2006. Landscape ecology, cross-disciplinarity, and sustainability science. *Landscape Ecology* 21.1: 1-4, doi:10.1007/s10980-006-7195-2
- Zegeye AD, Langendoen EJ, Stoof CR, Tilahun SA, Dagnew DC, Zimale FA, Guzman CD, Yitaferu B, Steenhuis TS. 2016. Morphological dynamics of gully systems in the subhumid Ethiopian Highlands: the Debre Mawi watershed. *SOIL* 2: 443-458





Scale				
Challenge	How does microbial diversity depend on increased pore-scale tortuosity and heterogeneity?/ To what extent does small scale heterogeneity matter at larger scales?	How does the change in connectivity as we switch from piston flow to bypass flow alter the carbon and nutrient cycling in soils?	How do small scale interventions on the landscape alter hydrological flow paths and sediment transport?	How are hydrological, sediment and habitat function altered by major infrastructure?/ How do vegetation patterns combined at river basin scale influence water and sediment transport?
Image description and credits	Top: Schematic pore structure. Bottom: Iron oxidizing Leptothrix bacteria (Credit: Bertram Schmidt, CC, distributed via imagegeo.edu.eu).	Top: Cracks in clay soil (Credit: with kind permission from Bram te Brake). Bottom: Preferential flow patterns through soil (Credit: with kind permission from Esther Bloem)	Top: Farm fields in Exmoore, UK (Credit: Maria Burguet, CC, distributed via imagegeo.edu.eu). Bottom: Pivot irrigation (Credit: Photo by John A. Kelley, USDA Natural Resources Conservation Service via Flickr under Creative Commons licence).	Left: Landsat 8 imagery before (top) and after (bottom) flooding in Argentina. Right: Landsat 1, 5, 7 imagery of three decades of change in the birdsfoot delta of the Mississippi River (Data available from the U.S. Geological Survey.)

Figure 1: Illustration of challenges in biophysical landscape interactions where process understanding would benefit from bridging scales and disciplines.

Accept




Figure 2: Sketch representing a dryland banded vegetation system with two potential system states: a) intact banded system with relatively short runoff pathways and limited hydrological connectivity (dashed lines) and b) disturbed system with longer runoff pathways and increased hydrological connectivity (continuous line) in rills or gullies.