

1 **Population biology of grey gurnard (*Eutrigla gurnardus* L.; Triglidae) in**  
2 **the coastal waters of Northwest Wales**

3

4 **Running Title:** Population biology of grey gurnard

5

6 **I. D. McCarthy | J. Cant<sup>1</sup> | A. L. Marriott<sup>2</sup>**

7

8 School of Ocean Sciences, Bangor University, Menai Bridge, Anglesey, UK.

9

10 <sup>1</sup>Present address: Faculty of Biological Sciences, University of Leeds, Leeds, UK

11

12 <sup>2</sup>Present address: Inorganic Geochemistry, Centre for Environmental Geochemistry,  
13 British Geological Survey, Keyworth, Nottingham, UK

14

15

16 **Correspondence**

17 Ian D. McCarthy, School of Ocean Sciences, Bangor University, Menai Bridge,  
18 Anglesey, UK.

19 Email: i.mccarthy@bangor.ac.uk

20

21 **Summary**

22 The grey gurnard *Eutrigla gurnardus* (L.) has been identified by ICES as a potential  
23 commercial species in the NE Atlantic with recommendations made to derive information  
24 on population biology for stock assessment purposes. However, data on the population  
25 biology of this species is limited. In this study, data on the age, growth and maturity of  
26 grey gurnard were collected by otter trawling in the coastal waters of northwest Wales and  
27 Eastern Anglesey. Total length (TL) of fish sampled ranged between 2.1-33.0 cm (male)  
28 and 1.9-36.9 cm (female) with the majority of female (70.8%) fish between 11-20 cm TL  
29 and male fish (70.5 %) between 11-18 cm TL. The percentage of fish >20 cm TL was larger  
30 for females (30.4%) compared to males (17.6%). Total weight (TW) for female and male

31 grey gurnard in the stratified subsample ranged from 1.9-499.9 g for females and 2.1-390.0  
32 g for males, with the majority of female (66.3%) and male (76.1%) fish between 10 and 60  
33 g. TL/TW relations for male and female fish and both sexes combined were:  $TW =$   
34  $0.006TL^{3.07}$ ,  $TW = 0.007TL^{3.03}$  and  $TW = 0.007TL^{3.05}$  respectively. Age structure (based  
35 on otolith reading) ranged between 0.5 and 7.5 years old for females and 0.5 to 5.5 years  
36 old for male with the majority of female (41.7%) and male (45.95%) fish aged as 1.5 years  
37 old. The age structure of female and male grey gurnards was significantly different with  
38 the majority of older fish (> 2.5 years) being female. The von Bertalanffy growth functions  
39 were calculated as  $L_t = 32.4[1 - e^{-0.24(t+1.41)}]$  for males,  $L_t = 45.9[1 - e^{-0.16(t+1.37)}]$  for females  
40 and  $L_t = 44.0[1 - e^{-0.18(t+1.20)}]$  for both sexes combined. Instantaneous rates of total mortality  
41 were similar for males and females and the combined Z value  $1.00 \text{ year}^{-1}$  with the natural  
42 mortality rate estimated as  $0.33 \text{ year}^{-1}$ . The size at 50% maturity ( $L_{50}$ ) was estimated to be  
43 25.3 cm TL for males, females and for both sexes combined. Age at 50% maturity ( $A_{50}$ )  
44 was 3.2 years for both males and females. The results of this study provide the first  
45 information on the population biology of *E. gurnardus* in the Irish Sea, the first detailed  
46 study in the NE Atlantic since 1985 and helps to address the data gap identified by ICES  
47 in knowledge of the population biology of this species.

48

## 49 **1 | INTRODUCTION**

50 The grey gurnard *Eutrigla gurnardus* L. (Triglidae) is a demersal teleost fish found in the  
51 Eastern Atlantic (Norway to Morocco, Madeira, Iceland, including Greenland) and also in  
52 the Mediterranean and Black Sea (Froese & Pauly, 2017). Across its range it is commonly  
53 found on sandy substrates usually at depths between 10 to 140 m, although it is recorded  
54 as deep as 340 m in the E. Mediterranean (Froese & Pauly, 2017). Grey gurnard feed  
55 predominantly on crustaceans (mostly mysids and decapods) and fishes (especially gobies,  
56 flatfish, and young gadids) with the latter becoming more important as the fish increase in  
57 size (Moreno-Amich, 1994; Montanini, Stagioni & Vallisneri, 2010; Montanini, Stagioni,  
58 Benni & Vallisneri, 2017). *E. gurnardus* has been described as being a Lusitanian species  
59 (Yang, 1982) with historically a predominantly southern distribution in coastal waters  
60 around the UK but its distribution is shifting, moving North with climate change (Beare et  
61 al., 2004; Perry, Low, Ellis & Reynolds, 2005). It is ranked among the 10 most dominant

62 species in the North Sea, where its abundance has increased since the 1980s (Heessen &  
63 Daan, 1995; Floeter et al., 2005), possibly through occupying the ecological niche of  
64 demersal gadoids that had declined through overfishing (Floeter et al., 2005).

65 Historically the grey gurnard, together with other gurnard species, has formed a major  
66 part of discards in bottom trawl fisheries in the coastal shelf seas around the UK and Ireland  
67 (Borges, Rogan & Officer, 2005; Enever, Revill & Grant, 2007, 2009), although discard  
68 rates of the grey gurnard in some fisheries, e.g. the Dutch beam trawl fishery in the North  
69 Sea, have declined as new markets for grey gurnard have developed (Catchpole, van  
70 Keeken, Gray & Piet, 2008). With increasing pressure on demersal fish stocks in the north  
71 east Atlantic, ICES has recognized the grey gurnard, along with the red gurnard  
72 *Chelidonichthys cuculus* and tub gurnard *C. lucerna*, as new potential commercial species,  
73 ('MoU species'; ICES, 2006). As interest in all 3 gurnard species has increased, ICES has  
74 recommended that landings and discards are monitored and information regarding  
75 population biology is obtained for stock assessment purposes, however, information  
76 remains limited for all three species (ICES, 2010, 2013, 2014a, 2015, 2016), especially  
77 biological data for the grey gurnard (ICES, 2010). A knowledge of ecology and population  
78 biology is essential in the development of sustainable management plans for any exploited  
79 species (King, 2007) but to the authors' knowledge, there has been no study of the  
80 population biology (i.e. patterns of growth, mortality and reproduction) of grey gurnard in  
81 the Irish Sea, and no detailed study on the population biology of grey gurnard since Baron  
82 (1985a, 1985b) in the Bay of Douarnenez, France. Therefore, the aims of this study are to  
83 provide data on the population biology (specifically the size/age-structure and patterns of  
84 growth, maturity and mortality) of grey gurnard *Eutrigla gurnardus* within the coastal  
85 waters of Eastern Anglesey and Northwest Wales.

86

## 87 **2 | MATERIALS AND METHODS**

### 88 **2.1 | Sample collection**

89 Grey gurnards were collected between 2000 and 2015 (excluding 2005 and 2008) by the  
90 *RV Prince Madog* in the coastal waters around Eastern Anglesey and Northwest Wales  
91 (Figure 1) as part of the ongoing survey (since 1972) of local demersal fish stocks. Surveys  
92 were conducted during October of each year in the same five areas: (A) Red Wharf Bay,

93 (B) Conwy Bay, (C) Inshore Colwyn Bay, (D) Offshore Point Lynas and (E) Offshore  
94 Colwyn Bay (A-C designated as 'inshore' sites and D-E designated as 'offshore' site; see  
95 Marriott, Latchford & McCarthy, 2010). During the 14 years of sampling, 275 trawls were  
96 conducted at trawl depths between 10 and 32 m in inshore sites (205 trawls in total) and  
97 between 17 and 45 m in offshore sites (70 trawls in total) respectively. The five sampling  
98 areas have similar substrate with most sites comprising of gravelly sand, medium sand and  
99 broken shells, with the sites around Point Lynas and Red Wharf Bay comprising mainly of  
100 sandy gravel and sand respectively. Trawls of approximately 1 hour duration towed at 2-3  
101 knots were conducted using a rockhopper otter trawl (cod end stretched mesh size of 73  
102 mm), in the five survey areas. On completion of each trawl, the catch was sorted and all  
103 grey gurnard were retained and frozen. In the laboratory, the fish were defrosted overnight  
104 and the following data were collected: TL (to nearest 0.1 mm), total weight (TW, to nearest  
105 0.1 g), sex and maturity status (immature or mature based on macroscopic examination of  
106 the gonads; Booth, 1997; King, 2007). Finally, the sagittal otoliths were removed and  
107 stored in paper envelopes until subsequent ageing. The age of each fish was determined as  
108 described by Marriott et al. (2010) using digital imaging techniques assuming one pair of  
109 opaque/hyaline bands formed each year (Colloca, Cardinale, Marcello & Ardizzone, 2003).

110

## 111 **2.2 | Data analysis**

112 The length-weight relationship was described using the power function  $TW = aTL^b$  (King,  
113 2007), where  $a$  and  $b$  are constants, with data for females and males examined separately  
114 and the slopes of the regression lines for the log-transformed data compared using a GLM  
115 to test for differences between the sexes. The  $b$ -values for males and females were also  
116 tested against a value of  $b=3$  to test for isometric growth. The age structure of male and  
117 female fishes was compared using a chi-squared test. The relationship between mean length  
118 at age for male and female grey gurnard was described using the von Bertalanffy growth  
119 (VBG),  $L_t = L_\infty[1 - e^{-k(t - t_0)}]$  (King, 2007), where  $L_t$  is the average TL (cm) at age  $t$  (years),  $k$   
120 is the growth coefficient ( $\text{year}^{-1}$ ),  $L_\infty$  is the asymptotic total length and  $t_0$  is the theoretical  
121 age at length zero (year). The total instantaneous mortality rates ( $Z$ ,  $\text{year}^{-1}$ ) were calculated  
122 from the linearised catch curve data (King, 2007) for females and males and the slopes of  
123 the regression lines for the log-transformed data were compared using a GLM to test for

124 differences between the sexes. The instantaneous rates of natural mortality ( $M$ , year<sup>-1</sup>) were  
125 estimated using the Pauly (1980) equation based on growth in length and the average  
126 surface seawater temperature for the area (10.66°C; Moelfre and Amlwch stations; Joyce,  
127 2006). The exploitation ratio,  $E = F/Z$  (King, 2007), was calculated for males and females  
128 combined where  $F$  (year<sup>-1</sup>) is the instantaneous rate of fishing mortality (estimated as  $Z -$   
129  $M$ ). The TL ( $L_{50}$ , cm) and age ( $A_{50}$ , years) at 50% maturity were calculated using the logistic  
130 equation  $Y = 1/[1 + e^{-r(L - X_{50})}]$  (King, 2007), where  $Y$  is the proportion of mature fish in the  
131 total length class  $L$  (cm),  $r$  is a constant and  $X_{50}$  is the TL or age at 50% maturity. In order  
132 to determine whether population biology parameters had changed over time, the correlation  
133 between year and length-weight coefficients, average/maximum size, or  
134 minimum/maximum age were determined using Pearson's correlation coefficient. Data  
135 were grouped into three time periods: 2000-2004, 2006-2010 and 2011-2015 and  
136 differences in size at age in each age group examined using ANOVA or t-test and b-values  
137 of the log-transformed TL/TW relations examined using GLM. All data are presented as  
138 mean values  $\pm$  SD with statistical analyses conducted in SPSS v22.

139

### 140 **3 | RESULTS**

141 In total, 1268 grey gurnard (732 females and 536 males) were caught with males and  
142 females ranging in TL between 2.1-33.0 cm and 1.9-36.9 cm (Figure 2A) respectively. The  
143 majority of female (70.8%) were between 11-20 cm TL and the majority of male fish (70.5  
144 %) were between 11-18 cm TL (Figure 2A). The percentage of fish >20 cm TL was larger  
145 for females (30.4%) compared to males (17.6%). TW for female and male grey gurnard in  
146 the stratified subsample ranged from 1.9-499.9 g for females and 2.1-390.0 g for males,  
147 with the majority of female (66.3%) and male (76.1%) fish between 10 and 60 g.

148 A total of 1021 fish could be aged (614 female, 407 male) with the age structure ranging  
149 between 0.5 and 7.5 years old for females and 0.5 to 5.5 years old for male fish (Figure  
150 2B). For both females (41.7%) and males (45.9%), the majority of fish were 1.5 years old.  
151 The age structures of female and male grey gurnards in the subsample were significantly  
152 different ( $\chi^2_4=34.8$ ,  $P<0.001$ ), with the older age classes consisting predominantly of  
153 female fish (Figure 2B).

154 The length-weight relationships for female and male grey gurnard and for both sexes  
155 combined are presented in Figure 3. Males exhibited positive allometric growth with a *b*  
156 value significantly different from 3 ( $\delta$ ,  $b=3.077$  [ $SE_b=0.033$ ];  $t_{534}=2.33$ ,  $P=0.02$ ) whereas  
157 females exhibited isometric growth ( $\text{♀}$ ,  $b=3.030$  [ $SE_b=0.021$ ];  $t_{731}=1.43$ ,  $P=0.15$ ). The  
158 slope values for the log-transformed linearised length-weight data for both female and male  
159 grey gurnard were similar ( $F_{1,1266}=1.33$ ,  $P=0.25$ ). The length-weight relationship for the  
160 combined data was described by  $TW = 0.007TL^{3.05}$  ( $SE_b=0.015$ ,  $r^2=0.970$ ,  $P<0.001$ ), with  
161 the *b* value significantly different from 3 ( $t_{1269}=3.0$ ,  $P<0.001$ ). VBG curves for female and  
162 male grey gurnard are presented in Figure 4 with the growth parameters presented in Table  
163 1. Female grey gurnard attained a larger  $L_\infty$  value than males ( $\text{♀}=45.9$  cm;  $\delta=32.4$  cm).  
164 The growth curve for the combined male and female data is described by  $L_t = 44.0[1 - e^{-0.18(t$   
165  $+ 1.20)}]$  ( $r^2=0.948$ ,  $P<0.001$ ).

166 There were no differences in the instantaneous rates of total mortality for males and  
167 females ( $\text{♀}=0.94$  year<sup>-1</sup>,  $\delta=1.13$  year<sup>-1</sup>;  $F_{1,6}=4.81$ ,  $p = 0.07$ ) and the instantaneous rate of  
168 total mortality for males and females combined was  $Z=1.00$  year<sup>-1</sup> with the instantaneous  
169 rates of natural mortality estimated as  $M=0.33$  year<sup>-1</sup> and the exploitation ratio as  $E=0.67$ .  
170 Length maturity ogives for male and female grey gurnard are presented in Figure 5. The  
171 calculated  $L_{50}$  values for female, male and combined sexes were calculated as 25.34 cm  
172 ( $\text{♀}$ ), 25.28 cm ( $\delta$ ) and 25.31 cm and the  $A_{50}$  values for female and male grey gurnard were  
173 both calculated as 3.2 years respectively.

174 When the population biology data were examined to determine any temporal changes  
175 There was no indication of changes in length-weight coefficients over time (Figure 6A)  
176 and the slope values for the log-transformed linearised length-weight data for the 5-year  
177 groupings were similar ( $F_{1,1266}=0.19$ ,  $P=0.83$ ; Table 2). Decreases in average length  
178 ( $r=0.61$ ,  $p=0.02$ ), maximum length ( $r=0.47$ ,  $p=0.09$ ) (Figure 2B) and maximum age  
179 (Figure 2C) were observed over time but these decreases coincided with an increase in the  
180 proportion of inshore fishing conducted in any given year ( $r=0.54$ ,  $p=0.04$ ) (Figure 2D).  
181 VBG curves could not be fitted to the size at age data for each 5 year grouping, however  
182 the average size for each time group were similar for each age group (0.5-2.5 years,  
183 ANOVAs, all  $p=0.13-0.56$ ; 3.5-4.5 years, *t*-tests,  $p=0.98$  and  $p=0.23$  respectively)  
184 suggesting that patterns of growth were similar over time (Figure 6E). Patterns of maturity

185 appeared similar over time (Figure 6F) with calculated  $L_{50}$  values for 2000-2004 and 2011-  
186 2015 of 25.00 and 25.81 cm respectively (Table 2).

187

#### 188 **4 | DISCUSSION**

189 ICES has identified the three main gurnard species in the NE Atlantic, red gurnard *C.*  
190 *cuculus*, tub gurnard, *C. lucerna* and grey gurnard *E. gurnardus* as potential new species  
191 for commercial exploitation (ICES 2006, 2013). However, detailed information on the  
192 population biology and landings/discard data for stock assessment purposes for each  
193 species within the different ICES subareas of the NE Atlantic is currently lacking. To help  
194 address this data gap, data from surveys conducted on all three triglid species in northwest  
195 Wales have been recently published: red gurnard (Marriott et al., 2010), tub gurnard  
196 (McCarthy & Marriott, 2017) and grey gurnard (this study). Population biology data for  
197 grey gurnard is limited. Although there have been many published studies presenting  
198 length-weight relationships for the species in the NE Atlantic and the Mediterranean (see  
199 Table 3), however, the number of published studies where multiple biological parameters,  
200 e.g. growth and reproduction, are co-reported is limited to studies in Brittany and the Faroe  
201 Islands (Table 1).

202 The maximum sizes reported in the literature for grey gurnard range from 50-60 cm  
203 depending on the population, although these values are for studies published between 1913  
204 and 1969 (data from Algeria, English Channel, North Sea and Mediterranean reported in  
205 Table 2 of Baron, 1985a). More recent studies report maximum sizes of *ca.* 40-46 cm  
206 [Brittany, Baron (1985a); North Sea and Skaggeiak, ICES (2010, 2014c); Bay of Biscay,  
207 ICES (2014d); Celtic Sea, ICES (2014b)]. In the present study, grey gurnard in the inshore  
208 coastal waters of NW Wales attained a maximum size of 37 cm TL and, since sampling  
209 was conducted at <45 m water depth, it is possible that larger fish may be located further  
210 offshore in deeper water. However, beam trawl surveys conducted at multiple sites within  
211 the Irish Sea report similar TL size-frequency distributions to that observed in the present  
212 study with a maximum size of 37 cm TL in Parker-Humphreys (2004) and 34 cm TL in  
213 ICES (2014b) indicating that the full size range for the species in the Irish was sampled  
214 within the study area. In addition, Parker-Humphreys (2004) presents data on the  
215 distribution and abundance of grey gurnards at 66 sites within the Irish Sea indicating that

216 in the eastern Irish Sea grey gurnards are caught at depths < 40 m and so the full depth  
217 range over which fish are likely to be found has also been sampled in the present study.

218 Positive allometric growth has been reported in the majority of grey gurnard studies  
219 (Tables 1 and 3) and average 'a' and 'b' values for the length-weight relationship from  
220 these studies are 0.007 and 3.06 respectively. The 'a' and 'b' values obtained for grey  
221 gurnard in northwest Wales are similar to this average (Table 1). The maximum age  
222 reported in the present study (8 years) is the same as that reported by Damm (1987) for  
223 grey gurnard in the North Sea, with a maximum age of 16 years reported by Baron (1985a)  
224 for grey gurnard in the Bay of Douarnenez, Brittany. The available data indicates that  
225 females grow faster than males and that the majority of larger individuals in the population  
226 are female (Baron 1985a; Damm, 1987; present study). Variability in size at age increased  
227 with increasing age class for both sexes (Figure 4) and it is possible that this could be due  
228 to dietary specialisation by some grey gurnards. An ontogenetic diet switch has been  
229 reported for the species with a change in dominant prey taxa from crustaceans (mostly  
230 mysids and decapods) to fishes (especially gobies, flatfish, and young gadids) as the fish  
231 increase in size (Moreno-Amich, 1994; Montanini et al., 2010). However, there is evidence  
232 to suggest individual dietary specialisation in grey gurnard with three putative feeding  
233 types (Weinert, Floeter, Krönke and Sell, 2010): a predator specialised on fish (FP), on  
234 invertebrates (IP) or having a mixed diet (MDP). Recent work by Montanini et al. (2017)  
235 also indicates increasing individual dietary specialisation in grey gurnard with increasing  
236 size. Weinert et al. (2010) found that fish condition, in terms of the length-specific  
237 individual weight, increased with specialisation on fish prey (FP > MDP > IP) and,  
238 although size at age and growth rates were not considered in their study, it is possible that  
239 the increased energy intake observed with specialisation on fish may translate into  
240 differences in growth (FP > MDP > IP) that may explain the larger size range observed  
241 with increasing age.

242 Although the population biology presented in this study was collected over a 16 year  
243 time period (2000-2015), there is little evidence for changes in length-weight relations and  
244 patterns of growth and maturity over time (Figure 6, Table 2) with changes in the size and  
245 age distributions over time related to an increase in inshore fishing in recent years. The  
246 population biology data for grey gurnards are limited (Table 1), however, the VBG



247 parameters ( $k$  and  $L_{\infty}$ ) for northwest Wales are similar to those reported for the North Sea.  
248 Growth performance of grey gurnards from different populations can be compared using  
249 the phi prime growth performance index ( $\Phi' = 2\log_{10}L_{\infty} + \log_{10}k$ ; Pauly and Munro, 1984)  
250 and the data are presented in Table 1. The average  $\Phi'$  value for the three grey gurnard  
251 populations from the NE Atlantic studied is  $2.84 \pm 0.24$  (range 2.53 - 2.96). The value  
252 recorded in the present study (2.53) is at the lower end of the  $\Phi'$  values reported but within  
253 the range of published values.

254 For an accurate assessment of  $L_{50}$ , it is recommended that sampling should be conducted  
255 at the start of the reproductive season (Lowerre-Barbieri et al., 2011). Although this was  
256 not the case in the present study as the spawning season for grey gurnard is thought to be  
257 between February-August (Froese & Pauly, 2017), the  $L_{50}$  values obtained in the present  
258 study were very similar to those obtained by most other studies in the NE Atlantic (Table  
259 1. In contrast, Valisineri, Montanini & Stagoni (2012) observed a much smaller length at  
260 50% maturity, 12 or 15 cm TL (♀ or ♂) for the single published study from the  
261 Mediterranean with Montanini et al. (2017) stating that grey gurnard may start to mature  
262 as small as 10 cm TL in the Mediterranean.

263 With the growing interest in gurnard species as MoU species, ICES has recommended  
264 that landings and discards are monitored and information on population biology is obtained  
265 for stock assessment purposes, however, information remains limited for red, tub and grey  
266 gurnards (ICES, 2010, 2013, 2015, 2016). Previously, gurnard landings were not sorted by  
267 species and were often reported as the generic category 'gurnards' and thus, species-  
268 specific data are only available from countries participating in gurnard fisheries since 2010  
269 (ICES, 2015). The issue of accurately quantifying discard rates for each gurnard species in  
270 other demersal fisheries still remains unresolved although discard rates are thought to be  
271 very high (ICES, 2015, 2016). For example, the average discard rate for grey gurnards in  
272 the North Sea is estimated at 80% (ICES, 2016). As a result, the management advice  
273 provided for grey gurnard is limited and advises a precautionary approach with reduced  
274 landings until more detailed information on population biology, stock size, fishing pressure  
275 and discard rates are determined as these are currently unknown (ICES 2014b, 2014c,  
276 2014d, 2016). The results of this study provide the first information on the population  
277 biology of *E. gurnardus* in the Irish Sea, the first detailed study in the NE Atlantic since

278 1985 and helps to address the data gap identified by ICES in knowledge of the population  
279 biology of this species.

280

## 281 **ACKNOWLEDGEMENTS**

282 The authors wish to thank the crew of the RV *Prince Madog* and the staff and students  
283 from the School of Ocean Sciences who conducted the fisheries surveys and collected  
284 much of the raw data used in this study.

285

## 286 **REFERENCES**

- 287 Baron, J. (1985a). Le Triglides (Teleosteen Scorpaeniformes) de la Baie de Douarnenez. I. La  
288 croissance de: *Eutrigla gurnardus*, *Trigla lucerna*, *Trigloporus lastoviza* et *Aspitrigla cuculus*.  
289 *Cybium*, 9, 127-144.
- 290 Baron, J. (1985b). Le Triglides (Teleosteen Scorpaeniformes) de la Baie de Douarnenez. II. La  
291 reproduction de *Eutrigla gurnardus*, *Trigla lucerna*, *Trigloporus lastoviza* et *Aspitrigla*  
292 *cuculus*. *Cybium* 9, 252-281.
- 293 Beare, D.J., Burns, F., Greig, A., Jones, E.G., Peach, K., Kienzle, M., McKenzie, E., & Reid, D.G.  
294 (2004). Long-term increases in prevalence of North Sea fishes having southern biogeographic  
295 affinities. *Marine Ecology Progress Series*, 284, 269–278.
- 296 Bilge, G., Yapici, S., Filiz, H., & Cerim, H. (2014). Weight–length relations for 103 fish species  
297 from the southern Aegean Sea, Turkey. *Acta Ichthyologica et Piscatoria*, 44, 263-269.
- 298 Bok, T.D., Gokturk, S.D., Kahraman, A.E., Alicli, T.Z., Acun, T., & Ates, C. (2011). Length-  
299 weight relationships of 34 fish species from the Sea of Marmara, Turkey. *Journal of Animal*  
300 *and Veterinary Advances* 10, 3037-3042. DOI: 10.3923/javaa.2011.3037.3042
- 301 Booth, A.J. (1997). On the life history of the lesser gurnard (Scorpaeniformes: Triglididae) inhabiting  
302 the Agulhas Bank, South Africa. *Journal of Fish Biology*, 51, 1155-1173. DOI:  
303 10.1111/j.1095-8649.1997.tb01133.x
- 304 Borges, L., Rogan, E., & Officer, R. (2005). Discarding by the demersal fishery in the waters  
305 around Ireland. *Fisheries Research* 76, 1-13. DOI: 10.1016/j.fishres.2005.05.011
- 306 Catchpole, T.L, Van Keeken O, Gray T.S., Piet G. (2008). The discard problem - A comparative  
307 analysis of two fisheries: The English *Nephrops* fishery and the Dutch beam trawl fishery.  
308 *Ocean & Coastal Management*, 51, 772-778. DOI: 10.1016/j.ocecoaman.2008.06.015

309 Colloca, F., Cardinale, M., Marcello, A., & Ardizzone, G.D. (2003). Tracing the life history of red  
310 gurnard (*Aspitrigla cuculus*) using validated otolith annual rings. *Journal of Applied*  
311 *Ichthyology*, 19, 19, 1–9. DOI: 10.1046/j.1439-0426.2003.00342.x

312 Damm, U. (1987). Growth of the grey gurnard *Eutrigla gurnardus* L. in the North Sea. *ICES*  
313 *Committee Meeting G:55*. 10 p.

314 Enever, R., Revill, A. & Grant, A. (2007). Discarding in the English Channel, Western approaches,  
315 Celtic and Irish seas (ICES subarea VII). *Fisheries Research* 86, 143-152. DOI:  
316 10.1016/j.fishres.2007.05.013

317 Enever, R., Revill, A. & Grant, A. (2009). Discarding in the North Sea and on the historical efficacy  
318 of gear-based technical measures in reducing discards. *Fisheries Research* 95, 40-46. DOI:  
319 10.1016/j.fishres.2008.07.008

320 Floeter, J., Kempf, A., Vinther, M., Schrum, C. & Temming, A. (2005). Grey gurnard (*Eutrigla*  
321 *gurnardus*) in the North Sea: an emerging key predator? *Canadian Journal of Fisheries and*  
322 *Aquatic Sciences* 62, 1863-1864. DOI: 10.1139/F05-108

323 Froese, R. & Pauly, D. (Eds.), (2017). *FishBase*. World Wide Web electronic publication.  
324 Retrieved from [www.fishbase.org](http://www.fishbase.org), version (10/2017).

325 Heessen, H.J.L., & Daan, N. (1995). Long-term trends in ten non-target North Sea fish species.  
326 *ICES Journal of Marine Science* 53, 1063-1078. DOI: 10.1006/jmsc.1996.0133

327 ICES, (2006). Report of the working group on the assessment of new MOU species (WGNEW),  
328 13-15 December 2005, ICES Headquarters, *ICES Advisory Committee on Fisheries*  
329 *Management, CM2006/ACFM, 11*, 234 pp.

330 ICES, (2010). Report of the Working Group on Assessment of New MoU Species (WGNEW), 11-  
331 15 October 2010, ICES HQ, Denmark. *ICES CM 2010/ACOM: 21*, 185 pp.

332 ICES, (2013). Report of the Working Group on Assessment of New MoU Species (WGNEW), 18  
333 - 22 March 2013, ICES HQ, Copenhagen, Denmark. ACOM., 189 pp.

334 ICES, (2014a). Report of the Working Group on Assessment of New MoU Species (WGNEW),  
335 24–28 March 2014, Copenhagen, Denmark. *ICES CM 2014/ACOM:21*. 162 pp.

336 ICES, (2014b). Grey gurnard in Subarea VI and Divisions VIIa–c and e–k (Celtic Sea and West of  
337 Scotland). In Report of the ICES Advisory Committee, 2014. *ICES Advice 2014*, Book 5,  
338 Section 5.3.7.

339 ICES, (2014c). Grey gurnard in Subarea IV (North Sea) and Divisions VIIId (Eastern Channel) and  
340 IIIa (Skagerrak–Kattegat). In Report of the ICES Advisory Committee, 2014. *ICES Advice*  
341 *2014*, Book 6, Section 6.3.6.

342 ICES, (2014d). Grey gurnard in Subarea VIII and Division IXa (Bay of Biscay and Atlantic Iberian  
343 waters). In Report of the ICES Advisory Committee, 2014. ICES Advice 2014, Book 7, Section  
344 7.3.6.

345 ICES, (2015). Red gurnard (*Chelidonichthys cuculus*) in Subareas III, IV, V, VI, VII and VIII  
346 (Northeast Atlantic). In ICES advice on fishing opportunities, catch and effort. Celtic Seas,  
347 Greater North Sea, and the Bay of Biscay and the Iberian Coast Ecoregions. ICES Advice,  
348 2015, Book 9, section 9.3.33.

349 ICES, (2016). Grey gurnard (*Eutrigla gurnardus*) in Subarea 4 and divisions 7.d and 3.a (North  
350 Sea, eastern English Channel, Skagerak and Kattegat). In ICES advice on fishing opportunities,  
351 catch and effort. *ICES Advice, 2016, Book 6*, section 6.3.15.

352 Joyce, A.E. (2006). The coastal temperature network and ferry route programme: long-term  
353 temperature and salinity observations. *Science Series Data Report, Cefas Lowestoft, 43*, 129  
354 pp.

355 King, M. (2007). *Fisheries Biology, Assessment and Management*. 2<sup>nd</sup> Edition. Oxford, Blackwell  
356 Publishing, 382 pp.

357 Lowerre-Barbieri, S.K., Ganas, K., Saborido-Rey, F., Murua, H., & Hunter, J.R. (2011).  
358 Reproductive timing in marine fishes: variability, temporal scales, and methods. *Marine and*  
359 *Coastal Fisheries*, 3, 71–91. DOI: 10.1080/19425120.2011.556932

360 Magnussen, E. (2007). Interpopulation comparison of growth patterns of 14 fish species on Faroe  
361 bank: are all fishes on the bank fast growing?. *Journal of Fish Biology*, 71, 453-475. DOI:  
362 10.1111/j.1095-8649.2007.01502.x

363 Marriott, A.L., Latchford, J.W., & McCarthy, I.D. (2010). Population Biology of the red gurnard  
364 (*Aspitrigla cuculus* L.; Triglidae) in the inshore waters of Eastern Anglesey and Northwest  
365 Wales. *Journal of Applied Ichthyology*, 26, 504-512. DOI: 10.1111/j.1439-0426.2010.01455.x

366 McCarthy, I.D. & Marriott, A.L. (2018). Age, growth and maturity of tub gurnard (*Chelidonichthys*  
367 *lucerna* L.; Triglidae) in the inshore coastal waters of Eastern Anglesey and Northwest Wales.  
368 *Journal of Applied Ichthyology* in press. DOI: 10.1111/jai.13614

369 Montanini, S., Stagioni, M., & Vallisneri, M. (2010). Diet of the grey gurnard, *Eutrigla gurnardus*  
370 in the Adriatic Sea, north-eastern Mediterranean. *Cybium*, 34, 367-372.

371 Montanini, S., Stagioni, M., Benni, E. & Vallisneri, M. (2017). Feeding strategy and ontogenetic  
372 changes in diet of gurnards (Teleostea: Scorpaeniformes: Triglidae) from the Adriatic Sea. *The*  
373 *European Zoological Journal*, 84, 356-367, DOI: 10.1080/24750263.2017.1335357

374 Moreno-Amich, R. (1994). Feeding-habits of gray gurnard, *Eutrigla gurnardus* (L., 1758), along  
375 the Catalan coast (Northwestern Mediterranean). *Hydrobiologia*, 273, 57-66. DOI:  
376 10.1007/BF00126768

377 Parker-Humphreys, M. (2004). Distribution and relative abundance of demersal fishes from beam  
378 trawl surveys in the Irish Sea (ICES Division VIIa) 1993-2001. *Science Series Technical*  
379 *Report, CEFAS Lowestoft*, 120, 68pp.

380 Pauly, D. (1980). On the interrelationship between natural mortality, growth parameters and mean  
381 environmental temperature in 175 fish stocks. *Journal du Conseil permanent International*  
382 *pour l' Exploration de la Mer*, 39, 195-212.

383 Pauly, D., & Munro, J.L. (1984). Once more on the comparison of growth in fish and invertebrates.  
384 *Fishbyte*, 2, 21.

385 Perry, A.L., Low, P.J., Ellis, J.R., & Reynolds, J.D. (2005). Climate change and distribution shifts  
386 in marine fishes. *Science*, 308, 1912-1915. DOI: 10.1126/science.1111322

387 Uçkun, D. (2005). Investigation of the age and growth characteristics of the species belonging to  
388 the family Triglidae in Edremit Bay. *Ege University Journal of Fisheries & Aquatic Sciences*,  
389 22, 363–369.

390 Vallisneri, K., Montanini, S., & Stagoni, M. (2012). Size at maturity of triglid fishes in the Adriatic  
391 Sea, northeastern Mediterranean. *Journal of Applied Ichthyology*, 28, 123-125. DOI:  
392 10.1111/j.1439-0426.2011.01777.x

393 Weinert, M., Floeter, J., Kröncke, I., & Sell, A.F. (2010). The role of prey composition for the  
394 condition of grey gurnard (*Eutrigla gurnardus*). *Journal of Applied Ichthyology*, 24, 75-84.  
395 DOI: 10.1111/j.1439-0426.2010.01448.x

396 Yang, J. (1982). The dominant fish fauna in the North Sea and its determination. *Journal of Fish*  
397 *Biology*, 20, 635–643. DOI: 10.1111/j.1095-8649.1982.tb03973.x

398

399

400 **FIGURE LEGENDS**

401

402 **FIGURE 1** Location of sampling sites trawled for grey gurnard *E. gurnardus* (L.) in the coastal  
403 waters of Eastern Anglesey and Northwest Wales. Inshore sites: (A) Red Wharf Bay; (B) Conwy  
404 Bay; (C) Colwyn Bay; Offshore sites: (D) Colwyn Bay (North of the Constable Bank); (E) Offshore  
405 Point Lynas.

406 **FIGURE 2** (A) Length-frequency distributions (Total Length, cm) and (B) Age-frequency  
407 distribution for male and female grey gurnard *E. gurnardus* (L.) sampled in October in the coastal  
408 waters of Northwest Wales, UK, between 2000 and 2015 (excluding 2005 and 2008).

409 **FIGURE 3** Length-weight relationships for (A) female and (B) male grey gurnard *E. gurnardus*  
410 (L.) sampled in October in the coastal waters of Northwest Wales, UK, between 2000 and 2015  
411 (excluding 2005 and 2008).

412 **FIGURE 4** Length-at-age relationships for female (solid circles) and male (open circles) grey  
413 gurnard *E. gurnardus* (L.) sampled in October in the coastal waters of Northwest Wales, UK,  
414 between 2000 and 2015 (excluding 2005 and 2008).

415 **FIGURE 5** Maturity ogives (Total length at 50% maturity,  $L_{50}$ ) for (A) female and (B) male  
416 grey gurnard *E. gurnardus* (L.) sampled in October in the coastal waters of Northwest Wales, UK,  
417 between 2000 and 2015 (excluding 2005 and 2008).

418 **FIGURE 6** Temporal changes in population biology data for grey gurnard *E. gurnardus* (L.)  
419 (both sexes combined) sampled in October in the coastal waters of Northwest Wales, UK, between  
420 2000 and 2015 (excluding 2005 and 2008). Data are presented for (A) the a/b-values for the length-  
421 weight relationship, (B) mean and maximum Total length, (C) maximum and minimum age. (D)  
422 proportion of fishing tows conducted inshore, (E) length-at-age relationships and (F) maturity  
423 ogives (Total length at 50% maturity,  $L_{50}$ ).

424 **TABLE 1.** Summary of studies where multiple population biology measures have been studied for grey gurnard *Eutrigla gurnardus*.  
 425 Data are presented for the coefficients from the length-weight relationship (a, b), the von Bertalanffy growth function [ $L_{\infty}$  (cm), k (year<sup>-1</sup>),  $t_0$  (years)], the growth performance index  $\Phi'$  (Pauly & Munro, 1984) and length at 50% maturity ( $L_{50}$ ). All length values are Total  
 426 Length.  
 427  
 428

Region/Location (latitude)	Sex	a	b	$L_{\infty}$	k	$t_0$	$\Phi'$	$L_{50}$	Reference
North West Wales, UK	♂	0.006	3.07	32.4	0.24	-1.41	2.40	25.3	This Study
	♀	0.007	3.03	45.9	0.16	-1.37	2.52	25.3	
Brittany, France		0.007	3.05	44.0	0.18	-1.20	2.53	25.3	Baron (1985a & b) and Froese & Pauly (2017)
	♂	0.006	3.08	34.4	0.77	0.14	2.96	29.4	
	♀	0.005	3.17	38.0	0.80	0.16	3.09	31.2	
North Sea	♂+♀	0.005	3.11	-	-	-	-	-	Froese & Pauly (2017)
	♂	-	-	-	-	-	-	18.0	
	♀	-	-	-	-	-	-	24.0	
Gulf of Gascony, France	♂+♀	-	-	46.0	0.16	-	2.53	23.0	Damm (1987)
	♂+♀	0.005	3.13	-	-	-	-	20	
English Channel	♂+♀	0.005	3.19	-	-	-	-	23	Froese & Pauly (2017)
Faroe Bank, Faroe Islands	♂+♀	-	-	35.0	0.48	-	2.76	31.0	Magnussen (2007)

429

430 **TABLE 2.** Summary the population biology data for grey gurnard *Eutrigla gurnardus*  
 431 sampled in October in the coastal waters of Northwest Wales, UK, between 2000 and 2015  
 432 (excluding 2005 and 2008). Data are presented for the coefficients from the length-weight  
 433 relationship (a, b), average and maximum Total Length (cm), maximum and minimum ages  
 434 (years) and length at 50% maturity (L<sub>50</sub>). The proportion of inshore tows conducted in each  
 435 time period is also presented with the total number of tows in parentheses (see Figure 1 and  
 436 Methods text for definition of ‘inshore’ versus ‘offshore’).  
 437  
 438

Parameter	2000-2004	2006-2010	2011-2015
Sample size	352	179	737
a	0.008 (0.099)	0.007 (0.099)	0.007 (0.095)
b	3.00 (0.033)	3.04 (0.034)	3.01 (0.035)
Average TL	20.7 (5.2)	14.3 (5.0)	15.8 (3.9)
Maximum TL	36.9	32.9	32.4
Minimum age	0.5	0.5	0.5
Maximum age	7.5	2.5	4.5
L <sub>50</sub>	25.0	-	25.87
Proportion inshore	0.66 (n=66)	0.69 (n=101)	0.81 (n=118)

439  
 440  
 441  
 442  
 443  
 444  
 445


**TABLE 3.** Summary of studies presenting data on the Total Length-Total Weight relationship for grey gurnard *Eutrigla gurnardus*. Data are presented for males and females separately or for both sexes combined. Data are taken from Fishbase (Froese & Pauly, 2017) except where indicated by asterisks.

Location	♂ a	♂ b	♀ a	♀ b	♂+♀ a	♂+♀ b
Scotland (N. Sea & West Coast)	-	-	-	-	0.006	3.10
North Sea	-	-	-	-	0.011	2.88
Portugal (S.coast)	0.024	2.72	0.017	2.86	-	-
Pagassitikos Gulf, Greece	0.006	3.08	0.005	3.17	-	-
Trikeri Channel, Greece	-	-	-	-	0.004	3.08
South Aegean Sea, Turkey*					0.004	3.26
Edremit Bay, Turkey**	0.007	3.08	0.006	3.17	-	-
Sea Of Marmara, Turkey***	-	-	-	-	0.011	2.96

446  
 447

\* Bilge et al. (2014), \*\* Uçkun (2005), \*\*\* Bok et al. (2011)







448  
449  
450  
451


Figure 1

452




453  
454  
455

Figure 2





456  
457  
458  
459

Figure 3




460  
461  
462  
463

Figure 4



464  
465  
466  
467

Figure 5



468  
469  
470

Figure 6