



British Geological Survey

NATURAL ENVIRONMENT RESEARCH COUNCIL

1. Introduction

Carbon Capture and Storage (CCS) is crucial for low-carbon industry, climate mitigation and a sustainable energy future. The offshore capacity of the UK is substantial and has been estimated at 78 Gt of CO₂ in saline aquifers and hydrocarbon fields^[1]. The early-mid Eocene Tay Sandstone Member of the Central North Sea (CNS) is a submarine-fan system and potential storage reservoir with a theoretical capacity of 123 Mt of CO₂^[1]. The Tay Sandstone comprises of 4 sequences, amalgamating into a fan complex 125 km long and 40 km wide at a minimum of 1500 m depth, striking NW-SE, hosting several hydrocarbon fields including Gannett A, B, D and Pict.

To better understand the storage potential and characteristics, the Tay Sandstone over Quadrant 21 has been interpreted using geophysical well logs and 3D seismic reflection data. Understanding the internal and external geometry of the sandstone as well as the lateral extent of the unit is essential when considering CO₂ vertical and horizontal fluid flow pathways and storage security.

2. Background

The early-mid Eocene aged Tay Sandstone Member is a submarine-fan system in the Central North Sea (Fig. 2.1). The fan rests on the regional Balder Tuff and is overlain by the thick mudstones of the Horda Formation (Fig. 2.2).

Four internal sequences make up the Tay Sandstone, reducing in size and clay influence with decreasing age. Each sequence separated by mudstones and a minor hiatus.

The Tay Sandstones were sourced from the Scottish Highlands via the Outer Moray Firth, with minor input from the southwest (possibly from the Southern Uplands) (Fig. 2.3).





Oil at the Pict and some of the Gannet fields are in the Tay Sandstone proving the cap rock seal and suggesting good reservoir porosity and permeability.

The Tay Sandstone sits above other Cenozoic sandstones such as the Mey, Cromarty and Forties Members which overlie deeper Jurassic reservoirs. This offers the opportunity Scotland for stacked storage of Co₂ in the CNS.

Figure 2.2. A summary stratigraphic column for the Eocene of the Central North Sea. Including the **Tay Sandstone Member** and overlying mud-rich Horda Formation^[2]



Figure 2.3. Early Eocene palaeogeography and depositional setting showing the extent of the submarine fans of the Tay Sandstone and Grid Sandstone members.^[3]

British Geological Survey, Keyworth, Nottinghamshire, NG12 5GG, United Kingdom

*chrisg@bgs.ac.uk www.bgs.ac.uk

CO₂ Storage Potential of the Eocene Tay Sandstone, Central North Sea

Chris M. A. Gent^{1*} & John D. O. Williams¹

3. Seismic Mapping



The 3D seismic volume used does not incorporate the whole extent of the large Tay Sandstone (Fig. 3.1). The fan strikes NW-SE across the seismic area, and is confirmed by numerous exploration wells.

The Top Tay Sandstone was mapped across the 3D seismic volume. The high amplitude horizon is clear below the thick homogenous Horda Formation. The Top Sele Formation represents the base of the Tay Sandstone Unit, as the Balder Tuff Formation is thin in this area and beneath seismic resolution.



sequence 4 showing the internal geometry of the sandstone

4. Well Information



Fifteen wells within the extent of the seismic volume were used to help build the interpretation. The formation tops were taken from the exploration wells composite logs. Gamma ray traces clearly show the clean 'blocky' sandstones separated by thinner siltstones and mudstones (Fig. 4.1).

A few wells cored the Tay Sandstone, including well 21/17-2 which cored several metres of the overlying Horda Formation. The sandstone show visible porosity and are often poorly consolidated (Fig. 4.2). Core analysis has revealed very good reservoir properties with porosity of 30-39% and permeability of 2-11D

Figure 4.1. The gamma ray log in wells 21/18-7 and 21/18-8 showing the clear 'blocky' response of the sandstone



Figure 3.3. A perpendicular Sub-seismic thicknesses (<40 cross-section across sequence 4. Note the mounded geometry caused by differential compaction.

Internally sequences 1-3 are quiescent but complex, with channel systems and erosional surfaces (Fig. 3.2).

Sequence 4 has a distinctive mounded geometry (Fig. 3.3) with a mappable base, it is 12.8 km by 3.8 km with an average thickness of 97 m (Fig. 3.4). The mapped closure was depth converted using nearby well velocity data.

m) limit the range of the sequence, which could have more laterally extensive thin sandstones. Attribute analysis could help refine the margins of the sequence.



Figure 4.2. Core taken from sequence 3 in 21/17-2. A typical section of clean sandstones. With porosity up to 39% and permeability up to 11D.

5. Storage Potential

With 35% porosity, 5 D permeability and a net to gross of 0.8, sequence 4 of the Tay Sandstone has a high total pore volume of 927x10⁶ m³. The remaining three sequences of stacked channels and interbedded mudstones reduce in net to gross with age as mud has a greater influence in the early fan system. Nevertheless, the sandstone properties remain relatively consistent and are far more laterally extensive than the youngest sequence.

The value of 123Mt CO₂ quoted in the CO₂ storage online atlas CO₂Stored³ is a rough estimate for the whole unit. Mapping undertaken and the pore volumes calculated will lead to a more accurate value of storage potential for this submarine-fan complex.

Several factors and risks still need considering when assessing this site for CO₂ storage: • Pressure connection: dependent on whether the Tay Sandstone is an open or closed unit. • Hydraulic connection with underlying Cenozoic sandstones (Forties, Cromarty, Mey). • Modelling of migration pathways to stratigraphic traps. Connectivity between sequences within the Tay Sandstone.

6. References

¹CO2Stored - Online CO2 Storage Atlas. www.co2stored.co.uk

- Northern North Sea, Keyworth, British Geological Survey for the United Kingdom Offshore Operators Association.





²Knox, R. W. O. B., and Holloway, S., 1992, Lithostratigraphic nomenclature of the UK North Sea : vol. 1 Paleogene of the Central and

³Jennette, D. C., Garfield, T., Mohrig, D., and Cayley, G., 2000. The interaction of shelf accommodation, sediment supply and sea level in controlling the facies, architecture and sequence stacking patterns of the Tay and Forties/Sele basin-floor fans, Central North Sea, in Proceedings Deep-water reservoirs of the world: Gulf Coast Section SEPM 20th Annual Research Conference 2000, SEPM, p. 402-421.

© NERC All Rights Reserved

Seismic data shown courtesy of CDA