

# Short-term iodine dynamics in soil solution

Olivier S. Humphrey<sup>†‡</sup>, Scott D. Young<sup>‡</sup>, Neil M. J. Crout<sup>‡</sup>, Elizabeth H. Bailey<sup>‡</sup>, E. Louise Ander<sup>†</sup>, and Michael J. Watts<sup>†\*</sup>

<sup>†</sup> Inorganic Geochemistry, Centre for Environmental Geochemistry, British Geological Survey, Keyworth, Nottingham NG12 5GG, UK.

<sup>‡</sup> School of Biosciences, University of Nottingham, Sutton Bonington Campus, Leicestershire, LE12 5RD, UK

## ABSTRACT

Assessing the reactions of iodine (I) in soil is critical to evaluating radioiodine exposure and understanding soil-to-crop transfer rates. Our mechanistic understanding has been constrained by method limitations in assessing the dynamic interactions of iodine between soil solution and soil solid phase over short periods (hours). We use microdialysis to passively extract soil solution spiked with radioiodine ( $^{129}\text{I}^-$  and  $^{129}\text{IO}_3^-$ ), to monitor short-term ( $\leq 40$  hours) *in-situ* fixation and speciation changes. We observed greater instantaneous adsorption of  $^{129}\text{IO}_3^-$  compared to  $^{129}\text{I}^-$  in all soils and the complete reduction of  $^{129}\text{IO}_3^-$  to  $^{129}\text{I}^-$  within 5 hours of addition. Loss of  $^{129}\text{I}$  from solution was extremely rapid; the average half-life of  $^{129}\text{I}^-$  and  $^{129}\text{IO}_3^-$  in soil solution was 4.06 and 10.03 hours, respectively. We detected the presence of soluble organically bound iodine (org- $^{129}\text{I}$ ) with a low molecular weight (MW) range (0.5-5 kDa) in all soils and a slower (20 to 40 hours) time-dependent formation of larger MW org-I compounds (12-18 kDa) in some samples. This

20 study highlights the very short window of immediate availability in which I from rainfall or  
21 irrigation remains in soil solution and available to crops, thus presenting significant challenges to  
22 phytofortification strategies in soil-based production systems.

## 23 **Introduction**

24 Iodine (I) is an essential micronutrient for mammalian life, required for the synthesis of thyroid  
25 hormones, approximately 1.9 billion people worldwide are estimated to be at risk of developing  
26 an iodine deficiency disorder.<sup>1,2</sup> Radioiodine (<sup>129</sup>I;  $t_{1/2} = 1.57 \times 10^7$  yr and <sup>131</sup>I;  $t_{1/2}$  8.02 d) released  
27 as a consequence of anthropogenic activities poses a major ecological and health concern due to  
28 its biophilic properties and relatively high environmental mobility.<sup>3</sup> Once <sup>129</sup>I enters the  
29 environment, it behaves similarly to <sup>127</sup>I, capable of bioaccumulating through the food chain and  
30 subsequently stored in the thyroid where it can induce tumors.<sup>4-6</sup> Understanding the  
31 biogeochemical processes and mechanisms that affect I cycling in the terrestrial environment is  
32 critical to evaluating the cause of I deficiency and potentially harmful exposure to radioiodine  
33 isotopes.

34 Soil properties including pH, redox potential, concentration of soil organic matter (SOM) and  
35 Fe/Al/Mn hydrous oxides affect I fixation, mobility and speciation.<sup>5, 7-11</sup> Following natural and  
36 anthropogenic I addition events (e.g. rainfall, irrigation, nuclear weapons testing), soluble  
37 inorganic-I species, iodide ( $I^-$ ) and iodate ( $IO_3^-$ ), can be converted to soluble organic compounds  
38 (org-I) and/or incorporated into the solid soil phase, bound to SOM or adsorbed to metal oxides.<sup>5-  
39 7,9,12</sup> It was demonstrated previously that  $IO_3^-$  and org-I are less mobile and have a greater affinity  
40 for solid-soil components compared to  $I^-$ .<sup>12-14</sup> Metal hydrous oxides can influence iodine dynamics  
41 in soils. Whilst ferric and aluminum oxides adsorb  $IO_3^-$  more strongly than  $I^-$ ,<sup>13</sup> the presence of

42 manganese oxide birnessite ( $\delta$ -MnO<sub>2</sub>) can oxidize I<sup>-</sup> to IO<sub>3</sub><sup>-</sup>.<sup>15</sup> However, in the presence of organic  
43 matter (OM) the oxidation of I<sup>-</sup> to IO<sub>3</sub><sup>-</sup> is limited as the intermediate product, I<sub>2</sub>, is incorporated  
44 into OM (pH <7).<sup>16</sup> Whilst the pH and redox potential of soils affects I speciation and mobility,  
45 SOM appears to be the dominant factor controlling I retention in soils.<sup>12, 17, 18</sup>

46 Microdialysis is an established method for neuroscience and pharmacokinetics and has been  
47 used to investigate the mobility and bioavailability of macro-nutrients and metal ions in soil<sup>19-22</sup>;  
48 its applicability for following I reaction dynamics was previously evaluated by the authors.<sup>23</sup> The  
49 passive sampling conditions of microdialysis enable the extraction of the soil solution phase with  
50 high temporal resolution and minimal disturbance to the soil structure. Unlike conventional soil  
51 solution sampling methods which are destructive (e.g. centrifugation, high-pressure squeezing) or  
52 require soil to be close to saturation (e.g. Rhizon™ samplers), microdialysis can perform  
53 continuous extractions of the soil solution at a wide range of water contents (as low as ~50% water  
54 holding capacity (WHC)). Evaluating soluble I without disturbing the ambient soil conditions,  
55 which could result in changes in I speciation, is vital for a more detailed understanding of I  
56 biogeochemistry in soil.

57 In this study, we investigated short-term soil soluble I dynamics in three soils with contrasting  
58 physicochemical properties. Microdialysis was used to sample an isotopic I tracer (<sup>129</sup>I) under  
59 ambient soil conditions to monitor changes in concentration and speciation following an addition  
60 event at 2.5 hour (hr) intervals over a 40 hr period. Iodine in the sampled dialysate was analyzed  
61 and speciated to determine: (i) the rate at which inorganic-I is removed from soil solution and  
62 adsorbed to the soil solid phase; and, (ii) the molecular weight (MW) range of soluble org-I  
63 compounds which are formed.

64

## 65 **Materials and Methods**

### 66 *Soil sampling and physiochemical characteristics*

67 Three topsoils (0 – 10 cm), from adjoining arable, grassland and woodland sites, were collected  
68 from a location in Nottinghamshire, UK. The soils were sieved to <2 mm, homogenized and air-  
69 dried. Soil pH was determined using an Orion pH meter after equilibrating 5 g of soil in 12.5 mL  
70 of 0.01 M CaCl<sub>2</sub> for 30 min. Loss-on-ignition (LOI), as an estimate of SOM content, was  
71 determined after heating in a muffle furnace at 450 °C for 16 hr, after an initial drying period. The  
72 water holding capacity (WHC) of the soil was determined gravimetrically in triplicate according  
73 to Grace, et al.<sup>24</sup>. Total soil <sup>127</sup>I was extracted with 5% tetramethylammonium hydroxide (TMAH)  
74 at 70 °C for 3 hr (with shaking after 1.5 hr); once cooled, 5 mL of Milli-Q water was added and  
75 the bottles were centrifuged (20 min at 3500 rpm) and the supernatant solutions retained for  
76 analysis by ICP-QQQ.<sup>25</sup> To determine the concentrations of reactive iron, aluminum, and  
77 manganese hydrous oxides, soils were extracted with dithionite-citrate-bicarbonate (DCB)  
78 solution, centrifuged (20 min at 2500 rpm), filtered (<0.22 μm) and the supernatant solutions  
79 retained for analysis by ICP-QQQ.<sup>26</sup> Particle size analysis was determined using the method  
80 described in Rawlins, et al.<sup>27</sup>, using a laser diffraction particle size analyzer.

81

### 82 *Soil incubation and microdialysis sampling*

83 The radioiodine tracer (<sup>129</sup>I) was obtained from the American National Institute of Standards as  
84 NaI (NIST, Gaithersburg, Maryland, USA; CRM 4949C, 0.004 mol L<sup>-1</sup> Na<sup>129</sup>I, 3451 Bq mL<sup>-1</sup>).  
85 The primary stock (<sup>129</sup>I<sup>-</sup>) was made up to 100 mL with 0.01 M NaOH, as recommended by the  
86 suppliers. Iodate (<sup>129</sup>IO<sub>3</sub><sup>-</sup>) was prepared from the <sup>129</sup>I<sup>-</sup> stock by oxidation, using a method adapted  
87 from Yntema and Fleming<sup>28</sup>. To 50 mL of the <sup>129</sup>I<sup>-</sup> stock, 5 mL of 0.1 M HCl was added in an

88 initial neutralization step, followed immediately by 5 mL of 0.2 M sodium chlorite for oxidation.  
89 Successful oxidation to  $^{129}\text{IO}_3^-$  was confirmed by anion exchange LC-ICP-QQQ. The water  
90 content of the air-dried soils (~90 g dry weight (dw)) was raised to 50% of WHC with Milli-Q  
91 water (18.2 M $\Omega$  cm; Millipore) and allowed to equilibrate for 10 days at  $20 \pm 0.5$  °C in the dark.  
92 The moist soils were then homogenized with equivalent volumes of  $^{129}\text{I}^-$  or  $^{129}\text{IO}_3^-$  in solution to  
93 bring the soils to 80% WHC with a final  $^{129}\text{I}$  concentration of 0.2 mg kg $^{-1}$  (dw basis). The soils  
94 were then distributed between triplicate 50 mL Corning® polypropylene centrifuge tubes (~30 g  
95 dw of soil per replicate).

96 The microdialysis system consisted of a syringe pump (KD Scientific Legato 200 Series),  
97 equipped with four plastic syringes (BD Plastipak; 20 mL) used to deliver the perfusate solution.  
98 Syringes were attached to microdialysis probes CMA 20 (10 mm length, 500  $\mu\text{m}$  outer and 400  
99  $\mu\text{m}$  inner diameter) with a polyethersulfone (PES) membrane (100 kDa molecular weight cut-off  
100 (MWCO). Within two minutes of  $^{129}\text{I}$  addition to soil, incision holes were made, using an  
101 introducing needle, and the microdialysis probes were inserted ~1.5 cm beneath the soil surface.  
102 Probes were perfused with Milli-Q water at a flow rate of 5  $\mu\text{L min}^{-1}$  and dialysate solutions  
103 collected at 2.5 hr intervals in glass microvials over a 40 hr extraction period. Samples were  
104 initially stored in the refrigerated micro-fraction collector (6 °C; CMA 470) before freezing at  $-20$   
105 °C until defrosting immediately before analysis. Throughout the extraction, soils were kept in the  
106 dark at a constant temperature of  $20 \pm 0.5$  °C. The probes and micro-fraction collector were from  
107 CMA Microdialysis AB (Kista, Sweden). In order to assess the concentration ( $\mu\text{g L}^{-1}$ ) of  $^{129}\text{I}$  in  
108 the soil solution a correction factor was applied to the measured dialysate solutions as  $C_{\text{soil\_solution}}$   
109 =  $C_{\text{dialysate}} / (1 - e^{-PS/F})$ . Where  $C_{\text{soil\_solution}}$  and  $C_{\text{dialysate}}$  are the concentrations of the target analyte in

110 the external soil solution and the dialysate solution respectively,  $PS$  in the surface area of the  
111 microdialysis membrane and  $F$  is the perfusion flow rate, adapted from Clough<sup>29</sup>.

112

### 113 *Total iodine analysis*

114 Total  $^{127}\text{I}$  concentrations were determined in single MS, no-gas mode (Agilent 8900 ICP-QQQ,  
115 Agilent Technologies). All  $^{129}\text{I}$  concentrations were measured using an ICP-QQQ (Agilent 8900,  
116 Agilent Technologies). Analysis was conducted using both quadrupoles and the reaction cell, to  
117 remove isobaric and polyatomic interferences, a full description of  $^{129}\text{I}$  analysis can be found in  
118 Humphrey, et al.<sup>30</sup>. Due to the small sample volume, we used an Agilent 1260 Infinity Bio-inert  
119 Quaternary LC pump and high performance autosampler to act as a low volume autosampler: vials  
120 were also equipped with flat bottomed glass inserts to further reduce the sample volume  
121 requirement. Throughout the analytical run, an in-house quality control sample with a known  
122 concentration of  $5\ \mu\text{g L}^{-1}$   $^{129}\text{I}$  was measured with an average recovery of 101% ( $n = 28$ ) and a limit  
123 of detection (LOD) ( $3 \times \text{SD blanks}$ ) of  $0.0075\ \mu\text{g L}^{-1}$  for  $^{129}\text{I}$ .

124

### 125 *Iodine speciation analysis*

126 Inorganic-I ( $^{129}\text{I}^-$  and  $^{129}\text{IO}_3^-$ ) concentrations were measured by ICP-QQQ following on-line  
127 chromatographic separation in isocratic mode using a Hamilton PRP X-100, anion exchange  
128 column ( $250\ \text{mm} \times 4.6\ \text{mm}$ ,  $5\ \mu\text{m}$ ) and a samples injection volume of  $25\ \mu\text{L}$ . The mobile phase  
129 was  $100\ \text{mM NH}_4\text{NO}_3$ , adjusted to pH 9.5 with TMAH (25%), at a flow rate of  $1.5\ \text{mL min}^{-1}$ .  
130 Quality control samples with a known I concentration of  $5\ \mu\text{g L}^{-1}$ ,  $^{129}\text{I}^-$  and  $^{129}\text{IO}_3^-$ , were measured  
131 with an average recovery of 97% and 96% ( $n = 22$ ), respectively. The LOD for  $^{129}\text{I}^-$  and  $^{129}\text{IO}_3^-$   
132 was  $0.007$  and  $0.006\ \mu\text{g L}^{-1}$ , respectively.

133 The MW distribution of soluble org-I was determined using size exclusion  
134 chromatography (SEC) hyphenated with UV absorbance detection and ICP-QQQ. An Agilent  
135 1260 Infinity Bio-inert Quaternary LC pump and high performance autosampler in isocratic mode  
136 with an AdvanceBio SEC 130 Å (2.7 µm, 4.6 x 300 mm column, MW range 100-120,000 Da,  
137 Agilent) was used as the sample introduction system. Samples (10 µL) were injected onto the  
138 column with a mobile phase (flow rate of 0.35 mL min<sup>-1</sup>) consisting of 150 mM sodium phosphate  
139 buffer, pH 7.0. Platinum tip sampler and skimmer cones were used on the ICP-QQQ due to the  
140 high sodium concentration in the mobile phase. The UV detector scanned at 220 nm to measure  
141 the calibration protein standards, and 254 nm to detect dissolved organic carbon.<sup>9, 31-35</sup> Protein  
142 standards of known MWs: Ovalbumin (45 kDa); Myoglobin (17 kDa); Aprotinin (6.7 kDa);  
143 Neurotensin (1.7 kDa); and Angiotensin II (1 kDa) (AdvanceBio SEC 130Å protein standard,  
144 Agilent) were used to calibrate the column.

145

## 146 **Results and Discussion**

### 147 *Soil characteristics*

148 The soil physiochemical characteristics in Table 1 reflect their land use. Compared to the arable  
149 and grassland soils the acidic woodland soil had a much greater LOI, extractable Fe/Al/Mn  
150 hydrous oxide content and total I concentration. Given that all three soils were collected with 2.5  
151 km of each other the greater I concentration in the woodland soil may reflect a greater ability to  
152 retain I from precipitation and the absence of annual crop removal.

153

154 **Table 1** Summary of soil physiochemical characteristics. Values expressed as mean ± standard  
155 error (SE) (n=3)

	Units	Arable	Grassland	Woodland
Location (long, lat decimal degrees)		52.9001, -1.0884	52.8890, -1.0877	52.8986, -1.0744
Elevation	m	95	83	121
pH (CaCl <sub>2</sub> )		6.9	5.9	3.6
LOI	%	5.1	7.0	50.6
WHC	%	36.8	41.5	70.8
<sup>127</sup> I	mg kg <sup>-1</sup>	3.38 ± 0.05	3.63 ± 0.05	12.9 ± 0.10
<sup>129</sup> I	µg kg <sup>-1</sup>	<DL	<DL	<DL
Fe <sub>2</sub> O <sub>3</sub>	%	0.180 ± 0.02	0.0131 ± 0.001	0.696 ± 0.06
Al(OH) <sub>3</sub>	%	0.0379 ± 0.003	0.0523 ± 0.007	0.925 ± 0.05
MnO <sub>2</sub>	%	0.0594 ± 0.005	0.0226 ± 0.003	0.622 ± 0.07
Texture (clay/silt/sand)	% mass	41/41/18	17/19/64	24/35/41
Texture classification		Clay	Sandy Loam	Clay Loam


156

157 *Total <sup>129</sup>I dynamics in soil solution*

158 Total concentrations of soluble <sup>129</sup>I sampled from the soils spiked with <sup>129</sup>I<sup>-</sup> or <sup>129</sup>IO<sub>3</sub><sup>-</sup> are shown  
 159 in Figure 1.

160





161  
 162  
 163 **Figure 1** Soluble  $^{129}\text{I}$  ( $\mu\text{g L}^{-1}$ ) sampled using microdialysis from arable, grassland and woodland  
 164 soils at 2.5 hour intervals over 40 hours. Symbols represent mean values obtained in triplicate,  
 165 error bars indicate  $\pm$  S.E. An asterisk at the top left of each chart denotes the initial spike  
 166 concentration; white circles ( $\circ$ ) represent soils spiked with  $^{129}\text{I}^-$ ; black triangles ( $\blacktriangle$ ) represent soils  
 167 spiked with  $^{129}\text{IO}_3^-$ . Note the log scale on the Y-axis. The dashed line indicates the detection limit.  
 168 Each microdialysis dialysate sample was sampled at  $5 \mu\text{L min}^{-1}$  for 2.5 hours.

169  
 170 Losses of  $^{129}\text{I}$  from soil solution were observed in all the soils within 2.5 hr, irrespective of the  
 171 initial spike speciation ( $^{129}\text{I}^-$  or  $^{129}\text{IO}_3^-$ ). The concentration of  $^{129}\text{I}$  in the first dialysate samples (0-  
 172 2.5 hr) were significantly higher in soils treated with  $^{129}\text{I}^-$  compared to  $^{129}\text{IO}_3^-$ . The significantly  
 173 larger initial loss of  $^{129}\text{IO}_3^-$  from solution compared to  $^{129}\text{I}^-$  observed in this study may be a  
 174 combination of adsorption on inorganic soil phases or, rapid immobilization within SOM.<sup>5</sup>

175 Volatilization has previously been reported to occur from soils, but is suppressed by the presence  
176 of SOM<sup>6</sup> and the likelihood of it meaningfully contributing to total loss rates is low.<sup>36</sup> Therefore,  
177 adsorption onto inorganic soil phases and rapid immobilization by SOM are far more likely reasons  
178 for the losses of <sup>129</sup>I from soil solution; both of these processes are pH-dependent. Prior to the  
179 incorporation of IO<sub>3</sub><sup>-</sup> with OM studies have shown that a reduction to an electrophilic intermediate  
180 species, such as HOI or I<sub>2</sub> occurs, this reduction occurs much faster under acidic conditions (pH<  
181 6).<sup>16, 37-40</sup> Clays, hydrous oxides and SOM can absorb I<sup>-</sup>, where sorption generally increases with  
182 decreasing pH, as observed in this study.<sup>41-43</sup> In previous studies it has been shown that when soil  
183 pH >6.0 I<sup>-</sup> sorption is predominantly influenced by OM, however, under more acidic conditions  
184 the role of specific adsorption to iron and aluminum oxides becomes increasingly important.<sup>44, 45</sup>  
185 The faster immediate sorption of <sup>129</sup>IO<sub>3</sub><sup>-</sup> compared to <sup>129</sup>I<sup>-</sup> highlights the different adsorption  
186 mechanisms of inorganic-I in soils.

187 The significant proportion of apparent instantaneous loss of <sup>129</sup>IO<sub>3</sub><sup>-</sup> from solution in all  
188 experimental soil and the highest loss rates of <sup>129</sup>I<sup>-</sup> from soil solution is consistent with previous  
189 results.<sup>5</sup> First-order kinetic equations were used to assess the rate at which total <sup>129</sup>I was removed  
190 from soil solution as:  $C_t = C_0 e^{-kt}$ , where  $C_t$  is the <sup>129</sup>I concentration at time  $t$  (hr) (mg L<sup>-1</sup>),  $C_0$  is  
191 the initial concentration of <sup>129</sup>I in soil solution (mg L<sup>-1</sup>), and  $k$  is the first-order rate constant (hr<sup>-1</sup>).  
192 The half-life in soil solution was then calculated from  $\ln(2)/k$ . The average half-life of soils spiked  
193 with <sup>129</sup>I<sup>-</sup> and <sup>129</sup>IO<sub>3</sub><sup>-</sup> was  $4.06 \pm 0.96$  and  $10.03 \pm 0.78$  hr, with  $r^2$  values of 0.77 and 0.45,  
194 respectively. Shetaya, et al.<sup>5</sup> investigated I dynamics in soils and reported that the loss of I<sup>-</sup> from  
195 solution was extremely rapid, reaching completion over minutes–hours, whereas IO<sub>3</sub><sup>-</sup> loss from  
196 solution was slower, typically occurring over hours–days, which is consistent with the results in  
197 the present study. However, in the current study the use microdialysis has enabled far greater


198 sampling resolution compared to previous studies at time intervals within a similar range to the  
 199 reaction times reported by Shetaya, et al. <sup>5</sup>. The results presented in this paper indicate that iodine  
 200 adsorption occurs more rapidly in soils under realistic conditions than previously observed using  
 201 batch sorption experiments and chemical extraction approaches.

202

203 *<sup>129</sup>I speciation in soil solution*

204 The concentrations of soluble inorganic ( $^{129}\text{I}^- + ^{129}\text{IO}_3^-$ ) and organic (total  $^{129}\text{I}$  minus the sum of  
 205 inorganic species)  $^{129}\text{I}$  in the three soils spiked with either  $^{129}\text{I}^-$  or  $^{129}\text{IO}_3^-$  are shown in Figure 2. A  
 206 combined inorganic fraction was calculated due to the concentrations of  $^{129}\text{IO}_3^-$  present in the soils  
 207 treated with either  $^{129}\text{I}^-$  or  $^{129}\text{IO}_3^-$  falling below the LOD (shown in Figure S1).

208



209

210 **Figure 2** Soluble  $^{129}\text{I}$  speciation of (a) iodate and (b) iodide spiked arable, grassland and woodland  
 211 soils over 40 hours. Symbols represent the mean values obtained in triplicate, bars indicate  $\pm\text{SE}$ .  
 212 White circles ( $\circ$ ) represent inorganic- $^{129}\text{I}$  concentrations; black triangles ( $\blacktriangle$ ) represent soluble

213 org-<sup>129</sup>I concentrations. Note the log scale on the primary Y-axis. Samples below the detection  
214 limit were omitted from the plot. Microdialysis dialysate was sampled at 5  $\mu\text{L min}^{-1}$  for 2.5 hours.

215  
216 The proportion of inorganic-<sup>129</sup>I present in soil solution was greater compared to org-<sup>129</sup>I in soils  
217 spiked with  $\text{I}^-$ , however, this trend was not observed consistently in  $\text{IO}_3^-$  spiked soils. In the arable  
218 soil spiked with  $^{129}\text{IO}_3^-$ , where the org-<sup>129</sup>I became the dominant fraction in samples after 25 hr,  
219 when inorganic-<sup>129</sup>I concentrations were below the LOD. In contrast, in woodland soil spiked with  
220  $^{129}\text{IO}_3^-$  the soluble org-<sup>129</sup>I fraction was below the LOD 10 hr after addition, and only inorganic  
221 species were detected.

222 The speciation of stable and radioiodine present in soil systems has been the focus of several  
223 studies<sup>9, 46-48</sup> with  $\text{I}^-$  and org-I identified as the dominant species present. When  $^{129}\text{IO}_3^-$  was added  
224 to the soils we observed an immediate reduction to  $^{129}\text{I}^-$  in soil solution, with very low  
225 concentrations of  $^{129}\text{IO}_3^-$  remaining in soil solution beyond 2.5 hrs (Figure S1, d). Reduction of  
226  $\text{IO}_3^-$  to  $\text{I}^-$  in soils has previously been observed<sup>49, 50</sup> and may arise from biotic processes. However,  
227 it seems more likely that reduction is caused by abiotic interactions with solid or dissolved OM as  
228 the reaction rates appear to be too rapid for biological processes.<sup>5</sup> The rapid reduction reported in  
229 this study is much faster compared to previous work, where the reduction of  $\text{IO}_3^-$  occurred over  
230 days.<sup>51</sup> Possible reasons for the discrepancies between the studies include the significantly lower  
231 concentrations of <sup>129</sup>I used in this study and the lower soil pH used in this study as the reduction  
232 of  $\text{IO}_3^-$  to the reactive iodine species is thermodynamically more favorable at lower pH.

233 In contrast, when soils were spiked with  $^{129}\text{I}^-$ , a very low concentration of  $^{129}\text{IO}_3^-$  was detected  
234 in the arable and grassland soils within 7.5 hr, indicating that limited  $^{129}\text{I}^-$  oxidation to  $^{129}\text{IO}_3^-$  had  
235 occurred (Figure S1). We only observed  $\text{I}^-$  oxidation in the arable and grassland soils; there was


236 no evidence for oxidation in the organic-rich acidic woodland soil (Figure S1). Factors controlling  
237  $^{129}\text{I}^-$  oxidation in soil remain unclear.<sup>52</sup> The conversion of  $\text{I}^-$  to org-I requires  $\text{I}^-$  to be initially  
238 oxidized to an intermediate such as  $\text{I}_2$  or HOI.<sup>16, 37</sup> Iodide-oxidizing bacteria in soils have  
239 previously been observed to influence  $\text{I}^-$  sorption/incorporation into SOM and oxidation to  $\text{IO}_3^-$   
240 and to produce volatile org-I compounds such as  $\text{CH}_3\text{I}$ .<sup>53, 54</sup> In addition, metal oxides (Al, Fe, Mn)  
241 and humic substances can act as oxidizing agents. In batch kinetic experiments pH and metal oxide  
242 concentrations significantly influence reaction rates, with faster oxidation occurring at lower pH  
243 and greater metal oxide concentrations.<sup>55, 56</sup> However, in our *in-situ* experiments, we did not  
244 observe  $\text{I}^-$  oxidation in the acidic, Fe/Al/Mn rich woodland soil. Figure 2 shows that the greatest  
245 initial loss of  $\text{I}^-$  occurred in the woodland soil, which had the largest metal oxide concentrations,  
246 an order of magnitude higher than the arable and grassland soils, highest organic matter and the  
247 lowest pH. The rapid loss of  $\text{I}^-$  from solution in organic-rich soils has been described as a first-  
248 order reaction.<sup>57</sup> Previous studies have demonstrated that  $\text{I}^-$  can easily be remobilized as it is only  
249 bound by weak electrostatic attraction to solid soil media,<sup>57</sup> therefore, further studies assessing the  
250 speciation of iodine during dynamic processes such as rainfall events or plant uptake should be  
251 investigated to assess iodine mobility in soils.

252

### 253 *Molecular weight distribution of dissolved organic $^{129}\text{I}$*

254 The presence of org- $^{129}\text{I}$  was confirmed using SEC-ICP-QQQ; the presence of multiple peaks  
255 indicates that soluble org-I exists at a range of MWs. Time-dependent changes in speciation and  
256 molecular weight of  $^{129}\text{I}$  in the soil solution phase are shown in Figure 3.

257



258  
 259 **Figure 3** Molecular weight distribution of soluble isotopically labeled  $^{129}\text{I}$  in arable, grassland and  
 260 woodland soils spiked with  $^{129}\text{IO}_3^-$  or  $^{129}\text{I}^-$  sampled over 40 hours, separated by size exclusion  
 261 chromatography. Note the log Y-axis. High molecular weight organically bound I (HMW org-I),  
 262 low molecular weight organically bound I (LMW org-I) and  $^{129}\text{I}^-$  retention times labeled.

263  
 264 Within all three experimental soils, for addition of both  $^{129}\text{I}^-$  and  $^{129}\text{IO}_3^-$ , chromatographic peaks  
 265 were observed for  $^{129}\text{I}^-$  between 15 and 18 min that decreased over the 40 hr extraction period. The  
 266 intensity of the  $^{129}\text{I}^-$  peak was considerably less in the  $^{129}\text{IO}_3^-$  treated soils due to the higher rates  
 267 of instantaneous adsorption on inorganic soil phases and rapid immobilization by SOM. All soils  
 268 spiked with  $^{129}\text{I}^-$ , had a broad peak representing LMW org- $^{129}\text{I}$  between 9 and 15 min; the overall  
 269 intensity of the org-I peak decreased over the 40 hr sampling period. There was a shift in the MW  
 270 distribution in the arable soil spiked with  $^{129}\text{I}^-$  between 20 and 40 hr when the signal intensity of

271 the LMW org-I peak decreased and a HMW org-I peak, with a retention time between 5 and 10  
272 min emerged. In contrast, within the grassland soil spiked with  $^{129}\text{I}^-$ , dissolved org- $^{129}\text{I}$  had a lower  
273 MW distribution; no org-I compounds  $>5$  kDa were present throughout the sampling period. The  
274 emergence and rapid decline of LMW org- $^{129}\text{I}$  between 0.5–5 kDa in the woodland soil further  
275 illustrates how quickly initially soluble I is removed from solution in organic rich soils. However,  
276 within the arable and grassland soils, the MW distribution of the org-I underwent various changes  
277 within the 40 hr sampling period. The UV absorbance results confirmed that soluble organic  
278 compounds had a broad MW range (Figure S2).

279 The transport behavior of inorganic-I in soils may be significantly impacted by the fixation of  
280 dissolved I into macromolecular humic substances, however, many of the mechanisms controlling  
281 the rate of fixation remain unknown.<sup>33, 38, 58, 59</sup> Xu, et al.<sup>3</sup> demonstrated that higher proportions of  
282  $^{125}\text{I}^-$  and  $^{125}\text{IO}_3^-$  were associated with organic compounds  $<3$  kDa, 16 and 20% compared to 6 and  
283 3% in the  $>3$  kDa fraction, respectively. In addition, they found that binding sites became saturated  
284 and less available when the treatment concentration increased.<sup>3</sup> In resuspension experiments, Xu,  
285 et al.<sup>18</sup> found that mobile  $^{129}\text{I}$  was associated with amphiphilic organic compounds with an average  
286 MW between 13.5 and 15 kDa. Figure 3 shows that higher proportions of soluble inorganic-I were  
287 incorporated into organic compounds with a low MW ( $<5$  kDa). However, there was evidence of  
288 larger dissolved org-I compounds with a MW between 12 and 18 kDa 20 hr after  $^{129}\text{I}$  addition, in  
289 agreement with Xu, et al.<sup>3</sup>.

290 The time-dependent formation of these larger dissolved org-I compounds could be due to smaller  
291 dissolved compounds binding together via weak chemical forces or the influence of microbial  
292 activity. The iodination of SOM in acidic conditions was shown to be predominantly an abiotic  
293 process, however, in less acidic conditions ( $\text{pH} \geq 5$ ), microbial assisted iodination of SOM was

294 observed.<sup>3</sup> Whilst microbial activity is generally not deemed essential for the transformation of  
295  $\text{IO}_3^-$  into org-I (which seems to be primarily controlled by abiotic processes<sup>6</sup>), microbial laccases  
296 have been shown to enhance  $\text{I}^-$  sorption in soils.<sup>53</sup> Microbiological activities have previously been  
297 reported to incorporate  $\text{I}^-$  with humic acids and soils with a relatively HMW and low mobility.<sup>33</sup>  
298 Heumann, et al.<sup>60</sup> identified that in the presence of microorganisms the production HMW org-I  
299 compounds significantly increased over an 8 week period; in contrast Figure 3 demonstrates that  
300 formation of relatively HMW compounds occurred over 20-40 hrs.

301 It has previously been suggested that the rapid rate of reaction of I with SOM could be beneficial,  
302 as humus may act as a natural barrier, accumulating any radioiodine released from nuclear waste  
303 repositories as immobile org-I compounds.<sup>18</sup> Following a radioiodine contamination event limited  
304 soil-to-crop transfer would be beneficial for reducing the risk and duration of potentially harmful  
305 exposure.<sup>18, 61, 62</sup> The short-term experiments presented in this paper show that a relatively small  
306 proportion of inorganic-<sup>129</sup>I is rapidly incorporated into LMW organic compounds (<2.5 hr) which,  
307 over time, can be sorbed to the solid soil phase or bind together to form HWM organic compounds  
308 (~30 hr).

309 Evaluating the short-term transport, speciation and fate of I in soil is critical to assess the  
310 environmental mobility and plant availability, as <sup>127</sup>I is an essential micronutrient for which dietary  
311 intake often depends on the transfer of I from soil-to-crops.<sup>63</sup> Soil I concentrations are dependent  
312 on the deposition of volatilized I compounds from seawater.<sup>11</sup> Specific rainfall characteristics  
313 influence the concentration of I in rainfall and soil properties affect the volume of I retained by  
314 soils that would be accessible for plant uptake.<sup>5</sup> Passive uptake is the predominant pathway for I  
315 absorption in plants,<sup>30</sup> as such, plant uptake is highly dependent upon the concentration in the soil  
316 solution phase. By using <sup>129</sup>I as a proxy for <sup>127</sup>I, it is possible to predict how much soluble I is




317 available for plant uptake following an addition event. We demonstrate that I only remains in the  
318 soil solution phase for a very limited period, before being sorbed on to inorganic soil phases or  
319 immobilized by SOM. Consequently, there is a severely limited window of opportunity for plants  
320 to absorb I from the soil solution immediately following rainfall or irrigation and limits the transfer  
321 of I to animal and human diets. However, this study did not investigate the impact of rewetting  
322 and remobilization of I bound to the solid soil phase, and this should be the subject of further  
323 research. Considering that there are an estimated 1.9 billion people at risk of I deficiency  
324 worldwide,<sup>64</sup> the findings in this paper may contribute to better understanding of the efficacy of  
325 phytofortification strategies.

326

### 327 **Supporting Information**

328 Additional information on analytical methods and supplementary results.

329 TOC GRAPHIC FOR MANUSCRIPT



330

331

332 AUTHOR INFORMATION

333 **Corresponding Author**

334 \*Corresponding author: Michael J. Watts; Email: mwatts@bgs.ac.uk; telephone number: 0115 936

335 3042

336 **Present Addresses**

337 † Inorganic Geochemistry, Centre for Environmental Geochemistry, British Geological Survey,

338 Keyworth, Nottingham, NG12 5GG, UK.

339 **Author Contributions**

340 The manuscript was written through contributions from all authors. All authors have given

341 approval to the final version of the manuscript.

342 **Conflict of Interest**

343 The authors declare no competing financial interest.

344 **ACKNOWLEDGMENT**

345 Funding for O. S. Humphrey was provided by The British Geological Survey-University of  
346 Nottingham Centre for Environmental Geochemistry. BGS University Funding Initiative  
347 (contract number: BUFI-S314). This work is published with the permission of the Executive  
348 Director, British Geological Survey.

349

350 **REFERENCES**

- 351 1. WHO; UNICEF; ICCIDD., Assessment of Iodine Deficiency Disorders and monitoring  
352 their Elimination. *World Health Organization Press: Geneva, Switzerland* **2007**.
- 353 2. Zimmermann, M. B.; Jooste, P. L.; Pandav, C. S., Iodine-deficiency disorders. *The*  
354 *Lancet* **2008**, *372*, (9645), 1251-1262.
- 355 3. Xu, C.; Miller, E. J.; Zhang, S.; Li, H.-P.; Ho, Y.-F.; Schwehr, K. A.; Kaplan, D. I.;  
356 Ootosaka, S.; Roberts, K. A.; Brinkmeyer, R.; Yeager, C. M.; Santschi, P. H., Sequestration and  
357 Remobilization of Radioiodine (129I) by Soil Organic Matter and Possible Consequences of the  
358 Remedial Action at Savannah River Site. *Environmental Science & Technology* **2011**, *45*, (23),  
359 9975-9983.
- 360 4. Takeda, A.; Tsukada, H.; Takaku, Y.; Satta, N.; Baba, M.; Shibata, T.; Hasegawa, H.;  
361 Unno, Y.; Hisamatsu, S., Determination of Iodide, Iodate and Total Iodine in Natural Water  
362 Samples by HPLC with Amperometric and Spectrophotometric Detection, and Off-line UV  
363 Irradiation. *Analytical Sciences* **2016**, *32*, (8), 839-845.
- 364 5. Shetaya, W. H.; Young, S. D.; Watts, M. J.; Ander, E. L.; Bailey, E. H., Iodine dynamics  
365 in soils. *Geochimica et Cosmochimica Acta* **2012**, *77*, 457-473.
- 366 6. Yamaguchi, N.; Nakano, M.; Takamatsu, R.; Tanida, H., Inorganic iodine incorporation  
367 into soil organic matter: evidence from iodine K-edge X-ray absorption near-edge structure.  
368 *Journal of Environmental Radioactivity* **2010**, *101*, (6), 451-457.
- 369 7. Bowley, H. E.; Young, S. D.; Ander, E. L.; Crout, N. M. J.; Watts, M. J.; Bailey, E. H.,  
370 Iodine binding to humic acid. *Chemosphere* **2016**, *157*, 208-214.
- 371 8. Yamada, H.; Kiriya, T.; Onagawa, Y.; Hisamori, I.; Miyazaki, C.; Yonebayashi, K.,  
372 Speciation of iodine in soils. *Soil science and plant nutrition* **1999**, *45*, (3), 563-568.
- 373 9. Shimamoto, Y. S.; Takahashi, Y.; Terada, Y., Formation of Organic Iodine Supplied as  
374 Iodide in a Soil–Water System in Chiba, Japan. *Environmental Science & Technology* **2011**, *45*,  
375 (6), 2086-2092.
- 376 10. Whitehead, D. C., The sorption of iodide by soils as influenced by equilibrium conditions  
377 and soil properties. *Journal of the Science of Food and Agriculture* **1973**, *24*, (5), 547-556.

- 378 11. Humphrey, O. S.; Young, S. D.; Bailey, E. H.; Crout, N. M. J.; Ander, E. L.; Watts, M. J.,  
379 Iodine soil dynamics and methods of measurement: a review. *Environmental Science: Processes  
380 & Impacts* **2018**, *20*, (2), 288-310.
- 381 12. Schwehr, K. A.; Santschi, P. H.; Kaplan, D. I.; Yeager, C. M.; Brinkmeyer, R., Organo-  
382 Iodine Formation in Soils and Aquifer Sediments at Ambient Concentrations. *Environmental  
383 Science & Technology* **2009**, *43*, (19), 7258-7264.
- 384 13. Kodama, S.; Takahashi, Y.; Okumura, K.; Uruga, T., Speciation of iodine in solid  
385 environmental samples by iodine K-edge XANES: Application to soils and ferromanganese  
386 oxides. *Science of The Total Environment* **2006**, *363*, (1–3), 275-284.
- 387 14. Shimamoto, Y. S.; Itai, T.; Takahashi, Y., Soil column experiments for iodate and iodide  
388 using K-edge XANES and HPLC–ICP-MS. *Journal of Geochemical Exploration* **2010**, *107*, (2),  
389 117-123.
- 390 15. Gallard, H.; Allard, S. b.; Nicolau, R.; Von Gunten, U.; Croué, J. P., Formation of  
391 iodinated organic compounds by oxidation of iodide-containing waters with manganese dioxide.  
392 *Environmental science & technology* **2009**, *43*, (18), 7003-7009.
- 393 16. Warner, J. A.; Casey, W. H.; Dahlgren, R. A., Interaction Kinetics of I<sub>2</sub>(aq) with  
394 Substituted Phenols and Humic Substances. *Environmental Science & Technology* **2000**, *34*,  
395 (15), 3180-3185.
- 396 17. Kaplan, D. I., Influence of surface charge of an Fe-oxide and an organic matter  
397 dominated soil on iodide and pertechnetate sorption. *Radiochim. Acta* **2003**, *91*, 173-178.
- 398 18. Xu, C.; Zhang, S.; Ho, Y.-F.; Miller, E. J.; Roberts, K. A.; Li, H.-P.; Schwehr, K. A.;  
399 Ootaka, S.; Kaplan, D. I.; Brinkmeyer, R.; Yeager, C. M.; Santschi, P. H., Is soil natural organic  
400 matter a sink or source for mobile radioiodine (129I) at the Savannah River Site? *Geochimica et  
401 Cosmochimica Acta* **2011**, *75*, (19), 5716-5735.
- 402 19. Miró, M.; Jimoh, M.; Frenzel, W., A novel dynamic approach for automatic  
403 microsampling and continuous monitoring of metal ion release from soils exploiting a dedicated  
404 flow-through microdialyser. *Analytical and Bioanalytical Chemistry* **2005**, *382*, (2), 396-404.
- 405 20. Brackin, R.; Näsholm, T.; Robinson, N.; Guillou, S.; Vinall, K.; Lakshmanan, P.;  
406 Schmidt, S.; Inselsbacher, E., Nitrogen fluxes at the root-soil interface show a mismatch of  
407 nitrogen fertilizer supply and sugarcane root uptake capacity. *Scientific Reports* **2015**, *5*, 15727.
- 408 21. Buckley, S.; Brackin, R.; Näsholm, T.; Schmidt, S.; Jämtgård, S., Improving in situ  
409 recovery of soil nitrogen using the microdialysis technique. *Soil Biology and Biochemistry* **2017**,  
410 *114*, 93-103.
- 411 22. Demand, D.; Schack-Kirchner, H.; Lang, F., Assessment of diffusive phosphate supply in  
412 soils by microdialysis. *Journal of Plant Nutrition and Soil Science* **2017**, (180), 220-230.
- 413 23. Humphrey, O. S.; Young, S. D.; Bailey, E. H.; Crout, N. M. J.; Ander, E. L.; Hamilton,  
414 E. M.; Watts, M. J., Investigating the use of microdialysis and SEC-UV-ICP-MS to assess iodine  
415 interactions in soil solution. *Chemosphere* **2019**, *229*, 41-50.
- 416 24. Grace, C.; Hart, M.; Brookes, P. C., Laboratory manual of the soil microbial biomass  
417 group. *Rothamsted Research* **2006**, 65.
- 418 25. Watts, M.; Mitchell, C., A pilot study on iodine in soils of Greater Kabul and Nangarhar  
419 provinces of Afghanistan. *Environmental Geochemistry and Health* **2009**, *31*, (4), 503-509.
- 420 26. Mehra, O. P.; Jackson, M. L., Iron oxide removal from soils and clays by a dithionite-  
421 citrate system buffered with sodium bicarbonate. In *Clays Clay Miner*, Pergamon: 1960; pp 317-  
422 327.

- 423 27. Rawlins, B. G.; Wragg, J.; Lark, R. M., Application of a novel method for soil aggregate  
424 stability measurement by laser granulometry with sonication. *European Journal of Soil Science*  
425 **2013**, *64*, (1), 92-103.
- 426 28. Yntema, L. F.; Fleming, T., Volumetric Oxidation of Iodide to Iodate by Sodium  
427 Chlorite. *Industrial & Engineering Chemistry Analytical Edition* **1939**, *11*, (7), 375-377.
- 428 29. Clough, G. F., Microdialysis of large molecules. *The AAPS Journal* **2005**, *7*, (3), E686-  
429 E692.
- 430 30. Humphrey, O. S.; Young, S. D.; Bailey, E. H.; Crout, N. M. J.; Ander, E. L.; Hamilton,  
431 E. M.; Watts, M. J., Iodine uptake, storage and translocation mechanisms in spinach (*Spinacia*  
432 *oleracea* L.). *Environmental Geochemistry and Health* **2019**, *41*, (5), 2145-2156.
- 433 31. Yamada, H.; Hisamori, I.; Yonebayashi, K., Identification of organically bound iodine in  
434 soil humic substances by size exclusion chromatography / inductively coupled plasma mass  
435 spectrometry (SEC / ICP-MS). *Soil Science and Plant Nutrition* **2002**, *48*, (3), 379-385.
- 436 32. Rädlinger, G.; Heumann, K. G., Determination of halogen species of humic substances  
437 using HPLC/ICP-MS coupling. *Fresenius J Anal Chem* **1997**, *359*, (4-5), 430-433.
- 438 33. Rädlinger, G.; Heumann, K. G., Transformation of Iodide in Natural and Wastewater  
439 Systems by Fixation on Humic Substances. *Environmental Science & Technology* **2000**, *34*, (18),  
440 3932-3936.
- 441 34. Takeda, A.; Tsukada, H.; Takaku, Y.; Hisamatsu, S. i., Fractionation of metal complexes  
442 with dissolved organic matter in a rhizosphere soil solution of a humus-rich Andosol using size  
443 exclusion chromatography with inductively coupled plasma-mass spectrometry. *Soil Science and*  
444 *Plant Nutrition* **2009**, *55*, (3), 349-357.
- 445 35. Shah, M.; Wuilloud, R. G.; Kannamkumarath, S. S.; Caruso, J. A., Iodine speciation  
446 studies in commercially available seaweed by coupling different chromatographic techniques  
447 with UV and ICP-MS detection. *Journal of Analytical Atomic Spectrometry* **2005**, *20*, (3), 176-  
448 182.
- 449 36. Sheppard, M.; Thibault, D.; Smith, P.; Hawkins, J., Volatilization: a soil degassing  
450 coefficient for iodine. *Journal of environmental radioactivity* **1994**, *25*, (3), 189-203.
- 451 37. Schlegel, M. L.; Reiller, P.; Mercier-Bion, F.; Barré, N.; Moulin, V., Molecular  
452 environment of iodine in naturally iodinated humic substances: Insight from X-ray absorption  
453 spectroscopy. *Geochimica et Cosmochimica Acta* **2006**, *70*, (22), 5536-5551.
- 454 38. Steinberg, S. M.; Schmett, G. T.; Kimble, G.; Emerson, D. W.; Turner, M. F.; Rudin, M.,  
455 Immobilization of fission iodine by reaction with insoluble natural organic matter. *Journal of*  
456 *Radioanalytical and Nuclear Chemistry* **2008**, *277*, (1), 175-183.
- 457 39. Francois, R., The influence of humic substances on the geochemistry of iodine in  
458 nearshore and hemipelagic marine sediments. *Geochimica et Cosmochimica Acta* **1987**, *51*, (9),  
459 2417-2427.
- 460 40. Brummer, J. G.; Field, R. J., Kinetics and mechanism of the oxidation of ferrous ion by  
461 iodate ion in strong perchloric acid, aqueous media. *Journal of Physical Chemistry* **1979**, *83*,  
462 (18), 2328-2335.
- 463 41. Whitehead, D. C., The distribution and transformations of iodine in the environment.  
464 *Environment International* **1984**, *10*, (4), 321-339.
- 465 42. Mackowiak, C. L.; Grossl, P. R.; Cook, K. L., Iodine toxicity in a plant-solution system  
466 with and without humic acid. *Plant and Soil* **2005**, *269*, (1-2), 141-150.
- 467 43. Kaplan, D. I.; Serne, R. J.; Parker, K. E.; Kutnyakov, I. V., Iodide sorption to subsurface  
468 sediments and illitic minerals. *Environmental Science & Technology* **2000**, *34*, (3), 399-405.

- 469 44. Yoshida, S.; Muramatsu, Y.; Uchida, S., Studies on the sorption of I<sup>-</sup> (iodide) and IO<sub>3</sub><sup>-</sup>  
470 (iodate) onto Andosols. *Water, Air, and Soil Pollution* **1992**, *63*, (3), 321-329.
- 471 45. Whitehead, D. C., The influence of organic matter, chalk, and sesquioxides on the  
472 solubility of iodide, elemental iodine, and iodate incubated with soil. *Journal of Soil Science*  
473 **1974**, *25*, (4), 461-470.
- 474 46. Yoshida, S.; Muramatsu, Y.; Katou, S.; Sekimoto, H., Determination of the chemical  
475 forms of iodine with IC-ICP-MS and its application to environmental samples. *Journal of*  
476 *Radioanalytical and Nuclear Chemistry* **2007**, *273*, (1), 211-214.
- 477 47. Yamada, H.; Kiriya, T.; Yonebayashi, K., Determination of Total Iodine in Soils by  
478 Inductively Coupled Plasma Mass Spectrometry. *Soil Science and Plant Nutrition* **1996**, *42*, (4),  
479 859-866.
- 480 48. Muramatsu, Y.; Uchida, S.; Sumiya, M.; Ohmomo, Y.; Obata, H., Tracer experiments on  
481 transfer of radio-iodine in the soil—rice plant system. *Water, Air, and Soil Pollution* **1989**, *45*,  
482 (1-2), 157-171.
- 483 49. Zhang, S.; Du, J.; Xu, C.; Schwehr, K.; Ho, Y.-F.; Li, H.-P.; Roberts, K.; Kaplan, D.;  
484 Brinkmeyer, R.; Yeager, C., Concentration-dependent mobility, retardation, and speciation of  
485 iodine in surface sediment from the Savannah River Site. *Environmental science & technology*  
486 **2011**, *45*, (13), 5543-5549.
- 487 50. Muramatsu, Y.; Uchida, S.; Sriyotha, P.; Sriyotha, K., Some considerations on the  
488 sorption and desorption phenomena of iodide and iodate on soil. *Water, Air, and Soil Pollution*  
489 **1990**, *49*, (1), 125-138.
- 490 51. Yamaguchi, N.; Nakano, M.; Tanida, H.; Fujiwara, H.; Kihou, N., Redox reaction of  
491 iodine in paddy soil investigated by field observation and the I K-Edge XANES fingerprinting  
492 method. *Journal of Environmental Radioactivity* **2006**, *86*, (2), 212-226.
- 493 52. Amachi, S., Microbial Contribution to Global Iodine Cycling: Volatilization,  
494 Accumulation, Reduction, Oxidation, and Sorption of Iodine. *Microbes and Environments* **2008**,  
495 *23*, (4), 269-276.
- 496 53. Seki, M.; Oikawa, J.-i.; Taguchi, T.; Ohnuki, T.; Muramatsu, Y.; Sakamoto, K.; Amachi,  
497 S., Laccase-Catalyzed Oxidation of Iodide and Formation of Organically Bound Iodine in Soils.  
498 *Environmental Science & Technology* **2012**, *47*, (1), 390-397.
- 499 54. Amachi, S.; Kasahara, M.; Hanada, S.; Kamagata, Y.; Shinoyama, H.; Fujii, T.;  
500 Muramatsu, Y., Microbial Participation in Iodine Volatilization from Soils. *Environmental*  
501 *Science & Technology* **2003**, *37*, (17), 3885-3890.
- 502 55. Fox, P. M.; Davis, J. A.; Luther III, G. W., The kinetics of iodide oxidation by the  
503 manganese oxide mineral birnessite. *Geochimica et Cosmochimica Acta* **2009**, *73*, (10), 2850-  
504 2861.
- 505 56. Allard, S.; von Gunten, U.; Sahli, E.; Nicolau, R.; Gallard, H., Oxidation of iodide and  
506 iodine on birnessite ( $\delta$ -MnO<sub>2</sub>) in the pH range 4–8. *Water Research* **2009**, *43*, (14), 3417-3426.
- 507 57. Sheppard, M. I.; Thibault, D. H., Chemical behaviour of iodine in organic and mineral  
508 soils. *Applied Geochemistry* **1992**, *7*, (3), 265-272.
- 509 58. Tikhomirov, F. A.; Kasparov, S. V.; Prister, B. S.; Sal'nikov, V. n., Role of organic  
510 matter in iodine fixation in soils. *Soviet Soil Science* **1980**, *12*, (1), 64-72.
- 511 59. Hu, Q. H.; Moran, J. E.; Gan, J. Y., Sorption, degradation, and transport of methyl iodide  
512 and other iodine species in geologic media. *Applied Geochemistry* **2012**, *27*, (3), 774-781.

- 513 60. Heumann, K. G.; Rädlinger, G.; Erbes, M.; Heiber, I.; Obst, U.; Filip, Z.; Claus, H.,  
514 Ageing of Dissolved Halogenated Humic Substances and the Microbiological Influence on this  
515 Process. *Acta hydrochimica et hydrobiologica* **2000**, 28, (4), 193-201.
- 516 61. Xu, C.; Zhong, J.; Hatcher, P. G.; Zhang, S.; Li, H.-P.; Ho, Y.-F.; Schwehr, K. A.;  
517 Kaplan, D. I.; Roberts, K. A.; Brinkmeyer, R., Molecular environment of stable iodine and  
518 radioiodine (<sup>129</sup>I) in natural organic matter: Evidence inferred from NMR and binding  
519 experiments at environmentally relevant concentrations. *Geochimica et Cosmochimica Acta*  
520 **2012**, 97, 166-182.
- 521 62. Keppler, F.; Biester, H.; Putschew, A.; Silk, P. J.; Schöler, H. F.; Müller, G.,  
522 Organoiodine formation during humification in peatlands. *Environmental Chemistry Letters*  
523 **2003**, 1, (4), 219-223.
- 524 63. Watts, M.; Joy, E.; Young, S.; Broadley, M.; Chilimba, A.; Gibson, R.; Siyame, E.;  
525 Kalimbira, A.; Chilima, B.; Ander, E., Iodine source apportionment in the Malawian diet.  
526 *Scientific reports* **2015**, 5, (15251).
- 527 64. Zimmermann, M. B.; Gizak, M.; Abbott, K.; Andersson, M.; Lazarus, J. H., Iodine  
528 deficiency in pregnant women in Europe. *The Lancet Diabetes & Endocrinology* **2015**, 3, (9),  
529 672-674.

530