

POLICY BRIEF

Open Access



# Using environmental monitoring data from apex predators for chemicals management: towards better use of monitoring data from apex predators in support of prioritisation and risk assessment of chemicals in Europe

Gabriele Treu<sup>1\*</sup>, Jaroslav Slobodnik<sup>2</sup>, Nikiforos Alygizakis<sup>2,3</sup>, Alexander Badry<sup>1</sup>, Dirk Bunke<sup>4</sup>, Alessandra Cincinelli<sup>5</sup>, Daniela Claßen<sup>1</sup>, Rene W. R. J. Dekker<sup>6</sup>, Bernd Göckener<sup>9</sup>, Georgios Gkotsis<sup>3</sup>, Georg Hanke<sup>10</sup>, Guy Duke<sup>8</sup>, Morten Jartun<sup>11</sup>, Paola Movalli<sup>6</sup>, Maria-Christina Nika<sup>3</sup>, Heinz Rüdell<sup>9</sup>, Jose V. Tarazona<sup>12</sup>, Nikolaos S. Thomaidis<sup>3</sup>, Victoria Tornero<sup>10</sup>, Katrin Vorkamp<sup>13</sup>, Lee A. Walker<sup>14</sup>, Jan Koschorreck<sup>1</sup> and Valeria Dulio<sup>7</sup>

## Abstract

A large number of apex predator samples are available in European research collections, environmental specimen banks and natural history museums that could be used in chemical monitoring and regulation. Apex predators bioaccumulate pollutants and integrate contaminant exposure over large spatial and temporal scales, thus providing key information for risk assessments. Still, present assessment practices under the different European chemical legislations hardly use existing chemical monitoring data from top predators. Reasons include the lack of user-specific guidance and the fragmentation of data across time and space. The European LIFE APEX project used existing sample collections and applied state-of-the-art target and non-target screening methods, resulting in the detection of > 4,560 pollutants including legacy compounds. We recommend establishing infrastructures that include apex predators as an early warning system in Europe. Chemical data of apex species from freshwater, marine and terrestrial compartments should become an essential component in future chemical assessment and management across regulations, with the purpose to (1) validate registration data with 'real world' measurements and evaluate the predictability of current models; (2) identify and prioritise hazardous chemicals for further assessment; (3) use data on food web magnification as one line of evidence to assess biomagnification; (4) determine the presence of (bio)transformations products and typical chemical mixtures, and (5) evaluate the effectiveness of risk management measures by trend analysis. We highlight the achievements of LIFE APEX with regard to novel trend and mixture analysis tools and prioritisation schemes. The proposed advancements complement current premarketing regulatory assessments and will allow the detection

\*Correspondence: gabriele.treu@uba.de

<sup>1</sup> German Environment Agency (Umweltbundesamt), Wörlitzer Platz 1, 06813 Dessau-Roßlau, Germany  
Full list of author information is available at the end of the article

of contaminants of emerging concern at an early stage, trigger risk management measures and evaluations of their effects with the ultimate goal to protect humans and the environment. This is the second policy brief of the LIFE APEX project.

**Keywords:** Bioaccumulation, Chemical regulation, Early warning system, Wide scope target and non-target screening, Wildlife

## Status quo in the use of apex predators in chemicals management

### Pollution is one of the greatest Anthropocene challenges

The intention of global and European chemical regulations such as the Regulation on Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH) [1] is to protect the environment and humans from hazardous substances. Despite these efforts, chemical pollution has become one of the Anthropocene's greatest environmental challenges. Humanity has likely exceeded safe thresholds of planetary boundaries for chemical pollution [2] with humans and wildlife in Europe are being exposed to a wide range of types of chemical pollution including plant protection products, biocides, pharmaceuticals, industrial chemicals and personal care products together with numerous transformation products, which pose substantial threat to biodiversity [3]. Concerns exist for apex predators in particular as they are exposed to high concentrations of biomagnifying substances [4, 5]. Recognising these challenges, the European Commission (EC) has developed strategies, enacted legally binding regulations and joined multilateral agreements. The European Green Deal [6] sets a 'zero pollution ambition' and measures to achieve this are set out in the EU Action Plan: 'Towards Zero Pollution for Air, Water and Soil' [7] and the European Chemicals Strategy for Sustainability Towards a Toxic-Free Environment [8]. The key challenge is to streamline the prioritisation, assessment and, where necessary, the restriction of the many thousands of chemicals marketed under various regulations such that chemical pollution is prevented more effectively in the future.

### Apex predators are underused sentinels for pollutant exposure

Top predators such as raptors and marine mammals play a key role in contaminant monitoring and wildlife toxicology as they integrate contaminant exposure over a long lifetime and, in some cases, large foraging areas. Certain chemicals bioaccumulate in their tissues and, with high energy turnover from lower trophic levels, biomagnify to high concentrations through the food chain. Population declines of top predators have been among the most alarming impacts of chemical pollution and

have driven public pressure to enact treaties aimed at reducing such pollution [9, 10]. Persistence and bioaccumulation of chemicals are of particular concern to regulators since they can lead to chronic exposure to high and unpredictable levels [11]. Accordingly, a number of regulatory frameworks specifically address the bioaccumulation in top predators, such as the Stockholm Convention [12] or assessing persistent (P), bioaccumulative (B) and toxic (T) or very persistent (vP) and very bioaccumulative (vB) property assessments under REACH, the EU Water Framework Directive (WFD, [13], and the EU Marine Strategy Framework Directive (MSFD) [14].

The European Chemicals Strategy calls for a more efficient use of contaminant monitoring data from humans and ecosystems as key to improve risk assessment and the understanding of pollution impacts [8]. Furthermore, there is a need for an effective early warning system based on monitoring data and other indicators. Pioneering work with raptors under the European Raptor Biomonitoring Facility (ERBFacility, [www.erbfacility.eu](http://www.erbfacility.eu)) has provided guidance on species and matrix selection for pan-European contaminant monitoring [15, 16] including a review of archive capacities [17, 18]. However, a European research infrastructure designed to support contaminant screening using top predators from different environmental compartments as bioindicators and as sentinels for early warning is still lacking. At present, many contaminant monitoring data in top predators are fragmented over time, are regionally dispersed, and often motivated by project-based initiatives rather than meeting the long-term needs of a comprehensive exposure assessment in the context of chemical risk and hazard assessment and the needs for effectiveness evaluations. To date, few monitoring activities (e.g. under the MSFD or Regional Sea Conventions) are organised in a consistent way over time that allows for temporal trend analysis. Furthermore, these monitoring data are not stored in a centralised database yet. This is a major drawback in sustainable chemicals management, including the safe placing on the market, and post-market surveillance, of substances and products, and for assessing the effectiveness of regulatory measures, especially considering the evolving patterns of chemical substitution and diversification.

With recent advances in analytical chemistry, the scientific and technical means are now available to generate unparalleled amounts of data for chemical and (eco) toxicological fingerprinting and the biological and ecological functions of ecosystems [19]. For instance, rapidly advancing non-target and suspect screening techniques based on high-resolution mass spectrometric data now allow the screening of thousands of contaminants, which can substantially support the identification and risk assessment of contaminants of emerging concern [20–22]. Applied to a large number of yet underused biota and apex predator specimens available in research collections, natural history museums and environmental specimen banks [18, 23–25] these techniques offer new possibilities to characterise biomagnifying contaminants of emerging concern, and to use top predators both as an early warning system and for long-term monitoring.

#### **EU prioritisation and risk assessment practices are often limited and lack a field context**

The EC aims at high-quality risk assessment outputs to avoid costly impacts on human and environmental health and ineffective risk management measures. Current screening, prioritisation and assessment methods as applied under the different European chemical and environmental regulations are usually based on *in silico* approaches, *in vitro* data and/or *in vivo* testing using algae, aquatic invertebrates, and fish to estimate chemical effects on a population or ecosystem. Higher tier tests like mesocosm studies or field studies are in general not required as part of the authorisation process of chemicals with the exception of plant protection products. Thus, present hazard and risk assessments to demonstrate the safe use of substances are mainly based on data generated under laboratory conditions and exposure modelling and do not reflect an empirical ecological context [26]. For REACH chemicals a major challenge remains since only a small fraction of the marketed chemicals has been sufficiently evaluated regarding their (eco)toxicological properties and exposure, while limited data are available for the majority of substances [27]. Many of the >22,000 compounds registered under REACH either lack data due to poor quality of registration dossiers or data are not sufficient for a comprehensive assessment, e.g. of PBT properties [28]. One reason is that chemical legislation requires only lower tier hazard testing for REACH chemicals with a production volume <10 tonnes per year. Moreover, terrestrial ecosystems are as yet hardly considered in environmental risk assessment. For instance, existing assessment strategies and test guidelines to assess bioaccumulation are restricted to aquatic organisms and do not consider the accumulation of substances in air-breathing organisms and biomagnification

in terrestrial food webs ([29]). In general, risk assessment guidelines are based on conservative approaches, but risk is assessed for each intended use and scenario which does not cover the aggregated exposure at landscape level from different current and historical uses, including hot-spots and biomagnification processes relevant for apex predators. In addition, current approaches neglect the fact that humans and ecosystems are exposed to chemical mixtures leading to potential cumulative effects even if the concentrations of individual substances remain below their respective effect thresholds [30–32]. Moreover, risk assessment and regulatory action is still assessed substance by substance and lags behind the rapid increase in the volume and diversity of chemicals [33, 34]. Given the increasing number of substances on the market [33] it becomes unlikely that these compounds can all be screened and assessed for hazards and risks individually without enormous efforts. Regulatory risk assessments focus on premarket authorisations, and should be complemented with post-market environmental vigilance. Landscape aggregated exposure models have been conceptualised [35, 36], but are still under early development, particularly in connection with population dynamics [37]. In this context, monitoring data using apex predators can be used to improve the current risk assessment methods as well as for realistic retrospective assessments. This calls for automated, cost efficient and effective tools for prioritisation of pollutants and use of *all* available resources, including the systematic use of environmental and human biomonitoring data. Current monitoring programmes under the WFD or MSFD are insufficient in this respect as they focus on few selected compounds and are mostly based on simplistic prioritisation approaches.

#### **Lack of guidance on the generation and use of chemical monitoring data**

A particular challenge and a reason for the limited use of biota field data is the lack of guidance on harmonised sampling, processing, archiving, and shipping of wildlife samples, which is an important prerequisite to generate high quality and comparable data. The comparison of premarket estimations with monitoring data is essential for assessing real exposure and effect levels, as well as for updating risk assessment methods. Additionally, practical advice for industry and regulators on the use of biota field data under the different European chemical regulations are lacking such as the Guidance on PBT assessment developed by the European Chemicals Agency [38, 39]. Limited access to data from research programmes is a general challenge. Consensus upon consistent guidance and assessment approaches using biota field data is currently hampered by the fact that registration and

risk assessment of chemicals on the EU market is fragmented across different legal frameworks [40]. This is now addressed by the EU Chemicals Strategy for Sustainability, which fosters coherent policies and action by all relevant international organisations.

## Achievements

### Lessons learned from the LIFE APEX project

Aims and scope of the LIFE APEX project are explained in detail in the first policy brief [41]. In total, about 200 pooled top predator and prey samples were screened for organic pollutants by wide-scope target and suspect screening, for presence of more than 65,000 chemicals. Overall, about 4,500 chemicals were detected of the assessed biota samples, i.e. freshwater fish, and marine mammals, raptors and fish. Those compounds included man-made pollutants and some naturally occurring compounds. These results reflect that apex species were exposed to thousands of pollutants, making up chemical cocktails that are covered by different regulations. Chemical mixtures included compounds known or expected to bioaccumulate in top predators but also a wide range of substances that had not been detected before, including plant protection products, biocides, human pharmaceuticals and industrial chemicals. While the LIFE APEX project demonstrated that contamination in European apex predators was widespread, the effects at the population level remain largely unknown. LIFE APEX also demonstrated that existing sample collections could be combined with state-of-the-art target, suspect and non-target screening methods to provide a more comprehensive picture of chemical exposure in samples of top predators and prey. We further provided these data to regulators for respective risk assessment. All LIFE APEX chemical data are publicly available through NORMAN and LIFE APEX database systems (<https://www.norman-network.com/apex/>). Specific LIFE APEX actions and case studies included:

### ***Robust prioritisation and screening scheme using chemical data from apex species***

All substances detected in top predators within LIFE APEX have been screened with the open access JANUS software ([www.vegahub.eu/portfolio-item/janus/](http://www.vegahub.eu/portfolio-item/janus/)), which allows the identification of hazardous chemical properties based on laboratory data and in silico estimates for PBT, endocrine disruption and cancerogenic, mutagenic and reprotoxic (CMR) properties. JANUS is based on a battery of quantitative structure–activity relationship (QSAR) models in specific workflows for each endpoint [42, 43]. A list of top-ranked pollutants (ranked by hazard scores, regulatory status and frequency of appearance)

was provided to ECHA and national regulatory agencies to aid further assessment.

In addition to hazard prioritisation, LIFE APEX data also allow the identification and exploration of inconsistencies between exposure estimates and registration status, e.g. if substances pre-registered under REACH are frequently found in apex predators, even if they should not be on the market or produced in Europe. Overall, we demonstrated that LIFE APEX approaches, including the prioritisation methods, were suitable to identify previously unknown and potentially problematic substances from a large set of chemicals detected in top predators and to make this information available to regulators.

### ***Use of LIFE APEX chemical data in support of prioritisation, substance evaluation and restriction under REACH***

The list of top-ranked chemicals, as described above, was subject to further in-depth assessment by the German Environment Agency with regard to PBT properties. REACH explicitly requires that all available information in registration dossiers and the open literature, including monitoring data, be considered in a weight of evidence approach to draw a conclusion on hazard endpoints. Thus, this in-depth assessment followed procedures for substance evaluation as laid out in REACH (Annex XIII; [38, 39]) considering physico-chemical, in vitro, in silico and in vivo data (if available) from the REACH dossiers as well as data derived from LIFE APEX.

The results from the LIFE APEX project were used to support regulatory substance evaluations under REACH. As an example, the occurrence and concentrations of the fragrance Galaxolide (CAS: 1222-05-5) measured in LIFE APEX were used as additional evidence to evaluate the bioaccumulation behaviour. Galaxolide has been initially found not to meet PBT properties [44] before REACH. New data on intrinsic properties for a re-assessment of the PBT status of Galaxolide, especially for persistence and bioaccumulation, are now available. Thus, France has initiated a re-evaluation of Galaxolide by the inclusion of the substance on the community rolling action plan in 2022 in order to clarify concerns related to its suspected PBT/vPvB properties, potential endocrine disruptor, consumer use and exposure of environment. According to a preliminary bioaccumulation assessment, standard laboratory data reported bioconcentration factors (BCF) for fish between 600 and 1,600 L/kg, which were below the B-criterion (BCF=2000 L/kg) and the vB criterion (BCF=5000 L/kg) according to REACH Annex XIII. In contrast, a field study reporting a bioaccumulation factor (BAF) far above 5000 L/kg indicates that Galaxolide might be even very bioaccumulative [45]. This finding is in line with the LIFE APEX data provided to the French chemical authority, where Galaxolide was detected in

58% of all pooled samples ( $n=67$ ) of four species (otter, seals, buzzards and fish) from Sweden, Germany, Netherlands and UK. High concentrations were detected in livers of buzzard (342  $\mu\text{g}/\text{kg}$ ), fish otters (3361  $\mu\text{g}/\text{kg}$ ) and harbour porpoises (3790  $\mu\text{g}/\text{kg}$ ), which highlights the ubiquitous distribution of the substance in the environment and may indicate an increased potential for biomagnification. Furthermore, LIFE APEX data were used in the restriction processes of per- and polyfluoroalkyl substances, to confirm environmental distribution behaviour and hazards, in particular PBT properties.

#### **Tool to determine typical chemical mixtures in top predators**

The combined chemical risk arising from exposure to chemical mixtures and transformation products is as yet hardly considered in European chemical risk assessment although new approaches are being developed [30–32]. LIFE APEX developed a mixture assessment tool ([www.norman-data.eu/apex](http://www.norman-data.eu/apex)) for occurrence data to visualise the co-exposure to various chemicals and to determine typical mixtures in apex predators. Moreover, the tool allows filtering of the chemicals by their frequency of appearance, species and matrix. The results are presented in an interactive and downloadable graph map.

#### **Statistical considerations regarding effectiveness evaluation**

The work in LIFE APEX with regard to effectiveness evaluations showed the extent to which raptor samples could be pooled while still allowing the detection of significant trends over time in contaminant concentrations following the introduction of risk management measures. This knowledge facilitates more cost-effective monitoring of the effectiveness of mitigation measures at the European scale.

#### **Development of guidance for use of chemical apex data in PBT screening and bioaccumulation assessments**

- Practical guidance documents and protocols were elaborated within LIFE APEX on the interpretation of field data from apex species, addressing uncertainties in PBT screening and bioaccumulation assessments. These guidance documents are provided online ([www.lifeapex.eu](http://www.lifeapex.eu)).
- Currently, results of LIFE APEX are being used in the revision process of the ECHA *Guidance on information requirements and chemical safety assessment*. Chapter R.11: PBT assessment [38] and R.7c: End-point specific guidance [39]. These ECHA guidelines have to be followed in registrations under REACH

and aim to aid industry to conduct and improve risk and hazard assessment as part of the registration process.

#### **Recommendations**

We propose a number of actions and advancements to allow detection and prioritisation of contaminants of emerging concern at an early stage, with the goal of timely risk management measures to protect humans and the environment from exposure to hazardous chemicals. We here focus on the ‘early warning system’ function, notwithstanding the importance of long-term monitoring efforts.

#### **Establish a European research infrastructure for the use of top predators as sentinels in early warning systems and in risk and hazard assessment**

Monitoring data are required for evidence-based approaches in support of EU Chemical Strategies for Sustainability Towards a Toxic-Free Environment. Over recent decades, experience has been mainly gained by the regulatory and scientific community with contaminant monitoring of water, sediment and soil, and mainly lower trophic marine biota. Today, raptors, terrestrial and marine mammals at the top of the food chain are available in research collections, natural history museums and some environmental specimen banks, to allow for pan-European contaminant monitoring in a complementary way [18, 23–25]. To this end, including top predator samples to monitor and screen for contaminants in freshwater, marine and in particular terrestrial ecosystems would be a cost- and resource-efficient solution to track levels of contaminants, to screen for contaminants of emerging concerns, chemical mixtures and transformation products, as well as to check the effectiveness of chemical regulations [5, 15]. This approach could support all chemical regulations, including those for industrial chemicals, plant protection products, biocides, and pharmaceuticals as well as WFD and MSFD.

Specifically, we recommend the following actions:

- Establish a pan-European research and monitoring infrastructure in support of chemical risk assessment that responds to regulatory needs for the systematic use of apex predator and other wildlife samples in chemical monitoring and early warning systems. A first step could be to incorporate top predators into the sampling regime of WFD and MSFD. As the detection of chemicals in apex predators reflects food web bioaccumulation and biomagnification, the chemical exposure at the top

of the food chain is also linked to human exposure potential. Experience exists with links to human exposure from Arctic monitoring of top predators and Inuit populations with a traditional subsistence lifestyle [46, 47]. This relates the call of the EC Chemicals Strategy for Sustainability Towards a Toxic-Free Environment for the development of an EU early warning tool to ensure that EU policies address emerging chemical risks as soon as identified by monitoring. By combining EU legal instruments for reducing contaminant releases to the environment and resulting exposures with the application of chemical monitoring in wildlife, a scientifically sound and more comprehensive basis for action can be developed.

- Making better use of targeted, suspect and non-target screening approaches using appropriate samples from top predator collections to respond to the needs of specific EU regulatory processes, for example:
  - a. To unravel inconsistencies in registration and approval data by matching exposure estimates with 'real world' levels in top predators.
  - b. To screen chemicals detected in apex species for classification as PBT compounds, endocrine disruptors and chemicals hazardous for human health (e.g., CMR), using *in silico* hazard screening tools such as JANUS (<https://www.vegahub.eu>) or the OECD toolbox (<https://qsartoolbox.org/>). This screening could support prioritisation [48] and decision-making processes for follow-up regulatory action for emerging compounds and chemical groups.
  - c. To provide evidence to better understand and assess biomagnification processes of chemicals, for example using samples from selected food webs [49] and/or matched samples of predator and prey.
  - d. To provide evidence on the occurrence of (bio)transformation products and typical chemical mixtures in top predators of different food webs, which can support advancements in risk modelling and assessment methods that are currently being developed with the ultimate goal to identify and manage hazardous chemicals more efficiently.
  - e. To establish retrospective time trends. Archived samples of apex species allow for retrospective analysis of contaminants of emerging concern as e.g. shown for Arctic top predators [50] and Scandinavian otters [51]. Retrospective time trends support the prioritisation of contaminants of emerging concern [48, 52]) and the effectiveness evaluation of management options for regulated compounds. Temporal data can

also be used to forecast future trends in relation to linear, cumulative and systemic approaches [27].

- f. To evaluate the predictability of current risk assessment schemes, and the effectiveness of chemical risk mitigation measures in particular by time trends analysis as e.g. shown for the Arctic [50, 53, 54].
- The proposed actions should build upon the considerable experiences and knowledge gained from the monitoring, modelling and assessment of chemical status by, e.g. dedicated efforts of ECHA, the European Food Safety Authority (EFSA) and EU Member States and international efforts in relation to the Stockholm Convention. To this end, the engagement of all relevant stakeholders including industry, the scientific community, sample providers, regulators and public representatives should be encouraged, to support these approaches and amendments and to enable their successful implementation. This can be achieved, for example, within the recently launched EU Partnership for the Assessment of Risks from Chemicals (PARC) [55], which will foster the collaboration of established working groups across the different chemical regulations.

#### **Develop specific guidance to support regulatory consideration of chemical data from apex predators**

- Current challenges, needs and perspectives from industry and regulators across regulations need to be systematically evaluated to identify where the use of biota/apex predator data can provide an added value (screening, prioritisation, assessment and management), which further approaches need to be developed, and how to incorporate them into current regulatory practices and assessment schemes. For instance, field bioaccumulation data are an ultimate example of real bioaccumulation potential and the impact of multiple releases from multiple sources.
- Recognising that field bioaccumulation data make an important contribution to the weight of evidence approach within assessments, in particular under REACH, plant protection products and biocides, we recommend further developing the section on the use of field data in the ECHA guidance [38, 39] and of the EFSA guidance on birds and mammals pesticides risk assessments [56], currently under update, to facilitate practical applications of chemical monitoring data from apex species for screening, assessment and management across EU-chemical

regulations. This could for instance be the work for a dedicated expert group in collaboration with PARC and the NORMAN network ([www.norman-network.com](http://www.norman-network.com), [55]).

- Screening, prioritisation and assessment of chemicals by means of advanced monitoring or modelling approaches require both expert knowledge and resources. Since not all chemical authorities and countries currently have these capacities, an increased cooperation and knowledge sharing on methods and procedures is advisable. With regard to REACH, this could be addressed within the regulatory exchange platforms of ECHA's expert groups and with industry, e.g. by promoting the use of monitoring databases before and after registration of a chemical. PARC could also provide a forum for capacity building, in close collaborations between research institutions and environmental authorities. In the area of plant protection products, this should lead to landscape exposure assessments, aggregating the exposure from different uses and developing the basis for combined exposure assessments. A clear opportunity is the extrapolation of the cumulative assessment groups, developed by EFSA for human health, to cover also apex predators.

## Conclusions and outlook

In summary, the following next steps are required for the integration of apex predators in surveillance, operational and investigative monitoring of legacy and emerging contaminants in support of EU risk and hazard assessments:

- Recognition that current concerns of exposures of humans and wildlife to legacy chemicals and contaminants of emerging concern, mixtures and (bio) transformation products require a more efficient use of existing resources, i.e. chemical monitoring data from apex and other wildlife species, which reflect 'real world' data.
- A centralised European data infrastructure is required which allows compilation, open-access and exchange of chemical monitoring data on wildlife as well as related regulatory information such as regulatory status, tonnages produced/ marketed and uses [57]. Such a data platform will foster collaborative efforts in data and knowledge sharing across different stakeholder groups. Storing raw target and non-target screening data converted into a common (open) format further allows for 'on demand' access to retrospective data analysis. The NORMAN platform could be used as central data platform and knowledge hub, hosting chemical moni-

toring data for prioritisation and risk assessment, as well as method information and guidance to ensure high quality and comparability of chemical monitoring data. Interoperability with the EU Information Platform for Chemical Monitoring (IPCHEM) and the European Marine Observation and Data Network (EMODnet) should be considered.

- The acknowledgment by regulators of the value of top predators as key sentinels for early warning as well as to monitor ecosystems and improve risk assessment and understanding of their impacts. Monitoring data are particularly valuable for developing landscape aggregated exposure models, complementing pre-market assessments.
- The development of baselines and comparable trend analysis based on spatial and temporal consistency of approaches, also supported by large-scale collaborative surveys.
- The development of coherent guidance documents should be initiated in collaborative approaches across European chemical regulations. These should address the practical application of chemical monitoring data from apex species in the screening, assessment and management of contaminants.

## Abbreviations

BCF: Bioconcentration factor; CMR: Cancerogenic, mutagenic and reprotoxic; EC: European Commission; ECHA: European Chemicals Agency; EFSA: European Food Safety Authority; EMODnet: European Marine Observation and Data Network; ESBs: Environmental specimen banks; EU: European Union; IPCHEM: Information Platform for Chemical Monitoring; MSFD: Marine Strategy Framework Directive; NHMs: Natural history museums; NORMAN: Network of reference laboratories, research centres and related organisations for monitoring of emerging environmental substances; OECD: Organisation for Economic Co-operation and Development; PARC: Partnership for the Assessment of Risks from Chemicals; PBT/vPvB: Persistent, Bioaccumulative and Toxic/very Persistent and very Bioaccumulative; POPs: Persistent organic pollutants; QSAR: Quantitative structure–activity relationship; REACH: Regulation on the Registration, Evaluation and Authorisation of Chemicals; UNEP: United Nations Environment Programme; WFD: Water Framework Directive.

## Acknowledgements

We thank Björn Hidding, Stephanie Bopp and all regulatory advisory board members for their valuable contribution to the LIFE APEX project (LIFE17 ENV/SK/000355). Furthermore, we would like to acknowledge all sample providers for their support of the project.

## Author contributions

GT has been responsible for the concept of the manuscript and drafted the manuscript. JK and KV helped to further elaborate the manuscript and contributed specific aspects. All authors read and approved the final manuscript.

## Funding

Open Access funding enabled and organized by Projekt DEAL. This study is financed by the European Union through the project LIFE17 ENV/SK/000355 'Systematic use of contaminant data from apex predators and their prey in chemicals management'.

**Availability of data and materials**

Not applicable; presented information is based on previously published data only.

**Declarations****Ethics approval and consent to participate**

Not applicable.

**Consent for publication**

Not applicable.

**Competing interests**

The authors declare that they have no competing interests.

**Author details**

<sup>1</sup>German Environment Agency (Umweltbundesamt), Wörlitzer Platz 1, 06813 Dessau-Roßlau, Germany. <sup>2</sup>Environmental Institute, Okružná 784/42, 97241 Koš, Slovak Republic. <sup>3</sup>National and Kapodistrian University of Athens, Panepistimiopolis Zographou, 15771 Athens, Greece. <sup>4</sup>Öko-Institut e.V, Institute for Applied Ecology, PO Box 1771, 79017 Freiburg, Germany. <sup>5</sup>Department of Chemistry "Ugo Schiff", University of Florence, Via delle Lastrucce 3, Florence, Italy. <sup>6</sup>Naturalis Biodiversity Center, Darwinweg 2, 2333 CR Leiden, Netherlands. <sup>7</sup>INERIS, National Institute for Environment and Industrial Risks, Verneuil en Halatte, France. <sup>8</sup>UK Centre for Ecology & Hydrology, Maclean Bldg, Benson Ln, Crowmarsh Gifford, Wallingford OX10 8BB, UK. <sup>9</sup>Fraunhofer Institute for Molecular Biology and Applied Ecology (Fraunhofer IME), 57392 Schmallenberg, Germany. <sup>10</sup>European Commission, Joint Research Centre (JRC), Ispra, Italy. <sup>11</sup>Norwegian Institute for Water Research (NIVA), Økernveien 94, 0579 Oslo, Norway. <sup>12</sup>Methodology and Scientific Support Unit, European Food Safety Authority (EFSA), Via Carlo Magno 1/A, 43126 Parma, Italy. <sup>13</sup>Department of Environmental Science, Aarhus University, Frederiksborgvej 399, 4000 Roskilde, Denmark. <sup>14</sup>UK Centre for Ecology & Hydrology, Lancaster Environment Centre, Lancaster LA1 4AP, UK.

Received: 21 July 2022 Accepted: 20 August 2022

Published online: 02 September 2022

**References**

- EC (2006) Regulation (EC) No 1907/2006 of the European Parliament and of the Council of 18 December 2006 concerning the Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH), establishing a European Chemicals Agency, amending Directive 1999/45/EC and repealing Council Regulation (EEC) No 793/93 and Commission Regulation (EC) No 1488/94 as well as Council Directive 76/769/EEC and Commission Directives 91/155/EEC, 93/67/EEC, 93/105/EC and 2000/21/EC
- Persson L, Carney Almroth BM, Collins CD, Cornell S, de Wit CA, Diamond ML et al (2022) Outside the safe operating space of the planetary boundary for novel entities. *Environ Sci Technol* 56:1510–1521
- UN (2019) United Nation Report: Nature's Dangerous Decline 'Unprecedented'; Species Extinction Rates 'Accelerating'. <https://www.un.org/sustainabledevelopment/blog/2019/05/nature-decline-unprecedented-report/>
- Chen Y, Fu J, Ye T, Li X, Gao K, Xue Q et al (2021) Occurrence, profiles, and ecotoxicity of poly- and perfluoroalkyl substances and their alternatives in global apex predators: a critical review. *J Environ Sci* 109:219–236
- Elliott JE, Elliott KH (2013) Tracking marine pollution. *Science* 340:556–558
- EC (2019) European Commission. Communication from the Commission to the European Parliament, the European Council, the Council, the European Economic and Social Committee and the Committee of the Regions. The European Green Deal. COM(2019) 640 final
- EC (2021) Communication from the commission to the European Parliament, the council, the European economic and social committee and the committee of the regions. Pathway to a Healthy Planet for All EU Action Plan: 'Towards Zero Pollution for Air, Water and Soil'
- EC (2020), European Commission (EC) - Chemicals Strategy for Sustainability - Towards a Toxic-Free Environment Communicaton from the Commission to the European Parliament, the Council, the European Economic and Social Committee and the Committee of the Regions 2020; COM(2020) 667 final
- Bierregaard RO Jr, Ben David A, Gibson L, Kennedy RS, Poole AF, Scheibel MS et al (2014) Post-DDT recovery of Osprey (*Pandion haliaetus*) populations in southern New England and Long Island, New York, 1970–2013. *J Raptor Res* 48:361–374
- Blus L, Heath R, Gish C, Belisle A, Prouty R (1971) Eggshell thinning in the brown pelican: implication of DDE. *Bioscience* 21:1213–1215
- Moermond CTA, Janssen MPM, de Knecht JA, Montforts HMM, Peijnenburg WJGM, Zweers PGPC et al (2012) PBT assessment using the revised annex XIII of REACH: a comparison with other regulatory frameworks. *Integr Environ Assess Manag* 8:359–371
- UNEP (2011) UN Environment Programme Stockholm Convention on Persistent Organic Pollutants. 2001. Annex D, p 35–36. [www.pops.int](http://www.pops.int). Accessed 20 April 2010
- EC (2000) Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for Community action in the field of water policy 2000.
- EC (2008) Directive 2008/56/EC of the European Parliament and of the Council of 17 June 2008 establishing a framework for community action in the field of marine environmental policy (Marine Strategy Framework Directive)
- Badry A, Krone O, Jaspers VLB, Mateo R, García-Fernández A, Leivits M et al (2020) Towards harmonisation of chemical monitoring using avian apex predators: identification of key species for pan-European biomonitoring. *Sci Total Environ* 731:139198
- Espín S, Andevski J, Duke G, Eulaers I, Gómez-Ramírez P, Hallgrímsson GT et al (2021) A schematic sampling protocol for contaminant monitoring in raptors. *Ambio* 50:95–100
- Gómez-Ramírez P, Shore RF, van den Brink NW, van Hattum B, Bustnes JO, Duke G et al (2014) An overview of existing raptor contaminant monitoring activities in Europe. *Environ Int* 67:12–21
- Ramello G, Duke G, Dekker RWRJ, van der Mije S, Movalli P (2022) A novel survey of raptor collections in Europe and their potential to provide samples for pan-European contaminant monitoring. *Environ Sci Pollut Res* 29:17017–17030
- Brack W, Escher BI, Müller E, Schmitt-Jansen M, Schulze T, Slobodnik J et al (2018) Towards a holistic and solution-oriented monitoring of chemical status of European water bodies: how to support the EU strategy for a non-toxic environment? *Environ Sci Eur* 30:33
- Brack W, Hollender J, de Alda ML, Müller C, Schulze T, Schymanski E et al (2019) High-resolution mass spectrometry to complement monitoring and track emerging chemicals and pollution trends in European water resources. *Environ Sci Eur* 31:62
- Hollender J, Schymanski EL, Singer HP, Ferguson PL (2017) Nontarget screening with high resolution mass spectrometry in the environment: ready to go? *Environ Sci Technol* 51:11505–11512
- Hollender J, van Bavel B, Dulio V, Farnen E, Furtmann K, Koschorreck J et al (2019) High resolution mass spectrometry-based non-target screening can support regulatory environmental monitoring and chemicals management. *Environ Sci Eur* 31:42
- Koizumi A, Harada KH, Inoue K, Hitomi T, Yang H-R, Moon C-S et al (2009) Past, present, and future of environmental specimen banks. *Environ Health Prev Med* 14:307–318
- Koschorreck J, Heiss C, Wellmitz J, Fliedner A, Rüdell H (2015) The use of monitoring data in EU chemicals management—experiences and considerations from the German environmental specimen bank. *Environ Sci Pollut Res* 22:1597–1611
- Movalli P, Duke G, Ramello G, Dekker R, Vrezec A, Shore RF et al (2019) Progress on bringing together raptor collections in Europe for contaminant research and monitoring in relation to chemicals regulation. *Environ Sci Pollut Res Int* 26:20132–20136
- Schäfer RB, Liess M, Altenburger R, Filser J, Hollert H, Roß-Nickoll M et al (2019) Future pesticide risk assessment: narrowing the gap between intention and reality. *Environ Sci Eur* 31:21
- EEA (2020) European Environment Agency (EEA). The European environment - state and outlook 2020. Knowledge for transition to a sustainable Europe. Luxembourg: Publications Office of the European Union 2019; Chapter 10 - Chemical pollution: DOI: <https://doi.org/10.2800/96749>



28. Springer A, Herrmann H, Sittner D, Herbst U, Schulte A (2015) REACH Compliance: Data Availability of REACH Registrations - Part 1: Screening of chemicals > 1000 tpa. ISSN 1862–4804. Editor: Neumann, M. Umweltbundesamt, Dessau-Roßlau
29. van den Brink NW, Arblaster JA, Bowman SR, Conder JM, Elliott JE, Johnson MS et al (2016) Use of terrestrial field studies in the derivation of bioaccumulation potential of chemicals. *Integr Environ Assess Manag* 12:135–145
30. Bopp SK, Kienzler A, Richarz AN, van der Linden SC, Paini A, Parisis N et al (2019) Regulatory assessment and risk management of chemical mixtures: challenges and ways forward. *Crit Rev Toxicol* 49:174–189
31. Galert W, Hassold E (2021) Environmental risk assessment of technical mixtures under the European registration, evaluation, authorisation and restriction of chemicals—a regulatory perspective. *Integr Environ Assess Manag* 17:498–506
32. Posthuma L, Brack W, van Gils J, Focks A, Müller C, de Zwart D et al (2019) Mixtures of chemicals are important drivers of impacts on ecological status in European surface waters. *Environ Sci Eur* 31:71
33. Wang Z, Walker GW, Muir DCG, Nagatani-Yoshida K (2020) Toward a global understanding of chemical pollution: a first comprehensive analysis of national and regional chemical inventories. *Environ Sci Technol* 54:2575–2584
34. Wilson MP, Schwarzman MR (2009) Toward a New US chemicals policy: rebuilding the foundation to advance new science, green chemistry, and environmental health. *Environ Health Perspectives* 117:1202–1209
35. Streißl F, Egsmose M, Tarazona JV (2018) Linking pesticide marketing authorisations with environmental impact assessments through realistic landscape risk assessment paradigms. *Ecotoxicology* 27:980–991
36. Topping CJ, Aldrich A, Beryn P (2020) Overhaul environmental risk assessment for pesticides. *Science* 367:360–363
37. Tarazona D, Tarazona G, Tarazona JV (2021) A simplified population-level landscape model identifying ecological risk drivers of pesticide applications, part one: case study for large herbivorous mammals. *Int J Environ Res Public Health* 18:7720
38. ECHA (2017) Guidance on Information Requirements and Chemical Safety Assessment. Chapter R.11: PBT/vPvB assessment. Version 3.0 – June 2017. [https://echa.europa.eu/documents/10162/13632/information\\_requirements\\_r11\\_en.pdf/a8cce23f-a65a-46d2-ac68-92fee1f9e54f](https://echa.europa.eu/documents/10162/13632/information_requirements_r11_en.pdf/a8cce23f-a65a-46d2-ac68-92fee1f9e54f). Assessed 4 July 2022.
39. ECHA (2017) Guidance on Information Requirements and Chemical Safety Assessment. Chapter R.7c: Endpoint specific guidance. Version 3.0 – June 2017. [https://echa.europa.eu/documents/10162/17224/information\\_requirements\\_r7c\\_en.pdf/e2e23a98-adb2-4573-b450-cc0df a7988e5?t=1498476107907](https://echa.europa.eu/documents/10162/17224/information_requirements_r7c_en.pdf/e2e23a98-adb2-4573-b450-cc0df a7988e5?t=1498476107907). Assessed 4 July 2022.
40. van Dijk J, Gustavsson M, Dekker SC, van Wezel AP (2021) Towards 'one substance – one assessment': an analysis of EU chemical registration and aquatic risk assessment frameworks. *J Environ Manage* 280:111692
41. Badry A, Slobodnik J, Alygizakis N, Bunke D, Cincinelli A, Claßen D, et al. (2022). Towards harmonised sampling and processing of archived wildlife samples to increase the regulatory uptake of monitoring data in chemicals management. *Environmental Sciences Europe* (accepted August 2022).
42. Pizzo F, Lombardo A, Brandt M, Manganaro A, Benfenati E (2016) A new integrated in silico strategy for the assessment and prioritization of persistence of chemicals under REACH. *Environ Int* 88:250–260
43. Pizzo F, Lombardo A, Manganaro A, Cappelli CI, Petoumenou MI, Albanese F et al (2016) Integrated in silico strategy for PBT assessment and prioritization under REACH. *Environ Res* 151:478–492
44. EC (2008) European Commission. European Union Risk Assessment Report on 1,3,4,6,7,8-HEXAHYDRO-4,6,6,7,8-HEXAMETHYLCYCLOPENTA-γ-2-BENZOPYRAN. <https://echa.europa.eu/documents/10162/947def3b-bbbf-473b-bc19-3bda7a8da910>. Assessed 20 June 2022.
45. Yao L, Zhao JL, Liu YS, Zhang QQ, Jiang YX, Liu S et al (2018) Personal care products in wild fish in two main Chinese rivers: bioaccumulation potential and human health risks. *Sci Total Environ* 621:1093–1102
46. Gibson J, Adlard B, Olafsdottir K, Sandanger TM, Odland J (2016) Levels and trends of contaminants in humans of the Arctic. *Int J Circumpolar Health* 75:33804
47. Sonne C (2010) Health effects from long-range transported contaminants in Arctic top predators: an integrated review based on studies of polar bears and relevant model species. *Environ Int* 36:461–491
48. Badry A, Treu G, Gkotsis G, Nika M-C, Alygizakis N, Thomaidis NS et al (2022) Ecological and spatial variations of legacy and emerging contaminants in white-tailed sea eagles from Germany: implications for prioritisation and future risk management. *Environ Int* 158:106934
49. Kosfeld V, Rüdell H, Schlechtriem C, Rauer C, Koschorreck J (2021) Food web on ice: a pragmatic approach to investigate the trophic magnification of chemicals of concern. *Environ Sci Eur* 33:93
50. Rigét F, Vorkamp K, Bossi R, Sonne C, Letcher RJ, Dietz R (2016) Twenty years of monitoring of persistent organic pollutants in Greenland biota. A review. *Environ Pollut* 217:114–123
51. Roos A, Berger U, Järnberg U, van Dijk J, Bignert A (2013) Increasing Concentrations of Perfluoroalkyl Acids in Scandinavian Otters (*Lutra lutra*) between 1972 and 2011: a new threat to the Otter Population? *Environ Sci Technol* 47:11757–11765
52. Dürig W, Alygizakis NA, Menger F, Golovko O, Wiberg K, Ahrens L (2022) Novel prioritisation strategies for evaluation of temporal trends in archived white-tailed sea eagle muscle tissue in non-target screening. *J Hazard Mater* 424, 127331. <https://doi.org/10.1016/j.jhazmat.2021.127331>
53. Rigét F, Bignert A, Braune B, Dam M, Dietz R, Evans M et al (2019) Temporal trends of persistent organic pollutants in Arctic marine and freshwater biota. *Sci Total Environ* 649:99–110
54. Wong F, Hung H, Dryfhout-Clark H, Aas W, Bohlin-Nizzetto P, Breivik K et al (2021) Time trends of persistent organic pollutants (POPs) and Chemicals of Emerging Arctic Concern (CEAC) in Arctic air from 25 years of monitoring. *Sci Total Environ* 775:145109
55. Dulio V, Koschorreck J, van Bavel B, van den Brink P, Hollender J, Munthe J et al (2020) The NORMAN Association and the European Partnership for Chemicals Risk Assessment (PARC): let's cooperate! *Environ Sci Eur* 32:100
56. European Food Safety Authority (2009) Guidance document on risk assessment for birds & mammals on request from EFSA. *EFSA J* 7(12):1438. <https://doi.org/10.2903/j.efsa.2009.1438>
57. Slobodnik J, Hollender J, Schulze T, Schymanski EL, Brack W (2019) Establish data infrastructure to compile and exchange environmental screening data on a European scale. *Environ Sci Eur* 31:65

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Submit your manuscript to a SpringerOpen® journal and benefit from:**

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)