



DATA NOTE

REVISED The genome sequence of the Dusky Thorn moth,***Ennomos fuscantarius* (Haworth, 1809)**

[version 2; peer review: 2 approved, 1 approved with reservations]

Previously titled: "The genome sequence of the Dusky Thorn moth, *Ennomos fuscantarius* (Haworth, 1809)"

Douglas Boyes¹⁺, Dominic Phillips^{id}²,
 University of Oxford and Wytham Woods Genome Acquisition Lab,
 Darwin Tree of Life Barcoding collective,
 Wellcome Sanger Institute Tree of Life programme,
 Wellcome Sanger Institute Scientific Operations: DNA Pipelines collective,
 Tree of Life Core Informatics collective, Darwin Tree of Life Consortium

¹UK Centre for Ecology & Hydrology, Wallingford, England, UK²Natural History Museum, London, England, UK

+ Deceased author

v2 First published: 07 Nov 2023, 8:505
<https://doi.org/10.12688/wellcomeopenres.20174.1>

Latest published: 14 Feb 2026, 8:505
<https://doi.org/10.12688/wellcomeopenres.20174.2>

Abstract

We present a genome assembly from an individual male *Ennomos fuscantarius* (the Dusky Thorn; Arthropoda; Insecta; Lepidoptera; Geometridae). The genome sequence is 444.9 megabases in span. Most of the assembly is scaffolded into 31 chromosomal pseudomolecules, including the Z sex chromosome. The mitochondrial genome has also been assembled and is 15.49 kilobases in length. Gene annotation of this assembly on Ensembl identified 12,173 protein coding genes.

Keywords

Ennomos fuscantarius, Dusky Thorn, genome sequence, chromosomal, Lepidoptera



This article is included in the [Tree of Life gateway](#).

Open Peer Review**Approval Status** ✓ ? ✓

	1	2	3
version 2 (revision) 14 Feb 2026			✓ view
version 1 07 Nov 2023	✓ view	? view	

1. **Arjen Van 't Hof** ^{id}, Biology Centre of the Czech Academy of Sciences, Budějovice, Czech Republic
2. **Saskia Wutke** ^{id}, University of Eastern Finland, Joensuu, Finland
3. **Stephan Koblmüller** ^{id}, University of Graz, Graz, Austria

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Darwin Tree of Life Consortium (mark.blaxter@sanger.ac.uk)

Author roles: **Boyes D:** Investigation, Resources; **Phillips D:** Writing – Original Draft Preparation;

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by Wellcome through core funding to the Wellcome Sanger Institute (206194) and the Darwin Tree of Life Discretionary Award (218328).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2026 Boyes D *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Boyes D, Phillips D, University of Oxford and Wytham Woods Genome Acquisition Lab *et al.* **The genome sequence of the Dusky Thorn moth, *Ennomos fuscantarius* (Haworth, 1809) [version 2; peer review: 2 approved, 1 approved with reservations]** Wellcome Open Research 2026, 8:505 <https://doi.org/10.12688/wellcomeopenres.20174.2>

First published: 07 Nov 2023, 8:505 <https://doi.org/10.12688/wellcomeopenres.20174.1>

REVISED Amendments from Version 1

Version 2 of this data note incorporates corrections requested by peer reviewers. We have also updated the results of the Mercury.FK analysis of both primary and alternate assemblies.

Any further responses from the reviewers can be found at the end of the article

Species taxonomy

Eukaryota; Metazoa; Eumetazoa; Bilateria; Protostomia; Ecdysozoa; Panarthropoda; Arthropoda; Mandibulata; Pancrustacea; Hexapoda; Insecta; Dicondylia; Pterygota; Neoptera; Endopterygota; Amphimesenoptera; Lepidoptera; Glossata; Neolepidoptera; Heteroneura; Ditrysia; Obtectomera; Geometroidea; Geometridae; Ennominae; *Ennomos*; *Ennomos fuscantaria* (Haworth, 1809) (NCBI:txid722662).

Background

Ennomos fuscantaria (Dusky Thorn) is a Geometrid moth in the Ennominae subfamily with a forewing length of 17–21 mm; a greyish or mauve-shaded band between the forewing margin and distal cross-line distinguishes it from similar species (Lewis, 2023; Waring *et al.*, 2017).

E. fuscantaria is a monovoltine species, flying from late July to early October. Though it readily comes to light, it is not often seen out on the wing (Randle *et al.*, 2019; Waring *et al.*, 2017). The larva feeds predominantly on ash (*Fraxinus excelsior*), on which it overwinters as an egg, later to pupate within spun leaves. It can be coaxed to feed on privets (*Ligustrum* spp.) (Waring *et al.*, 2017).

E. fuscantaria can be found in most habitats where ash trees are present; in the UK, it has a widespread distribution, frequently occurring through England and Wales (Randle *et al.*, 2019). Globally, *E. fuscantaria* is distributed throughout the western Palearctic, extending from western Europe to eastern Russia and the Mediterranean (GBIF Secretariat, 2023).

Numbers of *E. fuscantaria* are prone to fluctuate in the UK and are currently present on the Rothamstead Red List, with suspected population numbers decreasing by of 47% over a 10-year period as of 2019, with earlier research showing a 90% decrease between 1990 and 2001 (GBIF Secretariat, 2023; Palmer *et al.*, 2015; Waring *et al.*, 2017). This species is highly sensitive to the progression of climate change, as demonstrate by the large decrease in population linked to increased rates of ash dieback (Fox *et al.*, 2021; Palmer *et al.*, 2015). The genome sequence of *E. fuscantaria* will provide insights into how we can help to reduce these impacts that changes in climate can cause to occupancy and distribution of this vulnerable species. Sequence data will also support phylogenetic analyses of the relationships between Ennominae moths and studies of the evolution of flightlessness within the genus (Wahlberg *et al.*, 2010).

The genome of *Ennomos fuscantarius* was sequenced as part of the Darwin Tree of Life Project, a collaborative effort to sequence all named eukaryotic species in the Atlantic Archipelago of Britain and Ireland. Here we present a chromosomally complete genome sequence for *Ennomos fuscantarius*, based on one specimen collected by Douglas Boyes from Wytham Woods, Oxfordshire.

Genome sequence report

The genome was sequenced from one male *Ennomos fuscantarius* (Figure 1) collected from Wytham Woods, UK (51.77, -1.34). A total of 52-fold coverage in Pacific Biosciences single-molecule HiFi long reads and 85-fold coverage in 10X Genomics read clouds were generated. Primary assembly contigs were scaffolded with chromosome conformation Hi-C data. Manual assembly curation corrected 12 missing joins or mis-joins, reducing the scaffold number by 13.95%, and increasing the scaffold N50 by 1.61%.

The final assembly has a total length of 444.9 Mb in 36 sequence scaffolds with a scaffold N50 of 15.9 Mb (Table 1). A summary of the assembly statistics is shown in Figure 2, while the distribution of assembly scaffolds on GC proportion and coverage is shown in Figure 3. The cumulative assembly plot in Figure 4 shows curves for subsets of scaffolds assigned to different phyla. Most (99.95%) of the assembly sequence was assigned to 31 chromosomal-level scaffolds, representing 30 autosomes and the Z sex chromosome. Chromosome-scale scaffolds confirmed by the Hi-C data are named in order of size (Figure 5; Table 2). While not fully phased, the assembly deposited is of one haplotype. Contigs corresponding to the second haplotype have also been deposited. The mitochondrial genome was also assembled and can be found as a contig within the multifasta file of the genome submission.



Figure 1. Photograph of the *Ennomos fuscantarius* (iEnnFusc2) specimen used for genome sequencing.

The combined primary and alternate assemblies achieve an estimated QV of 55.0. The k -mer completeness is 96.74% for the primary assembly, 81.22% for the alternate haplotype, and 99.33% for the combined assemblies. The primary assembly has a BUSCO v5.3.2 completeness of 98.5% (single = 98.1%, duplicated = 0.4%), using the lepidoptera_odb10 reference set ($n = 5,286$).

Metadata for specimens, spectral estimates, sequencing runs, contaminants and pre-curation assembly statistics can be found at <https://links.tol.sanger.ac.uk/species/722662>.

Genome annotation report

The *Ennomos fuscantarius* genome assembly (GCA_905220475.3) was annotated using the Ensembl rapid annotation pipeline. The resulting annotation includes 23,475 transcribed mRNAs from 12,173 protein-coding and 1,740 non-coding genes. Further details of this annotation are available from the [Ensembl annotation page](#).

Methods

Sample acquisition and nucleic acid extraction

A male *Ennomos fuscantarius* (specimen ID Ox000200, individual iEnnFusc2) was collected from Wytham Woods, Oxfordshire (biological vice-county Berkshire), UK (latitude 51.77, longitude -1.34) on 2019-08-24 using a light trap. The specimen was collected and identified by Douglas Boyes (University of Oxford) and preserved on dry ice.

DNA was extracted at the Tree of Life laboratory, Wellcome Sanger Institute (WSI). The iEnnFusc2 sample was weighed and dissected on dry ice with tissue set aside for Hi-C sequencing. Head and thorax tissue was disrupted using a Nippi Powermasher fitted with a BioMasher pestle. High molecular weight (HMW) DNA was extracted using the Qiagen MagAttract HMW DNA extraction kit. Low molecular weight DNA was removed from a 20 ng aliquot of extracted DNA using the 0.8X AMPure XP purification kit prior to 10X Chromium sequencing; a minimum of 50 ng DNA was submitted for 10X sequencing. HMW DNA was sheared into an average fragment size of 12–20 kb in a Megaruptor 3 system with speed setting 30. Sheared DNA was purified by solid-phase reversible immobilisation using AMPure PB beads with a 1.8X ratio of beads to sample to remove the shorter fragments and concentrate the DNA sample. The concentration of the sheared and purified DNA was assessed using a Nanodrop spectrophotometer and Qubit Fluorometer and Qubit dsDNA High Sensitivity Assay kit. Fragment size distribution was evaluated by running the sample on the FemtoPulse system.

RNA was extracted from whole organism tissue of iEnnFusc1 in the Tree of Life Laboratory at the WSI using TRIzol, according to the manufacturer's instructions. RNA was then eluted in 50 μ l RNase-free water and its concentration assessed using a Nanodrop spectrophotometer and Qubit Fluorometer using the Qubit RNA Broad-Range (BR) Assay kit. Analysis of the integrity of the RNA was done using Agilent RNA 6000 Pico Kit and Eukaryotic Total RNA assay.

Table 1. Genome data for *Ennomos fuscantarius*, iEnnFusc2.3.

Project accession data		
Assembly identifier	iEnnFusc2.3	
Species	<i>Ennomos fuscantarius</i>	
Specimen	iEnnFusc2	
NCBI taxonomy ID	722662	
BioProject	PRJEB42951	
BioSample ID	SAMEA7520185	
Isolate information	iEnnFusc2: male: head and thorax (PacBio HiFi), abdomen (Hi-C scaffolding) iEnnFusc1: whole organism (RNA sequencing)	
Assembly metrics*		Benchmark
Consensus quality (QV)	Primary: 56.3; alternate: 54.4; combined: 55.0	≥ 40
k-mer completeness	Primary: 96.74%; alternate: 81.22%; combined: 99.33%	≥ 95%
BUSCO**	C:98.5%[S:98.1%,D:0.4%],F:0.5%,M:1.1%,n:5,286	C ≥ 95%
Percentage of assembly mapped to chromosomes	99.95%	≥ 90%
Sex chromosomes	Z chromosome	<i>localised homologous pairs</i>
Organelles	Mitochondrial genome assembled	<i>complete single alleles</i>
Raw data accessions		
PacificBiosciences SEQUEL II	ERR6560797	
10X Genomics Illumina	ERR6054405, ERR6054407, ERR6054408, ERR6054406	
Hi-C Illumina	ERR6054409	
PolyA RNA-Seq Illumina	ERR6054410, ERR6787420	
Genome assembly		
Assembly accession	GCA_905220475.3	
Accession of alternate haplotype	GCA_905220485.1	
Span (Mb)	444.9	
Number of contigs	46	
Contig N50 length (Mb)	15.5	
Number of scaffolds	36	
Scaffold N50 length (Mb)	15.9	
Longest scaffold (Mb)	19.5	
Genome annotation		
Number of protein-coding genes	12,173	
Number of non-coding genes	1,740	
Number of gene transcripts	23,475	

*Assembly metric benchmarks are adapted from column VGP-2020 of "Table 1: Proposed standards and metrics for defining genome assembly quality" from (Rhie *et al.*, 2021).

**BUSCO scores based on the lepidoptera_odb10 BUSCO set using v5.3.2. C = complete [S = single copy, D = duplicated], F = fragmented, M = missing, n = number of orthologues in comparison. A full set of BUSCO scores is available at <https://blobtoolkit.genomehubs.org/view/iEnnFusc2.3/dataset/CAJMZZ03/busco>.

Sequencing

Pacific Biosciences HiFi circular consensus and 10X Genomics read cloud DNA sequencing libraries were constructed according to the manufacturers' instructions. Poly(A) RNA-Seq libraries were constructed using the NEB Ultra II RNA Library Prep kit. DNA and RNA sequencing was performed by the Scientific Operations core at the WSI on Pacific




Figure 2. Genome assembly of *Ennomos fuscantarius*, ilEnnFusc2.3: metrics. The BlobToolKit Snailplot shows N50 metrics and BUSCO gene completeness. The main plot is divided into 1,000 size-ordered bins around the circumference with each bin representing 0.1% of the 444,925,525 bp assembly. The distribution of scaffold lengths is shown in dark grey with the plot radius scaled to the longest scaffold present in the assembly (19,450,406 bp, shown in red). Orange and pale-orange arcs show the N50 and N90 scaffold lengths (15,884,173 and 11,089,215 bp), respectively. The pale grey spiral shows the cumulative scaffold count on a log scale with white scale lines showing successive orders of magnitude. The blue and pale-blue area around the outside of the plot shows the distribution of GC, AT and N percentages in the same bins as the inner plot. A summary of complete, fragmented, duplicated and missing BUSCO genes in the lepidoptera_odb10 set is shown in the top right. An interactive version of this figure is available at <https://blobtoolkit.genomehubs.org/view/ilEnnFusc2.3/dataset/CAJMZZ03/snail>.

Biosciences SEQUEL II (HiFi), Illumina HiSeq 4000, Illumina HiSeq 4000 (RNA-Seq) and HiSeq X Ten (10X) instruments. Hi-C data were also generated from abdomen tissue of ilEnnFusc2 using the Arima2 kit and sequenced on the HiSeq X Ten instrument.

Genome assembly, curation and evaluation

Assembly was carried out with Hifiasm (Cheng *et al.*, 2021) and haplotypic duplication was identified and removed with purge_dups (Guan *et al.*, 2020). One round of polishing was performed by aligning 10X Genomics read data to the assembly with Long Ranger ALIGN, calling variants with FreeBayes (Garrison & Marth, 2012). The assembly was then scaffolded with Hi-C data (Rao *et al.*, 2014) using SALSA2 (Ghurye *et al.*, 2019). The assembly was checked for contamination and corrected using the gEVAL system (Chow *et al.*, 2016) as described previously (Howe *et al.*, 2021). Manual curation was performed using gEVAL, HiGlass (Kerpedjiev *et al.*, 2018) and Pretext (Harry, 2022). The mitochondrial genome was assembled using MitoHiFi (Uliano-Silva *et al.*, 2023), which runs MitoFinder (Allio *et al.*, 2020) or MITOS (Bernt *et al.*, 2013) and uses these annotations to select the final mitochondrial contig and to ensure the general quality of the sequence.




Figure 3. Genome assembly of *Ennomos fuscantarius*, ilEnnFusc2.3: BlobToolKit GC-coverage plot. Scaffolds are coloured by phylum. Circles are sized in proportion to scaffold length. Histograms show the distribution of scaffold length sum along each axis. An interactive version of this figure is available at <https://blobtoolkit.genomehubs.org/view/ilEnnFusc2.3/dataset/CAJMZZ03/blob>.

A Hi-C map for the final assembly was produced using bwa-mem2 (Vasimuddin *et al.*, 2019) in the Cooler file format (Abdennur & Mirny, 2020). To assess the assembly metrics, the k -mer completeness and QV consensus quality values were calculated in Merqury.FK (Rhie *et al.*, 2020). The genome was analysed within the BlobToolKit environment (Challis *et al.*, 2020) and BUSCO scores (Manni *et al.*, 2021; Simão *et al.*, 2015) were calculated.

Table 3 contains a list of relevant software tool versions and sources.

Genome annotation

The Ensembl gene annotation system (Aken *et al.*, 2016) was used to generate annotation for the *Ennomos fuscantarius* assembly (GCA_905220475.3). Annotation was created primarily through alignment of transcriptomic data to the genome, with gap filling via protein-to-genome alignments of a select set of proteins from UniProt (UniProt Consortium, 2019).

Wellcome Sanger Institute – Legal and Governance

The materials that have contributed to this genome note have been supplied by a Darwin Tree of Life Partner. The submission of materials by a Darwin Tree of Life Partner is subject to the ‘**Darwin Tree of Life Project Sampling Code of Practice**’, which can be found in full on the Darwin Tree of Life website [here](https://www.darwintreeoflife.org/). By agreeing with and signing up to the Sampling Code of Practice, the Darwin Tree of Life Partner agrees they will meet the legal and ethical requirements and standards set out within this document in respect of all samples acquired for, and supplied to, the Darwin Tree of Life Project.

Further, the Wellcome Sanger Institute employs a process whereby due diligence is carried out proportionate to the nature of the materials themselves, and the circumstances under which they have been/are to be collected and provided for use. The purpose of this is to address and mitigate any potential legal and/or ethical implications of receipt and use of the




Figure 4. Genome assembly of *Ennomos fuscantarius*, iEnnFusc2.3: BlobToolKit cumulative sequence plot. The grey line shows cumulative length for all scaffolds. Coloured lines show cumulative lengths of scaffolds assigned to each phylum using the buscogenes taxrule. An interactive version of this figure is available at <https://blobtoolkit.genomehubs.org/view/iEnnFusc2.3/dataset/CAJMZZ03/cumulative>.




Figure 5. Genome assembly of *Ennomos fuscantarius*, iEnnFusc2.3: Hi-C contact map of the iEnnFusc2.3 assembly, visualised using HiGlass. Chromosomes are shown in order of size from left to right and top to bottom. An interactive version of this figure may be viewed at <https://genome-note-higlass.tol.sanger.ac.uk/?d=KGN1bRa6-S8ahRJaRSp9yrA>.

Table 2. Chromosomal pseudomolecules in the genome assembly of *Ennomos fuscantarius*, iEnnFusc2.

INSDC accession	Chromosome	Length (Mb)	GC%
HG992024.1	1	18.42	37.0
HG992025.1	2	18.21	37.0
HG992026.1	3	17.84	37.0
HG992027.1	4	17.76	37.0
HG992028.1	5	17.17	36.5
HG992029.1	6	16.98	36.5
HG992030.1	7	16.84	36.5
HG992031.1	8	16.68	36.5
HG992032.1	9	16.42	36.5
HG992033.1	10	16.09	37.0
HG992034.1	11	16.07	36.5
HG992035.1	12	15.88	37.0
HG992036.1	13	15.7	36.5
HG992037.1	14	15.53	37.0
HG992038.1	15	15.19	37.0
HG992039.1	16	15.06	37.0
HG992040.1	17	14.83	37.0
HG992042.1	19	14.29	37.0
HG992041.1	18	14.29	37.5
HG992043.1	20	14.18	37.0
HG992044.1	21	13.83	37.5
HG992045.1	22	11.57	37.5
HG992046.1	23	11.43	37.0
HG992047.1	24	11.38	37.0
HG992048.1	25	11.09	37.5
HG992049.1	26	10.57	37.5
HG992050.1	27	8.69	38.0
HG992051.1	28	7.82	38.5
HG992052.1	29	7.77	37.5
HG992053.1	30	7.68	38.0
HG992023.1	Z	19.45	36.5
HG992054.3	MT	0.02	19.5

Table 3. Software tools: versions and sources.

Software tool	Version	Source
BlobToolKit	4.1.7	https://github.com/blobtoolkit/blobtoolkit
BUSCO	5.3.2	https://gitlab.com/ezlab/busco
FreeBayes	1.3.1-17-gaa2ace8	https://github.com/freebayes/freebayes
gEVAL	N/A	https://geval.org.uk/
Hifiasm	0.12	https://github.com/chhylp123/hifiasm

Table 3. *Continued*

Software tool	Version	Source
HiGlass	1.11.6	https://github.com/higlass/higlass
Long Ranger ALIGN	2.2.2	https://support.10xgenomics.com/genome-exome/software/pipelines/latest/advanced/other-pipelines
Mercury	1.1.2	https://github.com/thegenemyers/MERQURY.FK
MitoHiFi	1	https://github.com/marcelauliano/MitoHiFi
PretextView	0.2	https://github.com/wtsi-hpag/PretextView
purge_dups	1.2.3	https://github.com/dfguan/purge_dups
SALSA	2.2	https://github.com/salsa-rs/salsa
sanger-tol/ genomenote	v1.0	https://github.com/sanger-tol/genomenote
sanger-tol/ readmapping	1.1.0	https://github.com/sanger-tol/readmapping/tree/1.1.0

materials as part of the research project, and to ensure that in doing so we align with best practice wherever possible. The overarching areas of consideration are:

- Ethical review of provenance and sourcing of the material
- Legality of collection, transfer and use (national and international)

Each transfer of samples is further undertaken according to a Research Collaboration Agreement or Material Transfer Agreement entered into by the Darwin Tree of Life Partner, Genome Research Limited (operating as the Wellcome Sanger Institute), and in some circumstances other Darwin Tree of Life collaborators.

Data availability

European Nucleotide Archive: *Ennomos fuscantarius* (dusky thorn). Accession number PRJEB42951; <https://identifiers.org/ena.embl/PRJEB42951>. (Wellcome Sanger Institute, 2021) The genome sequence is released openly for reuse. The *Ennomos fuscantarius* genome sequencing initiative is part of the Darwin Tree of Life (DToL) project. All raw sequence data and the assembly have been deposited in INSDC databases. Raw data and assembly accession identifiers are reported in Table 1.

Author information

Members of the University of Oxford and Wytham Woods Genome Acquisition Lab are listed here: <https://doi.org/10.5281/zenodo.4789928>.

Members of the Darwin Tree of Life Barcoding collective are listed here: <https://doi.org/10.5281/zenodo.4893703>.

Members of the Wellcome Sanger Institute Tree of Life programme are listed here: <https://doi.org/10.5281/zenodo.4783585>.

Members of Wellcome Sanger Institute Scientific Operations: DNA Pipelines collective are listed here: <https://doi.org/10.5281/zenodo.4790455>.

Members of the Tree of Life Core Informatics collective are listed here: <https://doi.org/10.5281/zenodo.5013541>.

Members of the Darwin Tree of Life Consortium are listed here: <https://doi.org/10.5281/zenodo.4783558>.

References

- Abdennur N, Mirny LA: **Cooler: Scalable storage for Hi-C data and other genomically labeled arrays.** *Bioinformatics.* 2020; **36**(1): 311–316.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Aken BL, Ayling S, Barrell D, *et al.*: **The Ensembl gene annotation system.** *Database (Oxford).* 2016; **2016**: baw093.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Allio R, Schomaker-Bastos A, Romiguier J, *et al.*: **MitoFinder: Efficient automated large-scale extraction of mitogenomic data in target enrichment phylogenomics.** *Mol Ecol Resour.* 2020; **20**(4): 892–905.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bernt M, Donath A, Jühling F, *et al.*: **MITOS: Improved *de novo* metazoan mitochondrial genome annotation.** *Mol Phylogenet Evol.* 2013; **69**(2): 313–319.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Challis R, Richards E, Rajan J, *et al.*: **BlobToolKit - interactive quality assessment of genome assemblies.** *G3 (Bethesda).* 2020; **10**(4): 1361–1374.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cheng H, Concepcion GT, Feng X, *et al.*: **Haplotype-resolved *de novo* assembly using phased assembly graphs with hifiasm.** *Nat Methods.* 2021; **18**(2): 170–175.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Chow A, Brugger K, Caccamo M, *et al.*: **gEVAL — a web-based browser for evaluating genome assemblies.** *Bioinformatics.* 2016; **32**(16): 2508–2510.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Di Tommaso P, Chatzou M, Floden EW, *et al.*: **Nextflow enables reproducible computational workflows.** *Nat Biotechnol.* 2017; **35**(4): 316–319. 28398311
[Publisher Full Text](#)
- Fox R, Dennis EB, Harrower CA, *et al.*: 2021; *The State of Britain's Larger Moths 2021*. Wareham, Dorset, UK.
[Reference Source](#)
- Garrison E, Marth G: *Haplotype-based variant detection from short-read sequencing.* 2012; [Accessed 26 July 2023].
[Publisher Full Text](#)
- GBIF Secretariat: **Ennomos fuscantaria (Haworth, 1809).** *GBIF Backbone Taxonomy.* 2023; [Accessed 15 September 2023].
- Ghurye J, Rhie A, Walenz BP, *et al.*: **Integrating Hi-C links with assembly graphs for chromosome-scale assembly.** *PLoS Comput Biol.* 2019; **15**(8): e1007273.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Guan D, McCarthy SA, Wood J, *et al.*: **Identifying and removing haplotypic duplication in primary genome assemblies.** *Bioinformatics.* 2020; **36**(9): 2896–2898.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Harry E: **PretextView (Paired REad TEXTure Viewer): A desktop application for viewing pretext contact maps.** 2022; [Accessed 19 October 2022].
[Reference Source](#)
- Howe K, Chow W, Collins J, *et al.*: **Significantly improving the quality of genome assemblies through curation.** *GigaScience.* Oxford University Press; 2021; **10**(1): giaa153. 33420778
[Publisher Full Text](#) | [Free Full Text](#)
- Kerpedjiev P, Abdennur N, Lekschas F, *et al.*: **HiGlass: web-based visual exploration and analysis of genome interaction maps.** *Genome Biol.* 2018; **19**(1): 125.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lewis C: **Ennomos fuscantaria (Dusky Thorn).** *British Lepidoptera.* 2023; [Accessed 21 September 2023].
[Reference Source](#)
- Manni M, Berkeley MR, Seppey M, *et al.*: **BUSCO update: Novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes.** *Mol Biol Evol.* 2021; **38**(10): 4647–4654.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Palmer G, Hill JK, Brereton TM, *et al.*: **Individualistic sensitivities and exposure to climate change explain variation in species' distribution and abundance changes.** *Sci Adv.* 2015; **1**(9): e1400220.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Randle Z, Evans-Hill LJ, Parsons MS, *et al.*: *Atlas of Britain & Ireland's Larger Moths.* Newbury: NatureBureau; 2019.
- Rao SSP, Huntley MH, Durand NC, *et al.*: **A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping.** *Cell.* 2014; **159**(7): 1665–1680.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rhie A, McCarthy SA, Fedrigo O, *et al.*: **Towards complete and error-free genome assemblies of all vertebrate species.** *Nature.* 2021; **592**(7856): 737–746.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rhie A, Walenz BP, Koren S, *et al.*: **Merqury: Reference-free quality, completeness, and phasing assessment for genome assemblies.** *Genome Biol.* 2020; **21**(1): 245.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Simão FA, Waterhouse RM, Ioannidis P, *et al.*: **BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs.** *Bioinformatics.* 2015; **31**(19): 3210–3212.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Uliano-Silva M, Ferreira JGRN, Krashenninnikova K, *et al.*: **MitoHiFi: a python pipeline for mitochondrial genome assembly from PacBio high fidelity reads.** *BMC Bioinformatics.* 2023; **24**(1): 288.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- UniProt Consortium: **UniProt: a worldwide hub of protein knowledge.** *Nucleic Acids Res.* 2019; **47**(D1): D506–D515.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Vasimuddin M, Misra S, Li H, *et al.*: **Efficient Architecture-Aware Acceleration of BWA-MEM for Multicore Systems.** IEEE International Parallel and Distributed Processing Symposium (IPDPS). *IEEE* 2019; **2019**: 314–324.
[Publisher Full Text](#)
- Wahlberg N, Snäll N, Viidalepp J, *et al.*: **The evolution of female flightlessness among Ennominae of the Holarctic forest zone (Lepidoptera, Geometridae).** *Mol Phylogenet Evol.* 2010; **55**(3): 929–938.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Waring P, Townsend M, Lewington R: **Field Guide to the Moths of Great Britain and Ireland: Third Edition.** Bloomsbury Wildlife Guides. 2017.
- Wellcome Sanger Institute: **The genome sequence of the Dusky Thorn, Ennomos fuscantarius (Haworth, 1809).** *European Nucleotide Archive. [dataset]*, accession number PRJEB42951, 2021.

Open Peer Review

Current Peer Review Status:   

Version 2

Reviewer Report 26 February 2026

<https://doi.org/10.21956/wellcomeopenres.27653.r148735>

© 2026 Koblmüller S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Stephan Koblmüller 

University of Graz, Graz, Graz, Austria

In this manuscript, Boyes et al. report the whole-genome assembly of the dusky thorn moth, *Ennomos fuscantarius*, a geometrid species broadly distributed throughout Europe. The authors provide useful background information on the species, which I found quite valuable. The justification for sequencing this species is clearly explained. The data are presented in a clear and organized manner, and the methodological approaches are described in sufficient detail. I have no concerns regarding the validity of the data or the results. As far as I can tell, the associated datasets have been made available in publicly accessible repositories.

Is the rationale for creating the dataset(s) clearly described?

Yes

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: phylogenetics/-genomics, population genetics, phylogeography

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 30 April 2024

<https://doi.org/10.21956/wellcomeopenres.22337.r78060>

© 2024 Wutke S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Saskia Wutke**

University of Eastern Finland, Joensuu, Finland

This data note reports the genome of the Dusky Thorn (*Ennomos fuscantaria*), a geometrid moth. The genome was assembled based on PacBio HiFi and 10X Genomics data, scaffolded with Hi-C data, and annotated using transcriptome data. Overall, the dataset is presented clearly and concisely. The methods are state-of-the-art and follow the usual DToL procedure. The final chromosome-level genome assembly is 444.9 Mb long. With a scaffold N50 of 15.9 Mb and a total number of 36 scaffolds, the assembly is of good quality and provides a valuable addition to the growing list of lepidopteran genomes.

Yet, I have several points that should be addressed.

TITLE:

- It is not clear from the title that this is the genome of a moth or even insect. Some kind of information would be helpful.

BACKGROUND (Paragraph 4):

- 'annual variation in climate change' makes no sense. Please rephrase.
- Should be 'as demonstrated'
- 'population decrease in population [..]' 1x population is redundant
- Please explain how the ash dieback and climate change are connected, as they are mentioned in the same context.
- 'full genome' can only be 'genome', full is redundant.
- 'these vulnerable species' should be 'this vulnerable species'.
- 'insights into relationships between Ennominae moths'.

TABLE 1:

- Isolate information: ilEnnFusc2: male: head and thorax (should probably be 'DNA sequencing' instead of 'RNA sequencing')

METHODS:

- Sequencing section: Illumina HiSeq 4000 twice (redundant or mistake?)
- Please provide more details on the transcriptome data used for annotation and the protein set from UniProt. The sex of the individual used for RNA sequencing would be helpful.

DATA AVAILABILITY:

- Details for accessing the genome annotation results/predicted protein set would be helpful.

Is the rationale for creating the dataset(s) clearly described?

Yes

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

Partly

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Evolutionary genomics, phylogenomics, hymenoptera

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 03 Feb 2026

Tree of Life Team Sanger

Thank you for reviewing this data note.

We have made the corrections you requested to the text of the Background and to Table 1, and thank you for pointing out these errors. The word "moth" has now been added to the title. We have updated the link to the third-party genome annotation provider: <https://beta.ensembl.org/species/aa89cd65-c4eb-4931-aa7d-039604cb00cd>

Tree of Life team

Competing Interests: No competing interests were disclosed.

Reviewer Report 25 January 2024

<https://doi.org/10.21956/wellcomeopenres.22337.r71950>

© 2024 Hof A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Arjen Van 't Hof 

Biology Centre of the Czech Academy of Sciences, Budějovice, Czech Republic

The genome assembly of the Dusky Thorn is a valuable contribution to the scientific community, especially since hardly any sequence resources existed for this species. The assembly meets the same high standards as we are used to from the Darwin Tree of Life Consortium with full-length scaffolds for all 31 chromosomes plus a separate assembly for the alternate haplotype. The chromosome number has not been established by karyotyping before to my knowledge. It is consistent with the ancestral Lepidoptera composition of 30 autosomes and one Z chromosome. A male was used for the assembly, which means that the W chromosome is not included. The chromosomes are numbered based on size instead of gene content. This is somewhat surprising because numbering rules which take shared synteny with ancestral chromosomes into account have been proposed.

The number of identified protein coding genes is lower than reported in other Lepidoptera assemblies, though the BUSCO scores suggest near complete representation.

RNA was extracted from ilEnnFusc1, a different individual than the male used to obtain DNA.

Unfortunately, the sex of ilEnnFusc1 is not specified here or at the sample sheet for SAMEA7520184. The authors mention that the genome assembly can be used to study female 'lightlessness', and therefore, it would have been useful to know the sex of ilEnnFusc1.

The Dusky Thorn genome assembly is a welcome addition to a growing number of high-quality Lepidoptera genome sequences.

Is the rationale for creating the dataset(s) clearly described?

Yes

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Evolutionary Biology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.
