



DATA NOTE

# The genome sequence of the Common Flat-body moth, *Agonopterix heracliana* Linnaeus, 1758 [version 1; peer review: 2 approved]

Douglas Boyes<sup>1+</sup>, Finley Hutchinson<sup>2</sup>, Liam M. Crowley<sup>id</sup><sup>3</sup>, Peter W.H. Holland<sup>id</sup><sup>3</sup>,  
University of Oxford and Wytham Woods Genome Acquisition Lab,  
Darwin Tree of Life Barcoding collective,  
Wellcome Sanger Institute Tree of Life Management, Samples and Laboratory  
team,  
Wellcome Sanger Institute Scientific Operations: Sequencing Operations,  
Wellcome Sanger Institute Tree of Life Core Informatics team,  
Tree of Life Core Informatics collective, Darwin Tree of Life Consortium

<sup>1</sup>UK Centre for Ecology & Hydrology, Wallingford, England, UK

<sup>2</sup>University of Exeter, Penryn, England, UK

<sup>3</sup>University of Oxford, Oxford, England, UK

+ Deceased author

**V1** First published: 21 Jun 2024, 9:329  
<https://doi.org/10.12688/wellcomeopenres.22461.1>  
Latest published: 21 Jun 2024, 9:329  
<https://doi.org/10.12688/wellcomeopenres.22461.1>

## Abstract

We present a genome assembly from an individual male *Agonopterix heracliana* (the Common Flat-body; Arthropoda; Insecta; Lepidoptera; Depressariidae). The genome sequence is 539.1 megabases in span. Most of the assembly is scaffolded into 30 chromosomal pseudomolecules, including the Z sex chromosome. The mitochondrial genome has also been assembled and is 15.36 kilobases in length.

## Keywords

*Agonopterix heracliana*, Common Flat-body moth, genome sequence, chromosomal, Lepidoptera



This article is included in the [Tree of Life](#) gateway.

## Open Peer Review

Approval Status

	1	2
<b>version 1</b>		
21 Jun 2024	<a href="#">view</a>	<a href="#">view</a>

1. **Daniel Doucet**, Natural Resources Canada, Ontario, Canada
2. **Marta Vila** , Universidade da Coruña, A Coruña, Spain

Any reports and responses or comments on the article can be found at the end of the article.

**Corresponding author:** Darwin Tree of Life Consortium ([mark.blaxter@sanger.ac.uk](mailto:mark.blaxter@sanger.ac.uk))

**Author roles:** **Boyes D:** Investigation, Resources; **Hutchinson F:** Investigation, Resources; **Crowley LM:** Investigation, Resources; **Holland PWH:** Writing – Original Draft Preparation, Writing – Review & Editing;

**Competing interests:** No competing interests were disclosed.

**Grant information:** This work was supported by Wellcome through core funding to the Wellcome Sanger Institute [206194, <https://doi.org/10.35802/206194>] and the Darwin Tree of Life Discretionary Award [218328, <https://doi.org/10.35802/218328>]. *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

**Copyright:** © 2024 Boyes D *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**How to cite this article:** Boyes D, Hutchinson F, Crowley LM *et al.* **The genome sequence of the Common Flat-body moth, *Agonopterix heracliana* Linnaeus, 1758 [version 1; peer review: 2 approved]** Wellcome Open Research 2024, 9:329 <https://doi.org/10.12688/wellcomeopenres.22461.1>

**First published:** 21 Jun 2024, 9:329 <https://doi.org/10.12688/wellcomeopenres.22461.1>

## Species taxonomy

Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Protostomia; Ecdysozoa; Panarthropoda; Arthropoda; Mandibulata; Pancrustacea; Hexapoda; Insecta; Dicondylia; Pterygota; Neoptera; Endopterygota; Amphiesmenoptera; Lepidoptera; Glossata; Neolepidoptera; Heteroneura; Ditrysia; Gelechioidea; Depressariidae; Depressariinae; *Agonopterix*; *Agonopterix heracliana* Linnaeus, 1758 (NCBI:txid1073575).

## Background

Moths in the genus *Agonopterix*, family Depressariidae, have a characteristic resting shape with overlapping rounded wings giving an oval outline. Many species in the genus have distinctive wing markings although *Agonopterix heracliana* can be difficult to distinguish from *A. ciliella*, with both species having a row of three pale marks on the speckled grey forewings. Fine markings on the hindwings and genitalia differences can be used to distinguish the two species. There is a complex history concerning the naming of *A. heracliana*, including changes of genus, several misidentifications and a mix-up of historic specimens in Linnaeus' collection; this taxonomic history is described by [Karsholt \*et al.\* \(2006\)](#).

*A. heracliana* has been recorded across much of Europe, with a high concentration of records from the Netherlands, United Kingdom, Denmark and southern regions of Norway, Sweden and Finland ([GBIF Secretariat, 2024](#)). In Britain, the species is commonest in East Anglia, the Thames valley, south Wales and the Wales/England border ([NBN Atlas Partnership, 2024](#)). The adult moth is active in the colder months, from autumn to early spring, with the larvae developing during early summer. Scattered records from around Europe suggest that adults take measures to avoid extremes of heat or cold, for example by sheltering in caves or military bunkers ([Moog \*et al.\*, 2021](#)). The preferred larval foodplants are variety of umbellifers (family Apiaceae), with larvae using silk to spin a tube or fold in a leaf.

A genome sequence of *Agonopterix heracliana* was determined as part of the Darwin Tree of Life project. The genome sequence will facilitate research into larval food plant adaptations and will contribute to the growing set of resources for studying evolution in the Lepidoptera.

## Genome sequence report

The genome was sequenced from an adult *Agonopterix heracliana* ([Figure 1](#)) collected from Wytham Woods, Oxfordshire, UK (51.77, -1.34). A total of 40-fold coverage in Pacific Biosciences single-molecule HiFi long reads was generated. Primary assembly contigs were scaffolded with chromosome conformation Hi-C data. Manual assembly curation corrected 21 missing joins or mis-joins and removed 12 haplotypic duplications, reducing the assembly length by 2.70%, and decreasing the scaffold N50 by 2.83%.

The final assembly has a total length of 539.1 Mb in 47 sequence scaffolds with a scaffold N50 of 19.6 Mb ([Table 1](#)). The snail plot in [Figure 2](#) provides a summary of the assembly statistics, while the distribution of assembly scaffolds



**Figure 1.** Photograph of the *Agonopterix heracliana* (ilAgoHera1) specimen used for genome sequencing.

on GC proportion and coverage is shown in [Figure 3](#). The cumulative assembly plot in [Figure 4](#) shows curves for subsets of scaffolds assigned to different phyla. Most (99.88%) of the assembly sequence was assigned to 30 chromosomal-level scaffolds, representing 30 autosomes and the Z sex chromosome. Chromosome-scale scaffolds confirmed by the Hi-C data are named in order of size ([Figure 5](#); [Table 2](#)). Chromosome Z was assigned by synteny to *Agriphila straminella* (GCA\_950108535.1). While not fully phased, the assembly deposited is of one haplotype. Contigs corresponding to the second haplotype have also been deposited. The mitochondrial genome was also assembled and can be found as a contig within the multifasta file of the genome submission.

The estimated Quality Value (QV) of the final assembly is 68.8 with *k*-mer completeness of 100.0%, and the assembly has a BUSCO v5.4.3 completeness of 97.9% (single = 97.4%, duplicated = 0.5%), using the lepidoptera\_odb10 reference set (*n* = 5,286).

Metadata for specimens, BOLD barcode results, spectra estimates, sequencing runs, contaminants and pre-curation assembly statistics are given at <https://links.tol.sanger.ac.uk/species/1073575>.

## Methods

### Sample acquisition and nucleic acid extraction

An adult *Agonopterix heracliana* (specimen ID Ox000652, ToLID ilAgoHera1) was collected from Wytham Woods, Oxfordshire (biological vice-county Berkshire), UK (latitude 51.77, longitude -1.34) on 2020-07-20 using a light trap. The specimen was collected and identified by Douglas Boyes (University of Oxford) and preserved on dry ice.

The specimen used for Hi-C sequencing (specimen ID Ox003081, ToLID ilAgoHera2) was collected in a light trap

**Table 1. Genome data for *Agonopterix heracliana*, ilAgoHera1.1.**


Project accession data		
Assembly identifier	ilAgoHera1.1	
Species	<i>Agonopterix heracliana</i>	
Specimen	ilAgoHera1	
NCBI taxonomy ID	1073575	
BioProject	PRJEB65231	
BioSample ID	SAMEA7701514	
Isolate information	ilAgoHera1: whole organism (genome sequence) ilAgoHera2: whole organism (Hi-C sequencing)	
Assembly metrics*		Benchmark
Consensus quality (QV)	68.8	≥ 50
k-mer completeness	100.0%	≥ 95%
BUSCO**	C:97.9%[S:97.4%,D:0.5%], F:0.4%,M:1.7%,n:5,286	C ≥ 95%
Percentage of assembly mapped to chromosomes	99.88%	≥ 95%
Sex chromosomes	Z	<i>localised homologous pairs</i>
Organelles	Mitochondrial genome: 15.36 kb	<i>complete single alleles</i>
Raw data accessions		
PacificBiosciences Sequel Iie	ERR11867228	
Hi-C Illumina	ERR11872599	
Genome assembly		
Assembly accession	GCA_963693445.1	
Accession of alternate haplotype	GCA_963693455.1	
Span (Mb)	539.1	
Number of contigs	61	
Contig N50 length (Mb)	17.9	
Number of scaffolds	47	
Scaffold N50 length (Mb)	19.6	
Longest scaffold (Mb)	29.76	

\* Assembly metric benchmarks are adapted from column VGP-2020 of “Table 1: Proposed standards and metrics for defining genome assembly quality” from [Rhie et al. \(2021\)](#).

\*\* BUSCO scores based on the lepidoptera\_odb10 BUSCO set using version v5.4.3. C = complete [S = single copy, D = duplicated], F = fragmented, M = missing, n = number of orthologues in comparison. A full set of BUSCO scores is available at [https://blobtoolkit.genomehubs.org/view/Agonopterix\\_heracliana/dataset/GCA\\_963693445.1/busco](https://blobtoolkit.genomehubs.org/view/Agonopterix_heracliana/dataset/GCA_963693445.1/busco).

at the same location on 2022-07-22. The specimen was collected by Liam Crowley (University of Oxford) and Finley Hutchinson (University of Essex) and identified by Finley Hutchinson and preserved on dry ice.

The workflow for high molecular weight (HMW) DNA extraction at the Wellcome Sanger Institute (WSI) Tree of Life Core Laboratory includes a sequence of core procedures: sample preparation; sample homogenisation, DNA extraction,



**Figure 2. Genome assembly of *Agonopterix heracliana*, ilAgoHera1.1: metrics.** The BlobToolKit snail plot shows N50 metrics and BUSCO gene completeness. The main plot is divided into 1,000 size-ordered bins around the circumference with each bin representing 0.1% of the 539,129,159 bp assembly. The distribution of scaffold lengths is shown in dark grey with the plot radius scaled to the longest scaffold present in the assembly (29,756,636 bp, shown in red). Orange and pale-orange arcs show the N50 and N90 scaffold lengths (19,568,409 and 12,633,633 bp), respectively. The pale grey spiral shows the cumulative scaffold count on a log scale with white scale lines showing successive orders of magnitude. The blue and pale-blue area around the outside of the plot shows the distribution of GC, AT and N percentages in the same bins as the inner plot. A summary of complete, fragmented, duplicated and missing BUSCO genes in the lepidoptera\_odb10 set is shown in the top right. An interactive version of this figure is available at [https://blobtoolkit.genomehubs.org/view/Agonopterix\\_heracliana/dataset/GCA\\_963693445.1/snail](https://blobtoolkit.genomehubs.org/view/Agonopterix_heracliana/dataset/GCA_963693445.1/snail).

fragmentation, and clean-up. In sample preparation, the ilAgoHera1 sample was weighed and dissected on dry ice (Jay *et al.*, 2023). Tissue from the whole organism was homogenised using a PowerMasher II tissue disruptor (Denton *et al.*, 2023a).


HMW DNA was extracted using the Automated MagAttract v1 protocol (Sheerin *et al.*, 2023). DNA was sheared into an average fragment size of 12–20 kb in a Megaruptor 3 system with speed setting 30 (Todorovic *et al.*, 2023). Sheared DNA was purified by solid-phase reversible immobilisation (Strickland *et al.*, 2023): in brief, the method employs a 1.8X ratio of AMPure PB beads to sample to eliminate shorter fragments and concentrate the DNA. The concentration of the sheared and purified DNA was assessed using a Nanodrop spectrophotometer and Qubit Fluorometer and Qubit dsDNA

High Sensitivity Assay kit. Fragment size distribution was evaluated by running the sample on the FemtoPulse system.

Protocols developed by the WSI Tree of Life laboratory are publicly available on protocols.io (Denton *et al.*, 2023b).

### Sequencing

Pacific Biosciences HiFi circular consensus DNA sequencing libraries were constructed according to the manufacturers' instructions. DNA sequencing was performed by the Scientific Operations core at the WSI on a Pacific Biosciences Sequel IIe instrument. Hi-C data were also generated from ilAgoHera2 using the Arima v2 kit. The Hi-C sequencing was performed using paired-end sequencing with a read length of 150 bp on the Illumina NovaSeq 6000 instrument.



**Figure 3. Genome assembly of *Agonopterix heracliana*, ilAgoHera1.1: BlobToolKit GC-coverage plot.** Sequences are coloured by phylum. Circles are sized in proportion to sequence length. Histograms show the distribution of sequence length sum along each axis. An interactive version of this figure is available at [https://blobtoolkit.genomehubs.org/view/Agonopterix\\_heracliana/dataset/GCA\\_963693445.1/blob](https://blobtoolkit.genomehubs.org/view/Agonopterix_heracliana/dataset/GCA_963693445.1/blob).


### Genome assembly and curation

Assembly was carried out with Hifiasm (Cheng *et al.*, 2021) and haplotypic duplication was identified and removed with purge\_dups (Guan *et al.*, 2020). The assembly was then scaffolded with Hi-C data (Rao *et al.*, 2014) using YaHS (Zhou *et al.*, 2023). The assembly was checked for contamination and corrected using the TreeVal pipeline (Pointon *et al.*, 2023). Manual curation was performed using JBrowse2 (Diesh *et al.*, 2023), HiGlass (Kerpedjiev *et al.*, 2018) and PretextView (Harry, 2022). The mitochondrial genome was assembled using MitoHiFi (Uliano-Silva *et al.*, 2023), which runs MitoFinder (Allio *et al.*, 2020) or MITOS

(Bernt *et al.*, 2013) and uses these annotations to select the final mitochondrial contig and to ensure the general quality of the sequence.

### Final assembly evaluation

The final assembly was post-processed and evaluated with the three Nextflow (Di Tommaso *et al.*, 2017) DSL2 pipelines “sanger-tol/readmapping” (Surana *et al.*, 2023a), “sanger-tol/genomenote” (Surana *et al.*, 2023b), and “sanger-tol/blobtoolkit” (Muffato *et al.*, 2024). The pipeline sanger-tol/readmapping aligns the Hi-C reads with bwa-mem2 (Vasimuddin *et al.*, 2019) and combines the alignment files with




**Figure 4. Genome assembly of *Agonopterix heracliana*, ilAgoHera1.1: BlobToolKit cumulative sequence plot.** The grey line shows cumulative length for all sequences. Coloured lines show cumulative lengths of sequences assigned to each phylum using the buscogenes taxruler. An interactive version of this figure is available at [https://blobtoolkit.genomehubs.org/view/Agonopterix\\_heracliana/dataset/GCA\\_963693445.1/cumulative](https://blobtoolkit.genomehubs.org/view/Agonopterix_heracliana/dataset/GCA_963693445.1/cumulative).

SAMtools (Danecek *et al.*, 2021). The sanger-tol/genomenote pipeline transforms the Hi-C alignments into a contact map with BEDTools (Quinlan & Hall, 2010) and the Cooler tool suite (Abdennur & Mirny, 2020), which is then visualised with HiGlass (Kerpedjiev *et al.*, 2018). It also provides statistics about the assembly with the NCBI datasets (Sayers *et al.*, 2024) report, computes  $k$ -mer completeness and QV consensus quality values with FastK and MerquryFK, and a completeness assessment with BUSCO (Manni *et al.*, 2021).

The sanger-tol/blobtoolkit pipeline is a Nextflow port of the previous Snakemake Blobtoolkit pipeline (Challis *et al.*, 2020). It aligns the PacBio reads with SAMtools and minimap2 (Li, 2018) and generates coverage tracks for

regions of fixed size. In parallel, it queries the GoAT database (Challis *et al.*, 2023) to identify all matching BUSCO lineages to run BUSCO (Manni *et al.*, 2021). For the three domain-level BUSCO lineage, the pipeline aligns the BUSCO genes to the Uniprot Reference Proteomes database (Bateman *et al.*, 2023) with DIAMOND (Buchfink *et al.*, 2021) blastp. The genome is also split into chunks according to the density of the BUSCO genes from the closest taxonomically lineage, and each chunk is aligned to the Uniprot Reference Proteomes database with DIAMOND blastx. Genome sequences that have no hit are then chunked with seqtk and aligned to the NT database with blastn (Altschul *et al.*, 1990). All those outputs are combined with the blobtools suite into a blobdir for visualisation.





**Figure 5. Genome assembly of *Agonopterix heracliana*, ilAgoHera1.1: Hi-C contact map of the ilAgoHera1.1 assembly, visualised using HiGlass.** Chromosomes are shown in order of size from left to right and top to bottom. An interactive version of this figure may be viewed at [https://genome-note-higlass.tol.sanger.ac.uk/?d=Ry4u\\_F2cR--JG\\_dTYrjKQ](https://genome-note-higlass.tol.sanger.ac.uk/?d=Ry4u_F2cR--JG_dTYrjKQ).

**Table 2. Chromosomal pseudomolecules in the genome assembly of *Agonopterix heracliana*, ilAgoHera1.**

INSDC accession	Chromosome	Length (Mb)	GC%
OY856308.1	1	22.63	38.0
OY856309.1	2	22.55	38.0
OY856310.1	3	22.45	37.5
OY856311.1	4	21.97	37.5
OY856312.1	5	21.8	37.5
OY856313.1	6	21.78	38.0
OY856314.1	7	21.55	37.5
OY856315.1	8	21.33	38.0
OY856316.1	9	21.07	37.5
OY856317.1	10	20.14	37.5
OY856318.1	11	20.02	37.5
OY856319.1	12	19.57	38.0
OY856320.1	13	19.45	38.0
OY856321.1	14	17.91	38.0

INSDC accession	Chromosome	Length (Mb)	GC%
OY856322.1	15	18.98	37.5
OY856323.1	16	18.76	37.5
OY856324.1	17	18.68	37.5
OY856325.1	18	18.3	38.0
OY856326.1	19	17.88	38.5
OY856327.1	20	16.5	38.5
OY856328.1	21	14.62	38.0
OY856329.1	22	13.89	38.5
OY856330.1	23	12.99	38.5
OY856331.1	24	12.63	38.0
OY856332.1	25	11.79	38.0
OY856333.1	26	10.59	39.5
OY856334.1	27	9.71	38.5
OY856335.1	28	9.04	39.0
OY856336.1	29	8.76	39.5
OY856307.1	Z	29.76	37.5
OY856337.1	MT	0.02	20.0



All three pipelines were developed using the nf-core tooling (Ewels *et al.*, 2020), use MultiQC (Ewels *et al.*, 2016), and make extensive use of the Conda package manager, the Bioconda initiative (Grüning *et al.*, 2018), the Biocontainers infrastructure (da Veiga Leprevost *et al.*, 2017), and the Docker (Merkel, 2014) and Singularity (Kurtzer *et al.*, 2017) containerisation solutions.

Table 3 contains a list of relevant software tool versions and sources.

#### Wellcome Sanger Institute – Legal and Governance

The materials that have contributed to this genome note have been supplied by a Darwin Tree of Life Partner. The

**Table 3. Software tools: versions and sources.**

Software tool	Version	Source
BEDTools	2.30.0	<a href="https://github.com/arq5x/bedtools2">https://github.com/arq5x/bedtools2</a>
Blast	2.14.0	<a href="ftp://ftp.ncbi.nlm.nih.gov/blast/executables/blast+/">ftp://ftp.ncbi.nlm.nih.gov/blast/executables/blast+/</a>
BlobToolKit	4.3.7	<a href="https://github.com/blobtoolkit/blobtoolkit">https://github.com/blobtoolkit/blobtoolkit</a>
BUSCO	5.4.3 and 5.5.0	<a href="https://gitlab.com/ezlab/busco">https://gitlab.com/ezlab/busco</a>
bwa-mem2	2.2.1	<a href="https://github.com/bwa-mem2/bwa-mem2">https://github.com/bwa-mem2/bwa-mem2</a>
Cooler	0.8.11	<a href="https://github.com/open2c/cooler">https://github.com/open2c/cooler</a>
DIAMOND	2.1.8	<a href="https://github.com/bbuchfink/diamond">https://github.com/bbuchfink/diamond</a>
fasta_windows	0.2.4	<a href="https://github.com/tolkit/fasta_windows">https://github.com/tolkit/fasta_windows</a>
FastK	427104ea91c78c3b8b8b49f1a7d6bbeaa869ba1c	<a href="https://github.com/thegenemyers/FASTK">https://github.com/thegenemyers/FASTK</a>
GoaT CLI	0.2.5	<a href="https://github.com/genomehubs/goat-cli">https://github.com/genomehubs/goat-cli</a>
Hifiasm	0.19.5-r587	<a href="https://github.com/chhylp123/hifiasm">https://github.com/chhylp123/hifiasm</a>
HiGlass	44086069ee7d4d3f6f3f0012569789ec138f42b84aa44357826c0b6753eb28de	<a href="https://github.com/higlass/higlass">https://github.com/higlass/higlass</a>
MercuryFK	d00d98157618f4e8d1a9190026b19b471055b22e	<a href="https://github.com/thegenemyers/MERQURY.FK">https://github.com/thegenemyers/MERQURY.FK</a>
MitoHiFi	3	<a href="https://github.com/marcelauliano/MitoHiFi">https://github.com/marcelauliano/MitoHiFi</a>
MultiQC	1.14, 1.17, and 1.18	<a href="https://github.com/MultiQC/MultiQC">https://github.com/MultiQC/MultiQC</a>
NCBI Datasets	15.12.0	<a href="https://github.com/ncbi/datasets">https://github.com/ncbi/datasets</a>
Nextflow	23.04.0-5857	<a href="https://github.com/nextflow-io/nextflow">https://github.com/nextflow-io/nextflow</a>
PretextView	0.2	<a href="https://github.com/wtsi-hpag/PretextView">https://github.com/wtsi-hpag/PretextView</a>
purge_dups	1.2.5	<a href="https://github.com/dfguan/purge_dups">https://github.com/dfguan/purge_dups</a>
samtools	1.16.1, 1.17, and 1.18	<a href="https://github.com/samtools/samtools">https://github.com/samtools/samtools</a>
sanger-tol/genomenote	1.1.1	<a href="https://github.com/sanger-tol/genomenote">https://github.com/sanger-tol/genomenote</a>
sanger-tol/readmapping	1.2.1	<a href="https://github.com/sanger-tol/readmapping">https://github.com/sanger-tol/readmapping</a>
Seqtk	1.3	<a href="https://github.com/lh3/seqtk">https://github.com/lh3/seqtk</a>
Singularity	3.9.0	<a href="https://github.com/sylabs/singularity">https://github.com/sylabs/singularity</a>
TreeVal	1.0.0	<a href="https://github.com/sanger-tol/treeval">https://github.com/sanger-tol/treeval</a>
YaHS	1.2a.2	<a href="https://github.com/c-zhou/yahs">https://github.com/c-zhou/yahs</a>

submission of materials by a Darwin Tree of Life Partner is subject to the ‘**Darwin Tree of Life Project Sampling Code of Practice**’, which can be found in full on the Darwin Tree of Life website [here](#). By agreeing with and signing up to the Sampling Code of Practice, the Darwin Tree of Life Partner agrees they will meet the legal and ethical requirements and standards set out within this document in respect of all samples acquired for, and supplied to, the Darwin Tree of Life Project.

Further, the Wellcome Sanger Institute employs a process whereby due diligence is carried out proportionate to the nature of the materials themselves, and the circumstances under which they have been/are to be collected and provided for use. The purpose of this is to address and mitigate any potential legal and/or ethical implications of receipt and use of the materials as part of the research project, and to ensure that in doing so we align with best practice wherever possible. The overarching areas of consideration are:

- Ethical review of provenance and sourcing of the material
- Legality of collection, transfer and use (national and international)

Each transfer of samples is further undertaken according to a Research Collaboration Agreement or Material Transfer Agreement entered into by the Darwin Tree of Life Partner, Genome Research Limited (operating as the Wellcome Sanger Institute), and in some circumstances other Darwin Tree of Life collaborators.

## Data availability

European Nucleotide Archive: *Agonopterix heracliana* (common flat-body). Accession number PRJEB65231;

<https://identifiers.org/ena.embl/PRJEB65231> (Wellcome Sanger Institute, 2023). The genome sequence is released openly for reuse. The *Agonopterix heracliana* genome sequencing initiative is part of the Darwin Tree of Life (DToL) project. All raw sequence data and the assembly have been deposited in INSDC databases. The genome will be annotated using available RNA-Seq data and presented through the **Ensembl** pipeline at the European Bioinformatics Institute. Raw data and assembly accession identifiers are reported in [Table 1](#).

## Author information

Members of the University of Oxford and Wytham Woods Genome Acquisition Lab are listed here: <https://doi.org/10.5281/zenodo.7125292>.

Members of the Darwin Tree of Life Barcoding collective are listed here: <https://doi.org/10.5281/zenodo.4893703>.

Members of the Wellcome Sanger Institute Tree of Life Management, Samples and Laboratory team are listed here: <https://doi.org/10.5281/zenodo.10066175>.

Members of Wellcome Sanger Institute Scientific Operations: Sequencing Operations are listed here: <https://doi.org/10.5281/zenodo.10043364>.

Members of the Wellcome Sanger Institute Tree of Life Core Informatics team are listed here: <https://doi.org/10.5281/zenodo.10066637>.

Members of the Tree of Life Core Informatics collective are listed here: <https://doi.org/10.5281/zenodo.5013541>.

Members of the Darwin Tree of Life Consortium are listed here: <https://doi.org/10.5281/zenodo.4783558>.

## References

- Abdennur N, Mirny LA: **Cooler: scalable storage for Hi-C data and other genomically labeled arrays**. *Bioinformatics*. 2020; **36**(1): 311–316. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Allio R, Schomaker-Bastos A, Romiguié J, et al.: **MitoFinder: efficient automated large-scale extraction of mitogenomic data in target enrichment phylogenomics**. *Mol Ecol Resour*. 2020; **20**(4): 892–905. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Altschul SF, Gish W, Miller W, et al.: **Basic Local Alignment Search Tool**. *J Mol Biol*. 1990; **215**(3): 403–410. [PubMed Abstract](#) | [Publisher Full Text](#)
- Bateman A, Martin MJ, Orchard S, et al.: **UniProt: the Universal Protein Knowledgebase in 2023**. *Nucleic Acids Res*. 2023; **51**(D1): D523–D531. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bernt M, Donath A, Jühling F, et al.: **MITOS: improved *de novo* metazoan mitochondrial genome annotation**. *Mol Phylogenet Evol*. 2013; **69**(2): 313–319. [PubMed Abstract](#) | [Publisher Full Text](#)
- Buchfink B, Reuter K, Drost HG: **Sensitive protein alignments at Tree-of-Life scale using DIAMOND**. *Nat Methods*. 2021; **18**(4): 366–368. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Challis R, Kumar S, Sotero-Caio C, et al.: **Genomes on a Tree (GoaT): a versatile, scalable search engine for genomic and sequencing project metadata across the eukaryotic tree of life [version 1; peer review: 2 approved]**. *Wellcome Open Res*. 2023; **8**: 24. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Challis R, Richards E, Rajan J, et al.: **BlobToolKit – interactive quality assessment of genome assemblies**. *G3 (Bethesda)*. 2020; **10**(4): 1361–1374. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cheng H, Concepcion GT, Feng X, et al.: **Haplotype-resolved *de novo* assembly using phased assembly graphs with hifiasm**. *Nat Methods*. 2021; **18**(2): 170–175. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- da Veiga Leprevost F, Grüning BA, Alves Aflitos S, et al.: **BioContainers: an open-source and community-driven framework for software standardization**. *Bioinformatics*. 2017; **33**(16): 2580–2582. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Danecek P, Bonfield JK, Liddle J, et al.: **Twelve years of SAMtools and BCFtools**. *GigaScience*. 2021; **10**(2): giab008. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Denton A, Oatley G, Cornwell C, et al.: **Sanger Tree of Life sample homogenisation: PowerMash**. *protocols.io*. 2023a. [Publisher Full Text](#)
- Denton A, Yatsenko H, Jay J, et al.: **Sanger Tree of Life wet laboratory protocol collection**. *protocols.io*. 2023b. [Publisher Full Text](#)

- Di Tommaso P, Chatzou M, Floden EW, *et al.*: **Nextflow enables reproducible computational workflows.** *Nat Biotechnol.* 2017; **35**(4): 316–319.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Diesh C, Stevens GJ, Xie P, *et al.*: **JBrowse 2: a modular genome browser with views of synteny and structural variation.** *Genome Biol.* 2023; **24**(1): 74.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ewels P, Magnusson M, Lundin S, *et al.*: **MultiQC: summarize analysis results for multiple tools and samples in a single report.** *Bioinformatics.* 2016; **32**(19): 3047–3048.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ewels PA, Peltzer A, Fillinger S, *et al.*: **The nf-core framework for community-curated bioinformatics pipelines.** *Nat Biotechnol.* 2020; **38**(3): 276–278.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- GBIF Secretariat: **Agonopterix heracliana Linnaeus, 1758.** *GBIF Backbone Taxonomy.* 2024; [Accessed 29 April 2024].  
[Reference Source](#)
- Grünig B, Dale R, Sjödin A, *et al.*: **Bioconda: sustainable and comprehensive software distribution for the life sciences.** *Nat Methods.* 2018; **15**(7): 475–476.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Guan D, McCarthy SA, Wood J, *et al.*: **Identifying and removing haplotypic duplication in primary genome assemblies.** *Bioinformatics.* 2020; **36**(9): 2896–2898.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Harry E: **PretextView (Paired Read Texture Viewer): a desktop application for viewing pretext contact maps.** 2022; [Accessed 19 October 2022].  
[Reference Source](#)
- Jay J, Yatsenko H, Narváez-Gómez JP, *et al.*: **Sanger Tree of Life sample preparation: triage and dissection.** *protocols.io.* 2023.  
[Publisher Full Text](#)
- Karsholt O, Lvovsky AL, Nielsen C: **A new species of Agonopterix feeding on giant hogweed (*Heracleum mantegazzianum*) in the Caucasus, with a discussion of the nomenclature of *A. heracliana* (Linnaeus) (Depressariidae). *Nota Lepidoptera.* 2006; **28**: 177–192.  
[Reference Source](#)**
- Kerpedjiev P, Abdennur N, Lekschas F, *et al.*: **HiGlass: web-based visual exploration and analysis of genome interaction maps.** *Genome Biol.* 2018; **19**(1): 125.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kurtzer GM, Sochat V, Bauer MW: **Singularity: scientific containers for mobility of compute.** *PLoS One.* 2017; **12**(5): e0177459.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Li H: **Minimap2: pairwise alignment for nucleotide sequences.** *Bioinformatics.* 2018; **34**(18): 3094–3100.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Manni M, Berkeley MR, Seppely M, *et al.*: **BUSCO update: novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes.** *Mol Biol Evol.* 2021; **38**(10): 4647–4654.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Merkel D: **Docker: lightweight Linux containers for consistent development and deployment.** *Linux Journal.* 2014; **2014**(239): 2.  
[Publisher Full Text](#)
- Moog O, Christian E, Eis R: **Increased cave use by butterflies and moths: a response to climate warming?** *Int J Speleol.* 2021; **50**(1): 15–24.  
[Publisher Full Text](#)
- Muffato M, Butt Z, Challis R, *et al.*: **sanger-tol/blobtoolkit: v0.3.0 - Poliwig.** 2024.  
[Publisher Full Text](#)
- NBN Atlas Partnership: **Agonopterix heracliana (Linnaeus, 1758) Common Flat-body.** *NBN Atlas.* 2024; [Accessed 29 April 2024].  
[Reference Source](#)
- Pointon DL, Eagles W, Sims Y, *et al.*: **sanger-tol/treeval v1.0.0 - Ancient Atlantis.** 2023.  
[Publisher Full Text](#)
- Quinlan AR, Hall IM: **BEDTools: a flexible suite of utilities for comparing genomic features.** *Bioinformatics.* 2010; **26**(6): 841–842.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rao SSP, Huntley MH, Durand NC, *et al.*: **A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping.** *Cell.* 2014; **159**(7): 1665–1680.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rhie A, McCarthy SA, Fedrigo O, *et al.*: **Towards complete and error-free genome assemblies of all vertebrate species.** *Nature.* 2021; **592**(7856): 737–746.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sayers EW, Cavanaugh M, Clark K, *et al.*: **GenBank 2024 Update.** *Nucleic Acids Res.* 2024; **52**(D1): D134–D137.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sheerin E, Sampaio F, Oatley G, *et al.*: **Sanger Tree of Life HMW DNA extraction: automated MagAttract v.1.** *protocols.io.* 2023.  
[Publisher Full Text](#)
- Strickland M, Cornwell C, Howard C: **Sanger Tree of Life fragmented DNA clean up: manual SPRI.** *protocols.io.* 2023.  
[Publisher Full Text](#)
- Surana P, Muffato M, Qi G: **Sanger-tol/readmapping: sanger-tol/readmapping v1.1.0 - hebridean black (1.1.0).** *Zenodo.* 2023a.  
[Publisher Full Text](#)
- Surana P, Muffato M, Sadasivan Baby C: **sanger-tol/genomenote (v1.0.dev).** *Zenodo.* 2023b.  
[Publisher Full Text](#)
- Todorovic M, Sampaio F, Howard C: **Sanger Tree of Life HMW DNA fragmentation: diagenode Megaruptor<sup>3</sup> for PacBio HiFi.** *protocols.io.* 2023.  
[Publisher Full Text](#)
- Uliano-Silva M, Ferreira JGRN, Krashenninnikova K, *et al.*: **MitoHiFi: a python pipeline for mitochondrial genome assembly from PacBio high fidelity reads.** *BMC Bioinformatics.* 2023; **24**(1): 288.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Vasimuddin M, Misra S, Li H, *et al.*: **Efficient architecture-aware acceleration of BWA-MEM for multicore systems.** In: *2019 IEEE International Parallel and Distributed Processing Symposium (IPDPS).* IEEE, 2019; 314–324.  
[Publisher Full Text](#)
- Wellcome Sanger Institute: **The genome sequence of the Common Flat-body moth, Agonopterix heracliana Linnaeus, 1758.** European Nucleotide Archive. [dataset], accession number PRJEB65231, 2023.
- Zhou C, McCarthy SA, Durbin R: **YaHS: yet another Hi-C scaffolding tool.** *Bioinformatics.* 2023; **39**(1): btac808.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

# Open Peer Review

Current Peer Review Status:  

---

## Version 1

Reviewer Report 12 October 2024

<https://doi.org/10.21956/wellcomeopenres.24748.r101601>

© 2024 Vila M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Marta Vila** 

Universidade da Coruna, A Coruña, Galicia, Spain

*Agonopterix heracliana* (Linnaeus, 1758) is a common species of moth in the family Depressariidae, widely distributed across the UK and much of Europe. Commonly referred to as the "Common Flat-body" or "Common Brindled Brown", this species is often associated with umbelliferous plants, such as common hogweed *Heracleum sphondylium* (hence, the "heracliana") and cow parsley (*Anthriscus sylvestris*), where the larvae feed on leaves. Although *A. heracliana* does not pose significant agricultural threats, it plays an important ecological role as a herbivore in various ecosystems. Given its wide distribution and interactions with various plant species, this genome assembly represents a valuable resource for future ecological, evolutionary, and comparative genomic studies.

The sequencing and assembly of the genome of *A. heracliana* provide a foundational resource for investigating a range of biological questions. As one of the first genomic resources for a species within the Depressariidae family, this genome will allow researchers to explore the evolutionary relationships between this species and other Lepidoptera, particularly in understanding divergence patterns within the superfamily Gelechioidea. Furthermore, the availability of this genome will be crucial for examining the molecular basis of ecological adaptations, such as host plant selection and tolerance to environmental stressors.

The manuscript follows the typical structure for a Genome Note, and the procedures are appropriate. My comments are as follows:

- 1. Background Section:** I would suggest a minor change in this sentence "...*heracliana* can be difficult to distinguish **in the UK** from *A. ciliella*". This is because a third, very similar species, *A. caucasiella*, is also present, though it is primarily distributed in the Caucasus region (Karsholt et al. 2006).
- 2. Genome Sequence Report Section:** The authors should acknowledge that sequencing a female specimen is necessary to provide information about the W chromosome. |
- 3. Genome Sequence Report Section:** I would appreciate a brief explanation of how future

research could address the issue of the Z chromosome not being fully phased. The sentence "While not fully phased, the assembly deposited is of one haplotype. Contigs corresponding to the second haplotype have also been deposited" could be expanded to clarify how this challenge might be resolved in the future.

**Is the rationale for creating the dataset(s) clearly described?**

Yes

**Are the protocols appropriate and is the work technically sound?**

Yes

**Are sufficient details of methods and materials provided to allow replication by others?**

Yes

**Are the datasets clearly presented in a useable and accessible format?**

Yes

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** Molecular Ecology, Phylogeography

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

Reviewer Report 09 October 2024

<https://doi.org/10.21956/wellcomeopenres.24748.r94991>

© 2024 Doucet D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Daniel Doucet**

Canadian Forest Service, Natural Resources Canada, Ontario, Canada

The report by Boyes and colleagues describes the complete genome sequencing of the moth *Agonopterix heracliana*. The Depressariidae is a moderately speciose family of moths (approximately 2,300) within the Gelechioidea superfamily. The Gelechioidea are known to display very diverse life history strategies, such as predation or living in aquatic environments. Understanding the adaptive radiation of this group benefits from the provision of complete genomes from as many species as possible, such as *A. heracliana*.

The rationale for creating the dataset for this species is well described. The host plants that this species feeds on (Apiaceae) are known to synthesize a wide array of secondary metabolites, some of which function as defensive compounds. It will be interesting to examine the genome for genes encoding detoxification enzymes that may act against Apiaceae defenses.

The collection, labeling and processing of the insect sample is appropriate and follows the standard of the field in genomics. Although not an expert in bioinformatics, the work follows the standard workflow established by the Tree of Life, with thousands of genomes assembled using the same methods. The methods are described in sufficient details so that the work can be replicated. Likewise the datasets are clearly presented and accessible in a useable format.

**Is the rationale for creating the dataset(s) clearly described?**

Yes

**Are the protocols appropriate and is the work technically sound?**

Yes

**Are sufficient details of methods and materials provided to allow replication by others?**

Yes

**Are the datasets clearly presented in a useable and accessible format?**

Yes

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** Cell biology

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

---