



Review Article

Seasonal dynamics of Arctic soils: Capturing year-round processes in measurements and soil biogeochemical models

Zhou Lyu^a, Pacifica Sommers^b, Steven K. Schmidt^b, Marta Magnani^{c,d}, Mihai Cimpoiasu^e, Oliver Kuras^e, Qianlai Zhuang^f, Youmi Oh^{g,h}, Maria De La Fuenteⁱ, Margaret Cramm^{a,j}, James A. Bradley^{k,a,*}

^a Queen Mary University of London, London, UK

^b University of Colorado, Boulder, CO, USA

^c Institute of Geosciences and Earth Resources, National Research Council, Turin, Italy

^d National Biodiversity Future Center, Palermo 90133, Italy

^e British Geological Survey, Nottingham, United Kingdom

^f Purdue University, West Lafayette, USA

^g National Oceanic and Atmospheric Administration, USA

^h Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, USA

ⁱ Université Libre de Bruxelles, Brussels, Belgium

^j Natural History Museum, London, UK

^k Aix Marseille University, University of Toulon, CNRS, IRD, MIO, Marseille, France



ARTICLE INFO

Keywords:

Arctic soils

Seasonality

Soil microorganisms

Soil organic carbon

Biogeochemical modelling

Permafrost

ABSTRACT

The Arctic is undergoing rapid changes in climate, altering the status and functioning of high-latitude soils and permafrost. The vast majority of studies on Arctic soils and permafrost are conducted during the summer period due to ease of accessibility, sampling, instrument operation, and making measurements, in comparison to during winter and transition seasons. However, there is increasing evidence that microbial activity continues in Arctic soils outside of the summer period. Moreover, it is becoming clear that understanding the seasonal dynamics of Arctic soils is of critical importance, especially considering that the under-studied winter is the period that is most sensitive to climate warming. Soil biogeochemical models have advanced our understanding of the functioning and fate of soils in the Arctic, however it is vital that seasonality in biotic and abiotic processes is accurately captured in these models. Here we synthesize recent investigations and observations of the year-round functioning of Arctic soils, review soil biogeochemical modelling frameworks, and highlight certain processes and behaviors that are shaped by seasonality and thus warrant particular consideration within these models. More attention to seasonal processes will be critical to improving datasets and soil biogeochemical models that can be used to understand the year-round functioning of soils and the fate of the soil carbon reservoir in the Arctic.

1. Introduction

The Arctic has undergone rapid changes in climate during recent decades, including widespread increases in air temperature and overall increases in annual precipitation (as rainfall and snowfall) (Box et al., 2019; Meredith et al., 2019; Kalnay et al., 1996; Rapačić et al., 2015; Rawlins et al., 2010). In particular, the winter period is experiencing the highest rates of human-induced climate warming (Graham et al., 2017; Post et al., 2019): from 1971 to 2017, Arctic air temperatures have

increased on average by 3.1 °C during the winter months (October–May), compared with a 1.8 °C increase during the summer period (June–September) (Box et al., 2019). Concurrently, snow cover and permafrost area are generally in decline across most regions of the Arctic (except for some parts of Siberia which have recorded increased prevalence and persistence of seasonal snow cover) (Bormann et al., 2018; Bulygina et al., 2009; Lemke et al., 2007; Schindler and Donahue, 2006; Tomczyk et al., 2021). Changes in air temperature and precipitation have induced a series of alterations to soil and permafrost, including

* Corresponding author at: Aix Marseille University, University of Toulon, CNRS, IRD, MIO, Marseille, France.

E-mail address: jbradley.earth@gmail.com (J.A. Bradley).

<https://doi.org/10.1016/j.earscirev.2024.104820>

Received 27 October 2023; Received in revised form 16 May 2024; Accepted 17 May 2024

Available online 19 May 2024

0012-8252/© 2024 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

increases in soil temperature, changes to soil moisture regimes, deepening of permafrost active-layer depths, reductions in the spatial extent of permafrost, prolonging of the vegetation growing season, and shortening durations of snow cover (Box et al., 2019; Brown et al., 2018; Lund et al., 2012, 2014; Meredith et al., 2019; Nicolsky et al., 2017; Parmentier et al., 2011; Romanovsky et al., 2017). There is mounting evidence that feedback processes are already affecting Arctic soils, including increased primary production and growing season carbon uptake, as well as elevated soil respiration during the non-growing season, resulting in significant changes to the net carbon fluxes between the land and atmosphere (Chapin III et al., 2005; Lemke et al., 2007; López-Blanco et al., 2017; Lund et al., 2010; Lund, 2018; Röbger et al., 2022).

The Arctic region contains large quantities of soil organic carbon. Hugelius et al. (2014) estimated that ~1300 Pg C is stored in Arctic soils, with 472 ± 27 Pg C in the upper-most 1 m, accounting for close to half of the total organic carbon stored in soils globally (Batjes, 2014; Jobbágy and Jackson, 2000). This enormous carbon pool is largely stored in soils with mean annual temperatures below freezing ($< 0^\circ\text{C}$) (Hugelius et al., 2014). These soils are highly vulnerable to climate warming, which causes thawing of permafrost that accelerates carbon mobilization and decomposition processes, elevating rates of carbon exchange between soil and the atmosphere. Both long-term records and process-based models have confirmed an increase in the net flux of soil carbon into the atmosphere due to warming of the Arctic over the past two decades (Euskirchen et al., 2017; McGuire et al., 2012).

Soils in high latitudes exhibit distinct seasonality in biotic and abiotic characteristics. Summers of perpetual daylight are punctuated by extended periods of 24-h darkness, snow cover and sub-zero temperatures (Fig. 1). Microbial respiration rates during wintertime in Arctic soils have typically been assumed to be low, limited largely by below-freezing temperatures and therefore reduced liquid water availability to soil microorganisms (Mazur, 1980; Arndt et al., 2020; Dunfield et al., 1993). However, recent studies have observed continuous soil respiration fluxes from Arctic soils during winter periods (Arndt et al., 2020; Natali et al., 2019) as well as bursts of carbon emitted from soils

following early spring thawing (Nielsen et al., 2001; Raz-Yaseef et al., 2017; Teepe and Ludwig, 2004), substantially increasing the annual carbon efflux of arctic tundra ecosystems (Fahnestock et al., 1999). In fact, winter-time soil emissions are now known to be a significant regulator of the net ecosystem carbon balance in the Arctic because of the disproportionate rate of warming in the Arctic during winter, as well as the alterations to the timing of the transition periods between summer and winter seasons (i.e. spring thaw and fall freeze-up). Records have also captured considerable carbon dioxide (CO_2) emissions from Arctic soils during the ‘shoulder seasons’ (i.e., during fall freezing and springtime thawing) (Commane et al., 2017; Zona et al., 2016). Similarly, methane (CH_4) fluxes from Arctic soils during wintertime are considered to make up a significant portion (15–50%) of the annual methane fluxes to the atmosphere in the Arctic region (Flessa et al., 2008; Zona et al., 2016; Treat et al., 2018). Given the high global warming potential of methane (28–34times that of CO_2 over a 100 year period (GWP-100)) and the prevalence of methane-producing thermokarsts in the Arctic (Turetsky et al., 2020), understanding the seasonal changes in methane production and consumption is crucial to better establish current and future global carbon budgets. The timing, magnitude and nature by which greenhouse gasses are produced and released from high-latitude soils is also strongly shaped by variation in transport processes, linked to seasonal changes in the extent of freezing, saturation and snow-cover (Chuvilin et al., 2021; Heinze, 2021; Lange et al., 2016).

A wealth of new observations and experiments on shoulder-season and wintertime Arctic soil dynamics provide insights into better understanding and predicting the seasonality of microbial activity and carbon dynamics in Arctic soils. The emergence of year-round data (e.g., Pedron et al., 2022) will undoubtedly enable an improved understanding and prediction of the fate of Arctic soils under future warming scenarios, as well as the development and application of more robust and accurate mechanistically-based soil biogeochemical models. Numerical models are powerful tools enabling future predictions of feedbacks between the changing climate and the terrestrial carbon cycle, which can help to provide valuable information for policymaking. It may be necessary to consider microbial activity explicitly in soil carbon models




Fig. 1. Time-series of soil properties at the Bayelva permafrost monitoring site in Svalbard for the year 2019. The zero-curtain period (Table 1) is marked by ‘ZC’. The data is a continuation of the data record published in Boike et al. (2018).

because microbes drive a large fraction, if not the majority, of soil carbon turnover in the Arctic, and respond dynamically to environmental and climate change. However, there are large discrepancies between the formulations and outputs of different model simulations, as well as between model simulations and observations (Fisher et al., 2014; McGuire et al., 2016; Webb et al., 2016; Meredith et al., 2019). For example, model-intercomparison studies demonstrate that there is high variability in predicted carbon fluxes from Arctic tundra soils, and that the magnitude of discrepancies between models and observations can be as large as the observed fluxes themselves (Fisher et al., 2014; McGuire et al., 2016; Shirley et al., 2022; Ito et al., 2023). The nature, timing, magnitude and causes of observed discrepancies between both the formulations and the outputs of model simulations are numerous and complex. In particular, winter-time carbon fluxes are poorly simulated by many process-based models (McGuire et al., 2016, 2018a), however can be adequately captured and accounted for in empirical models (Natali et al., 2019).

As evidence for Arctic climate change is mounting, the effects of these changes on the carbon cycle can start to be observed (Bruhwiler et al., 2021). Nevertheless, looking forward, there are major discrepancies (i.e. divergent signs and magnitudes of changes) in the modeled predictions of global soil carbon changes by the end of the century, mainly driven by differences in the assumptions regarding soil carbon persistence, especially in the Arctic (Wieder et al., 2019).

To improve our understanding of Arctic soil carbon cycling, and make better predictions of the fate of these systems under future warming scenarios, it is necessary to simulate soil biogeochemical dynamics not just during summer periods, but year-round, and to include the microbial mechanisms that modulate soil carbon responses to climate. Here, we first synthesize new observations, measurements and experiments capturing winter, shoulder-season, or all-year-round Arctic soil dynamics. Second, we review soil microbial and biogeochemical modelling frameworks and highlight important factors to consider when applying models year-round to bridge gaps in resolving seasonal processes and improving the simulation of microbial activity and carbon fluxes in Arctic terrestrial ecosystems. The term “soil” encompasses diverse landforms and ecosystems, spanning mineral and organic soils to wetlands. Whilst wetlands and seasonally inundated areas are noted as critical hotspots for CH₄ production and emission (Wik et al., 2016; Comyn-Platt et al., 2018), we focus this review primarily on Arctic upland, tundra, mineral and organic soils and cryosols.

2. Seasonality of microbial processes and carbon cycling in Arctic soils

2.1. Soil microbial dynamics and the interplay with soil physical processes

2.1.1. Microbial activity

Soil microorganisms play a crucial role in regulating carbon emissions from Arctic soils (Hopple et al., 2020; Natali et al., 2019; Schuur et al., 2015; Pedron et al., 2022), and are highly temperature and moisture sensitive (Brooks et al., 1997; Clein and Schimel, 1995; Elberling and Brandt, 2003). Indeed, microbial respiration in Arctic soil changes dynamically across seasons (Raz-Yaseef et al., 2017), and with climatic variables such as snow cover (Yi et al., 2020). Similarly, the exchange of greenhouse gases between permafrost soils and the atmosphere are further regulated by above-ground vegetation, through both direct uptake of CO₂ by primary production of plants (Wei et al., 2021), and vegetation-driven changes to the hydrological regime (Keuschnig et al., 2022).

It is now well established that liquid water may persist in frozen peatlands where temperature drops well below the freezing point of water (Pavlova, 1970), and that soil microorganisms continue to actively decompose organic matter at relatively high rates at temperatures well below 0 °C, and at measurable rates down to -18 °C (Elberling and Brandt, 2003) – thus potentially producing a considerable

amount of CO₂ during the snow-covered winter periods lasting many months. Indeed, soil microbes likely remain sensitive to changes in soil temperature and moisture regimes at temperatures below 0 °C (Brooks et al., 1997; Elberling and Brandt, 2003; Monson et al., 2006; Schmidt et al., 2009).

Whilst the upper-most layers of soils may be frozen during winter and shoulder-season periods, deeper layers of soils (Fig. 1) (Boike et al., 2018; Keuschnig et al., 2022) and taliks (Parazoo et al., 2018; Farquharson et al., 2022) may remain thawed, and microorganisms in these deep layers continue to respire, producing CO₂ and CH₄. It is estimated that 14 to 80% of the CO₂ that is produced in snow-covered soils remains trapped in the snow and soil column (Elberling and Brandt, 2003). The amount of CO₂ trapping is highly dependent on soil moisture levels and microbial community dynamics (Keuschnig et al., 2022). The trapping of CO₂ in snow-covered soils implies a substantial decoupling in time between the CO₂ being produced in the soil over winter, and CO₂ emissions that can be measured from the snow/soil surface. High rates of CO₂ emissions typically measured during spring may be partially attributed to trapped CO₂ that is released following changes to soil moisture regimes due to thawing events (Goulden et al., 1998; Zimov et al., 1996). However, more recent evidence shows that the majority of CO₂ production under the snow takes place during late winter (Liptzin et al., 2009; Monson et al., 2006) when soil temperatures are near 0 °C and microbial biomass (especially fungi) reach their maximum population sizes (Schmidt et al., 2009) and CO₂ fluxes are more synchronized with production of CO₂. A recent study by Arndt et al. (2020) indicates that bursts of respiration caused by infiltration of the soil by O₂-rich snow melt water are partially responsible for late-winter fluxes of CO₂ (rather than just arising from bursts of stored gases). Importantly, methane produced by methanogens in thawed deeper layers may escape to the atmosphere largely without being oxidized by methanotrophs if near-surface layers containing methanotrophs remain frozen (Zona et al., 2016).

2.1.2. Microbial biomass, community structure, and function

Shifts in bacterial and fungal community composition of Arctic tundra soils are apparent between early and late summer periods (Männistö and Häggblom, 2006; Wallenstein et al., 2007), and across seasons (i.e. summer and winter) (Buckeridge et al., 2013; Lipson and Schmidt, 2004; Pold et al., 2021; Schadt et al., 2003). Recently, Poppeliers et al. (2022) revealed that Arctic soil microbial biomass and community structure undergo seasonal, with the most dynamic period being the transition between winter and spring, although they stress that the fall to winter transition is relatively unstudied. In addition, Bardgett et al. (2005) and Buckeridge et al. (2013) suggested that Arctic soil microbial communities tend to be bacteria-dominated in summer, and fungal-dominated in winter. Accordingly, shifts in fungal-bacteria (F:B) ratio are predicted during summer-to-winter and winter-to-summer transitions – most notably a spike in F:B ratio during the winter and early spring under the snow (Aanderud et al., 2013; Buckeridge and Grogan, 2008; Lipson et al., 2009; Schadt et al., 2003; Waring et al., 2013; Zinger et al., 2009), potentially leading to functional differences and thus alteration to biogeochemical fluxes (e.g. soil respiration) during the winter season (Monson et al., 2006; Schmidt et al., 2009).

Experimental studies have suggested that microbial communities may be able to respond rapidly to seasonal shifts in substrate availability (Schimel and Mikan, 2005; Sturm et al., 2005). This may be linked to the apparent differences in fungal and bacterial biomass, since fungi and bacteria are known to have different enzymatic degradation potentials. For example, cellulose and lignin are the two most abundant input carbon sources in soil, and studies have found that fungi are more capable of degrading lignin than bacteria under the same environmental conditions (Datta et al., 2017). Lipson et al. (2002) showed that cellulose activity and breakdown of phenolic compounds were highest in the winter in tundra soils corresponding to the highest activity and biomass of fungi in the same soils, and several studies (Finestone et al., 2022;

Schmidt et al., 2008; Singh et al., 2012) have shown that many fungal isolates from cold soils have high extracellular cellulase activity. “Snow mold” fungi are very common and active in snow-covered ecosystems throughout the world, including high-Arctic and Antarctic soils (Tojo and Newsham, 2012; Liptzin et al., 2015). It has been postulated that the filamentous growth form and rapid doubling time of these snow molds allow them to access plant litter on the soil surface during the snow-covered season. These traits give snow molds an advantage over bacteria since most bacteria depend on mobility through an aqueous phase to colonize organic matter (Schmidt et al., 2009).

2.1.3. Physiological changes and dormancy

As well as structural and functional changes, physiological changes can be expected among soil microbial communities across different seasons (Table 1). Many microorganisms are known to transition into a reversible state of dormancy when confronted with unfavorable conditions, subsisting in a state of comparatively lower metabolic activity until environmental conditions become more favorable – upon which they can resume a higher level of activity (Lennon and Jones, 2011). The metabolic activities of dormant vegetative microorganisms are largely limited to essential maintenance processes such as biomolecular repair and replacement, and the allocation of energy towards new biomass growth is minimal. Many bacteria and most fungi also produce spores that exhibit virtually no metabolic activity until they are activated in response to environmental stimuli. Many spore producers are opportunistic fungi that persist during stressful periods as spores and then grow rapidly during times when environmental conditions are favorable to them. For example, some rapidly growing psychrophilic snow-mold fungi such as those related to *Mortierella alpina* become active in the fall and winter and grow rapidly as secondary colonists of decaying organic matter in cold ecosystems including Arctic tundra, and produce both asexual spores and thick-walled meiospores (zygospores) that allow them to survive during the summer (Geml et al., 2021; Schmidt et al., 2008, 2009; Thormann et al., 2003). Dormancy therefore serves an ecologically important role throughout the year in polar environments – in particular, by enhancing the persistence of seasonal microbial communities and maintaining ecosystem diversity and functioning until the onset of conditions that are conducive to growth for seasonally active microbial populations.

2.2. Carbon fluxes

Carbon cycling in Arctic soils is a distinctly seasonal process that is strongly shaped by multiple varying and compounding environmental factors. Arctic soils are generally considered as a CO₂ source during the snow-covered winter season, and as a net-neutral to CO₂ sink during the summer due to higher photosynthesis and respiration (Fisher et al., 2014; Oechel et al., 1993; Ravn et al., 2020). For CH₄, Arctic soils are considered to be a net source on an annual basis with net emissions of CH₄ during the summer (Bruhwiler et al., 2021). However, recent studies have highlighted geographical differences in annual carbon budgets (see Lau et al., 2015), as well as across the transitional and winter seasons (Lüers et al., 2014; Waldrop et al., 2021; Zhang et al., 2019).

Alaska contains a range of different permafrost landforms, that are typically characterized by deep layers of carbon-rich soil. In recent years, as the depth of seasonal thawing deepens and the annual duration of the thawed period lengthens, this deep carbon is becoming more accessible to soil microorganisms and is increasingly vulnerable to degradation. Previous data synthesis efforts have reported the Alaskan tundra permafrost region as a carbon source to the atmosphere over the past 40 years (Belshe et al., 2013), while other synthesis and modelling studies have recognized the region as either a carbon sink or carbon neutral (Fisher et al., 2014; McGuire et al., 2012). Uncertainties in Alaskan soil carbon budgets largely arise due to the inherent difficulties in establishing long-term continuous measurements of soil carbon fluxes

Table 1

Characteristics and seasonal variations in physical and biological properties of Arctic tundra and upland soils.

Type of change	Property	Typical seasonal characteristics
Physical	Thermal properties	<i>Summer:</i> Soil temperatures generally above 0 °C throughout the soil active layer (until the upper-most limit of the permafrost layer). <i>Fall:</i> Zero-curtain phenomenon: soil temperature is de-coupled from air temperature and maintained at 0 °C due to latent heat. Upper-most layers freeze (<0 °C) earliest due to sub-zero air temperatures. The freezing front propagates downwards over time. <i>Winter:</i> The entire soil depth profile (i.e. the active layer and the permafrost layer) is predominantly frozen (<0 °C). <i>Spring:</i> Upper-most layers warm earlier than deeper layers. Zero-curtain phenomenon where soil temperature is de-coupled from air temperature and maintained at 0 °C due to latent heat (less pronounced than fall period).
	Water availability	<i>Summer:</i> Water available in liquid form (>0 °C). <i>Fall:</i> Liquid water begins to freeze. <i>Winter:</i> Majority of soil water is in frozen state, yet limited liquid water may persist in thin films, lenses or pockets. <i>Spring:</i> Frozen soil water begins to melt.
	Snow dynamics	<i>Summer:</i> Summers in the Arctic are generally snow-free. <i>Fall:</i> Thin temporary snow cover. <i>Winter:</i> Snowpack accumulates. Periodic melting and rain-on-snow events possible. <i>Spring:</i> Periods of accumulation and periods melting and rain-on-snow events. Snow may persist until early summer.
	Gas exchange	<i>Summer:</i> Active exchange of gases between soil and atmosphere, and within soil pores. <i>Fall:</i> Generally active gas exchange between soil and atmosphere, and within the soil. <i>Winter:</i> Snowpack and frozen soil restricts gas exchange. <i>Spring:</i> Bursts of gas exchange between soil and atmosphere as the snowpack melts and soil thaws.
Biological	Microbial community structure	Changes to microbial community structure (both bacterial and fungal) observed across seasons.
	Microbial community activity	<i>Summer:</i> Highest rates of activity. <i>Fall:</i> Microbial activity rates begin to reduce as microorganisms are limited by temperature and moisture availability. Many non-psychrophilic microbes may become dormant, whilst psychrophilic snow mold fungi may remain active. <i>Winter:</i> Limited data. Microbial activity restricted but may persist in limited capacity and area. Majority of microorganisms may be dormant. <i>Spring:</i> Microbial activity resumes as soil thaws and microbes transition out of dormancy.
	Microbial community function	Seasonal dynamics vary with site factors and do not seem to have a consistent pattern (see Section 3.2.3). Seasonally-driven changes to the soil hydrological regime affect redox state of the soil and likely drives changes in the dominance between aerobic SOC decomposition (producing CO ₂) and anaerobic SOC decomposition (producing CH ₄).

(continued on next page)

Table 1 (continued)

Type of change	Property	Typical seasonal characteristics
	Fungal to bacterial ratio	Higher fungal:bacterial (F:B) ratios during the winter and early spring hypothesized based on available information, potentially leading to shift in dominant functional characteristics and biochemical fluxes.
	Above-ground vegetation	<i>Summer</i> : Fixation of CO ₂ by plants during periods of high primary productivity. Elevated plant respiration. <i>Shoulder seasons and winter</i> : Minimal to no primary productivity, basal plant respiration.
	Greenhouse gas production and exchange	CO ₂ produced in aerobic soil layers and transported upward and downward via diffusion; CH ₄ produced in (often water-saturated) anaerobic soil layers, transported upward and downward via diffusion and advection, and may be oxidized during transport through adjacent layers (particularly if those layers are oxygenated). Influence of above-ground vegetation on CO ₂ fluxes as described above and vegetation-driven changes to soil hydrological regime.

over appropriate timescales and spatial scales. Indeed, carbon flux measurements from eddy covariance towers provide valuable year-round continuous measurements that can shed light on soil carbon budgets on a seasonal basis, but these fluxes might not always be synchronous with soil microbial respiration (especially during periods of snow-cover).

Data from continuously monitored heath and wet sedge tundra (2008–2015) and tussock tundra (2013–2014) in Alaska suggest that these sites were all net carbon sources on an annual basis, with considerable interannual variability. A substantial amount of carbon was released from the soil to the atmosphere during winter periods (Euskirchen et al., 2017), whilst during the growing season (June–August), these sites were a net carbon sink - with gross primary productivity (GPP) offsetting ecosystem respiration. During the observation periods, both the heath tundra and tussock tundra sites exhibited elevated carbon uptake associated with warming summer temperatures, suggesting that the regional growing-season carbon sink could be enhanced under a warming climate. Nevertheless, despite interannual and seasonal variability, carbon that had accumulated during the growing season was subsequently lost during the snow-covered cold season (from September to May) via soil respiration—indicating that soil microbes were still active during winter, and that active-layer carbon remains available for microbial degradation year-round. The observation of continuous microbial respiration in carbon-rich Arctic Alaskan soils over winter is further supported by CARVE aircraft data collected over Alaskan tundra from 2012 to 2014 (Commune et al., 2017), strengthening the case that Alaskan tundra soils were a net source of carbon to the atmosphere during winter. Both studies also highlighted the early winter (October to December) as a period of significant carbon release - emphasizing the importance of the zero-curtain period (the period during which soil active layer temperature stays near the freezing point of water due to latent heat transfer during the freezing or thawing process, despite substantially different temperatures of the adjacent air, Fig. 1) on the annual carbon balance. In both studies, peak daily net carbon emissions were found to occur in the transitional fall period (late September to early November) – coinciding with the period when plant GPP dropped dramatically while soil microorganisms remained active in the non-frozen soil. Both studies also showed that the net carbon balance turned from carbon loss (winter) to carbon accumulation (summer) around June, coinciding with the beginning of the snow-free period and the onset of plant primary productivity. The net carbon accumulation

peaked in July to August, associated with rising GPP from greening vegetation during the growing season.

For Alaskan wetlands, enhanced summer methanogenesis may be counteracted by high summer CH₄ oxidation, especially in younger bogs where available carbon and nitrogen are comparatively rich. Observations showed that young Alaskan lowlands with collapse-scar bogs from permafrost thawing are a carbon sink in summer and a carbon source in winter, while older peatland plateaus act as a carbon source both in winter and summer (Waldrop et al., 2021).

Studies conducted in the permafrost-affected Siberian Arctic also provide valuable insight into the seasonal dynamics of high-latitude soil carbon fluxes. Continuous eddy covariance measurements on polygonal tundra and river terraces at the Samoylov research site in the Lena River delta, northeastern Siberia, indicate that despite a general accumulation of carbon during the period of measurement (2010 to 2017), the growing season carbon sink is largely offset by low but continuous cold-season soil respiration (Holl et al., 2019). Continuous year-round eddy covariance measurements at this site suggest that whilst 39% of the total annual methane released occurs during the cold season, recent increases in summer air temperature correspond to earlier and intensified summer methane emissions (Rößger et al., 2022). Another study has further suggested that the increased cold season respiration in response to warming might, in the future, outpace the increase in plant carbon uptake during the growing season (Runkle et al., 2013), transforming the Siberian tundra into a carbon source.

Similar patterns in the seasonality of soil carbon fluxes have been observed in west Greenland in recent years, where warming has stimulated both growing season carbon uptake and annual soil carbon loss. A five-year eddy covariance measurement (Zhang et al., 2019) showed that west Greenland heath tundra was a carbon sink of $-35 \pm 15 \text{ g C m}^{-2} \text{ yr}^{-1}$, with rapid transitions in the carbon exchange during the beginning and the end of the growing season. The annual budget was strongly affected by the balance between the growing season carbon accumulation encouraged by warming and the carbon that is released from underneath snow-covered but non-frozen soil. In fact, it has been shown that elevated cold-season CO₂ emissions have reduced the regional carbon sink noticeably (Zhang et al., 2019). Net carbon release from west Greenland heath tundra was highest during the transitional spring-to-summer period, around mid to late June, before the rapid increase in plant GPP, though similar in magnitude to the carbon emitted during the transition from summer to fall. At this site, plant GPP continued to increase during the summer period and the net carbon sink peaked in late August before switching back to a carbon source as temperature and photosynthetically-active radiation decreased. During the five-year measurement period, warmer years (2015 and 2016) were associated with significant increases in both growing season GPP and non-growing season carbon emissions, and the total overall net carbon accumulation was lowest. Considering areas with moist tundra, ice-free west and northeast Greenland tundra were also observed to be a methane sink, consuming 1.4 to 18.3 times more CH₄ than the amount being emitted to the atmosphere from wetlands (D'Imperio et al., 2017; Juncher Jørgensen et al., 2015). These findings further underscore the importance of understanding how different components of the diverse Arctic permafrost ecosystems (upland and lowland tundra, river terraces, heath, wetlands, etc.) respond to the changing climate and together shape the regional seasonality and carbon balance.

Long-term and year-round carbon flux measurements are scarce for far north archipelagos such as Svalbard. In contrast to many other high-latitude permafrost regions, the Svalbard archipelago is semi-desert and the soil is not as carbon rich as many other high-latitude environments such as in North America and Siberia (Sipes et al., 2022; Schuur et al., 2015; Nakatsubo et al., 2005). Soil carbon cycling is mostly driven by local processes that result in highly heterogeneous soil ecosystems (Sipes et al., 2022). Nitrogen, labile carbon, and soil moisture limit microbial activity and plant productivity, especially in the forefield of retreating glaciers (Yoshitake et al., 2007; Nakatsubo et al., 2005; Bradley et al.,

2014), but potential for phosphorus limitation has not been adequately tested. During summer, tundra ecosystems across the Svalbard archipelago are usually observed to be weak carbon sinks (Magnani et al., 2022; Pirk et al., 2017). Soil microbial activity is thought to persist during winter periods as continuous carbon emissions have been captured for at least some part of the cold season (Lüers et al., 2014). A single year eddy covariance observation conducted at a permafrost site in north-west Spitsbergen found that the annual carbon budget was close to neutral from spring 2008 to spring 2009 (Lüers et al., 2014). Low-level but sustained winter carbon emissions were observed until late January, and net carbon fluxes varied positively and negatively between February and June. Carbon taken up during the growing season (June–September) by plants was almost entirely offset by respiration during fall and winter. Similar patterns were observed from January 2015 to January 2016 at the same site, despite the annual carbon balance strongly depending on CO₂ flux peaks of debated origin (Jentsch et al., 2021). During the transition seasons the net carbon balance is generally dominated by emissions, but fall season soil respiration can be higher (Lüers et al., 2014) or lower (Cannone et al., 2016, 2019) during spring, depending on the site and year of measurement. This variability can be explained by the combined effect of active layer thaw depth and surface temperature that limited the emissions during the shoulder seasons (Cannone et al., 2019). Also, seasonal variations of the emissions potentially mirror changes in the structure of soil bacterial communities, as shown at an active-layer site (Schostag et al., 2015). On top of seasonal changes, the size and composition of the microbial community in Svalbard tundra soils has been observed to change after a four-year warming treatment during which 1 °C soil warming was induced, resulting in a 44% increase in summertime emissions (Newsham et al., 2022). Hence, microbial shifts induced by climate change effects may change the balance of natural carbon cycles, which further support the need for in-depth modelling of the complex microbial dynamics.

3. Soil biogeochemical modelling

In this section, we review current process-based soil biogeochemical model formulations and provide recommendations for future developments, specifically considering the need to capture seasonal processes in Arctic soil biogeochemical models. We focus on conventional process-based modelling approaches (reviewed by Chandel et al., 2023) – i.e. those that usually distinguish various components of the system (microbial biomass, organic carbon, etc.) in multiple distinct ‘pools’, linked by a system of coupled ordinary differential equations and parameter values that describe biological and other processes leading to transformations of and transfers between these pools. These equations are then numerically solved over a defined time-period to provide a time-series of model outputs including the size of various reservoirs/pools and the rates/magnitudes of fluxes between them. Alternative modelling approaches such as statistically/empirically based modelling, structural equation modelling, generalized linear mixed models, and individual-based modelling are not considered in this review (Schnecker et al., 2014; Bradley et al., 2016a; Hellweger and Buccì, 2009).

3.1. Soil organic carbon and biomass: Pools, transformations and fluxes

3.1.1. Soil Carbon Pools

In order to simulate the transformation of carbon and other elements from one form to another, it is necessary to distinguish different pools (chemical and biological fractions) of carbon within the model framework.

Most extant soil biogeochemical models, such as the CENTURY and Roth-C models, characterize soil carbon decomposition by its mean residence time or turnover rate through different carbon pools (Parton et al., 1987; Jenkinson, 1990) and simulate microbial dynamics and microbially driven soil organic carbon decomposition implicitly using

first order kinetics (i.e., carbon pool turnover rate; Todd-Brown et al., 2012) based on a linear relationship with the total soil carbon stock (Jenkinson et al., 1987; Parton et al., 1988). These models typically define soil organic carbon (SOC) in one or more carbon pools and use first-order linear decay rates modified by environmental factors to simulate the biogeochemical reactions between the pools and the fluxes to the atmosphere (Fig. 2A). These models therefore do not explicitly resolve the microbial dynamics that ultimately regulate and drive soil carbon cycling (Allison et al., 2010; Conant et al., 2011; Cotrufo et al., 2013; Schmidt et al., 2011; Six et al., 2006; Treseder et al., 2012).

In these first-order kinetics models, prescribed parameters are used to describe environmental sensitivities of SOC decomposition processes – such as using a fixed Q₁₀ (temperature sensitivity) factor, where a Q₁₀ value of 2.0 is used to prescribe a doubling of SOC decomposition with every 10 °C rise in soil temperature (Davidson and Janssens, 2006). Such prescribed environmental dependencies imply that the simulated biogeochemical processes are assumed to function optimally with available substrates and a fixed concentration of microbial enzymes (i.e. a non-growing biomass). However, in nature, soil physical and biological conditions are heterogeneous and change with time and between environments – leading to deviations from such prescribed relationships. Soil microbial respiration changes dynamically across seasons (Raz-Yaseef et al., 2017), and with climatic variables such as snow cover (Yi et al., 2020), making respiration poorly approximated by the steady loss rates often included in biogeochemical models to represent pools of soil microbial carbon (e.g., Li et al., 1992, 1994; Wang et al., 2010). Measurement techniques that accompanied the earlier versions of soil biogeochemical models provided valuable knowledge and validation data on total soil carbon stocks and soil carbon fluxes. The emergence of molecular tools including gene sequencing now enable more detailed information, including microbial biomass content, function, and enzymatic activity to be coupled with model simulations (Schneegurt et al., 2003; Frey et al., 2008; Bradley et al., 2016b; Guo et al., 2020).

3.1.2. Soil Microbial Community Dynamics

The next generation of soil biogeochemical models specifically address the simplification of first-order-decay models by explicitly introducing microbial carbon into the set of defined carbon pools (Fig. 2B). Microbially-explicit models define microbial biomass as one or more explicit pools whose dynamics (growth, death, dormancy etc.) may be sensitive to a continuum of biotic and abiotic factors. Some microbially-explicit soil biogeochemical models have been tested for applications at global scales, such as the Microbial-MIneral Carbon Stabilization (MIMICS) model (Wieder et al., 2015), the Carbon, Organisms, Rhizosphere, and Protection in the Soil Environment (CORPSE) model (Sulman et al., 2014), and CLM-Microbe model (He et al., 2021). These models have, in general, performed well when compared with contemporary observations. Yet, importantly, very few microbially-explicit soil models have been applied specifically to the Arctic region. A previous application of MIC-TEM in the Arctic (Zha and Zhuang, 2018) with and without an explicit microbial formulation, showed that simulations with an explicit description of microbial biomass predicted more carbon to be released from soils during the 21st century compared to those with a microbially-implicit description.

In ‘microbially-explicit’ model structures, the microbial biomass carbon pool constitutes a rate-limiting factor in the SOC decomposition process, and thus both biotic and abiotic factors together strongly influence the simulated soil respiration behavior (Sierra et al., 2011; Zha and Zhuang, 2018). However, even among models that include explicit representation of one or more microbial biomass pools, such as the DAYCENT (Wang et al., 2010), MOMOS (Pansu et al., 2004, 2010) and DNDC (Li et al., 1992, 1994) models, microbial biomass pools are often not used to drive decomposition reaction rates; rather, they are defined as simply another organic carbon pool (distinct from non-living organic carbon) with an independently prescribed loss rate. These models thus may lack important and climatologically-sensitive biogeochemical




Fig. 2. Diagrams for the formulation of microbial soil carbon models. (A) A microbially-implicit soil organic carbon (SOC) degradation model; (B) A microbially-explicit model; (C) A microbially-explicit model where microbial biomass is represented in distinct pools for active and dormant biomass (B_{active} and B_{dormant} respectively), SOC is divided into polymers and monomers, and enzyme production and consumption processes are explicitly resolved. State variables are represented by solid-border shapes and derived variables (i.e. CO_2 production) are represented by a dashed-border shapes. The consumption/degradation of organic carbon (to microbial biomass and CO_2) is represented by solid green arrows. The contribution of necromass (i.e. dead microbial biomass) to SOC is represented by dashed blue arrows. Enzyme production and consumption is represented by solid red arrows. The transformation of polymers into monomers is represented by the solid black arrow. The transfer of microbial biomass between active and dormant states (i.e. transition into and out of dormancy) is represented by dashed black arrows. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

mechanisms that might be centrally involved in climate feedbacks.

Extant microbial soil models often define carbon pools along the reaction chain:

1. Input carbon pool(s). Input carbon represents the fresh substrate source for soil microbial decomposition from above-ground and surface environments, and usually consists of vegetation litter-fall and exudates. These inputs can be further divided depending on their reactivity (e.g. labile, refractory), and/or their molecular complexity (e.g. monomers, polymers) (Fig. 2C) (Allison, 2012; Tang and Riley, 2015; Wieder et al., 2015);
2. Microbial biomass pool(s). Some soil models include microbial biomass as a defined pool and thus it is explicitly resolved within the soil biogeochemical reaction framework. Here, the fraction of soil carbon comprised by microbial biomass is defined within one or more state variables (i.e. stores): simulated as one pool in some models (Allison et al., 2010), or in several discrete pools, for instance, divided according to function (Wieder et al., 2015; Bradley et al., 2015), and/or according to carbon assimilation strategy (e.g. structural microbial biomass and reserve microbial biomass pools) (Tang and Riley, 2015; Zha and Zhuang, 2018);
3. Extracellular enzyme pool(s). Extracellular enzymes play a vital role as catalyst in carbon decomposition process (Allison et al., 2010; Grant et al., 2001; He et al., 2014; Parton et al., 1987; Tang and Riley, 2015, 2019), can show distinct seasonal patterns in cold environments (Weintraub et al., 2007), and can be explicitly resolved in models of soil carbon in order to more accurately represent how soil carbon cycling and microbial-enzymatic decomposition is affected by environmental factors – including those that vary seasonally. Where enzymes are explicitly resolved within soil biogeochemical models, the abundance of enzymes depends on microbial biomass (a portion of microbial carbon uptake is allocated to enzyme production) and environmental conditions (Schimel and Weintraub, 2003); therefore enzyme-explicit models typically apply an empirical relationship between microbial biomass and enzyme production. Enzyme production rates are also constrained by physical factors to simulate litter decomposition at each time step (Allison et al., 2010; He et al., 2014; Tang and Riley, 2015).

Soil models compute the reactions between these carbon pools, including the depolymerization (i.e. degradation) of SOC, the uptake of substrates (i.e., microbial C assimilation), microbial growth and turnover, and the production and turnover of enzymes. Inputs and outputs are also computed, including litter carbon (polymeric compounds including cellulose, hemicellulose, lignin, monomers including intracellular material, easily-leached monomeric compounds and root exudates). Some soil biogeochemical models distinguish different molecular weights of input carbon, so that monomers are assimilated by microbes (part of which is respired, the rest goes into reserve microbial pool that supports maintenance, enzyme production, and microbial growth), and polymers are degraded by enzymes to lower molecular weight carbon (Tang and Riley, 2015, 2019). Other soil biogeochemical models distinguish input carbon by their chemical quality (e.g., metabolic versus structural litter inputs) (Wieder et al., 2015).

Carbon pools must be sufficiently represented in models so that model behavior can adequately capture both the steady-state and the dynamic nature of biogeochemical processes occurring in the natural environment. Thus, as factors such as carbon quality, microbial biomass, and enzyme activity change over seasonal cycles, the simulated soil carbon and microbial dynamics will be reliably captured.

3.2. Modelling microbial dynamics in Arctic soils and considerations for year-round simulations

Certain applications of microbially-explicit models to year-round studies of Arctic soils may require consideration of dynamic processes that are not currently resolved or poorly resolved in existing soil biogeochemical model frameworks. Current soil biogeochemical models, for example, are known to poorly constrain methane emissions, in particular during the shoulder and winter seasons (Treat et al., 2018).

As new data and understanding of the year-round processes occurring in Arctic soils emerges, as well as the recognition of previously unknown sensitivities to changes in climate, it is important to revisit soil biogeochemical modelling frameworks to assess where developments and improvements can be made.

3.2.1. Microbial Growth, Maintenance, and turnover

Soil microbial dynamics are intricate and complex; however, various fundamental biological processes are recognized by most existing microbially-explicit soil biogeochemical models, including microbial growth, maintenance (i.e., energy/substrate use for processes other than growth) and turnover (i.e. death/mortality). Conceptual and numerical representations for microbial growth and maintenance are, in general, built on two distinct steps: (1) substrate uptake and (2) substrate assimilation.

Microbially-explicit modelling approaches must first define rules to determine how and at what rates microorganisms take up and assimilate SOC. There are three general categories of mathematical expressions for uptake and assimilation of SOC in extant models: (1) Michaelis–Menten kinetics (MM, Michaelis and Menten, 1913), or Monod kinetics (Monod, 1949), (2) reverse Michaelis–Menten kinetics (r-MM, Schimel and Weintraub, 2003), and (3) a combination or generalization of the original and the reverse MM kinetics. Conceptual diagrams representing MM and r-MM kinetics are displayed in Fig. 3.

The original MM kinetics assumes that substrate (SOC) concentration exerts a limitation on substrate uptake, and thus on growth rate. The rate of substrate uptake is calculated according to:

$$v = \frac{V_{max} \cdot S}{K_S + S} \quad (1)$$

where v is the actual substrate uptake rate, V_{max} denotes the maximum substrate uptake rate, K_S represents the half saturation constant, and S is the substrate concentration.

The r-MM kinetics, alternatively, assumes that enzyme concentration exerts a limit on substrate uptake rate, and the rate of substrate uptake is calculated as:

$$v = \frac{V_{max} \cdot E}{K_E + E} \quad (2)$$

where E is the enzyme concentration, K_E represents the half saturation constant for enzyme concentration and V_{max} is the same of Eq. (1).

MM and r-MM kinetics have been applied to improve soil biogeochemical models with reasonable results (Allison et al., 2010; Drake et al., 2013; Schimel and Weintraub, 2003; Sulman et al., 2014; Wieder et al., 2015). However, it is important to note that in the natural environment, substrate and enzyme conditions can be highly heterogeneous across the various temporal and spatial scales of observation, measurement, and model application. Moreover, discrepancies still exist among and between model simulations, as well as between model simulations and observations.

Microbial growth efficiency (the ratio of growth to assimilation) and carbon use efficiency (CUE, the ratio between carbon retained by microbes in the microbial biomass pool and enzyme production, and the carbon taken up by the microbes) are often used to describe microbial assimilation of carbon substrate (Manzoni et al., 2012; Tang and Riley, 2015). These terms are essentially synonymous among many soil numerical models. As substrate transits through the soil and is captured by a microorganism, a fraction of the substrate that is assimilated is utilized to support the microbial maintenance requirement - usually calculated as proportional to the microbial biomass pool and scaled by microbial turnover and growth rates. Model simulations often assume that respiration loss is only due to carbon released during the assimilation process. More dynamic models consider respiration loss during the assimilation and maintenance processes (Grant et al., 2001; Petersen et al., 2005; Sitch et al., 2003; Tang and Riley, 2015). One commonly used approach




Fig. 3. Conceptual diagram of the mechanisms described for soil microbially explicit modelling. (A) MM-kinetics, (B) reverse MM-kinetics, and (C) dormancy switch.

considers maintenance as an additional consumption of substrate aside from the substrate that is consumed by microbial growth (Pirt, 1965), whereas another approach considers maintenance as a catabolic process (i.e. derived from respiration/consumption of cell biomass carbon stores) (Herbert, 1959). After maintenance activities, the residual substrate may then be partitioned into growth and enzyme production, computed as being proportional to their maximum potential rates (see Fig. 2).

Most microbially explicit models capture the entire microbial community within a single basic unit – the microbial carbon pool, while

some models further distinguish explicit pools for structural microbial biomass and reserve microbial biomass (Tang and Riley, 2015). Major advantages of the structural/reserve representation are that it allows for (i) an explicit representation of the allocation of carbon between maintenance and growth within simulated microbial communities, and (ii) a storage mechanism that becomes important in regulating microbial activity when carbon supply changes. This description of microbial biomass may be particularly influential to microbial growth and carbon fluxes considering seasonal changes in carbon availability in Arctic soils arising from biotic (e.g. litter input) and abiotic (e.g. freeze-thaw) changes. Many Arctic soil ecosystems, however, such as tundra, are not typically rich in soil litter carbon supplies from upper layers of vegetation (i.e. from mosses and short vascular plants), and thus a significant microbial reserve would not ordinarily be predicted in these systems (Elmendorf et al., 2012; Lloyd, 2001; Meentemeyer et al., 1982; Uchida et al., 2010; Villarreal et al., 2012).

Rates of microbial maintenance, growth, and turnover are expected to vary depending on seasonal changes in major environmental characteristics. Maintenance respiration is relatively commonly considered in microbially explicit models. It is usually prescribed in a linear relation to total active microbial biomass, but can also be empirically determined and/or scaled with environmental factors such as temperature and soil moisture (Moorhead and Sinsabaugh, 2006; Petersen et al., 2005; Schimel and Weintraub, 2003; Wang et al., 2014). Growth respiration is another commonly considered loss flux in many models and is generally presented as being proportional to the soil carbon decomposition flux (Moorhead and Sinsabaugh, 2006; Pansu et al., 2004; Tang and Riley, 2015). Similar to descriptions of maintenance cost, some models assume that this respiration flux is influenced by environmental factors (temperature, soil moisture, etc.) and scaled according to certain relationships with these variables (Tang and Riley, 2015). A turnover rate is usually prescribed to microbial pools in soil biogeochemical models to simulate microbial mortality. This mortality term is often dependent on microbial density or population size (Tang and Riley, 2015; Wieder et al., 2015). Microbial necromass (i.e. dead microbial biomass) will typically be cycled back into SOC pools – making this carbon available as substrates for decomposition. Microbial growth, maintenance and turnover are explicitly described in many soil biogeochemical models, and process rates are commonly linked to factors such as temperature, moisture availability, and light. Seasonal variation in these factors thus drives responses in the microbial community dynamics and resulting transformations of carbon pools (including organic carbon degradation and greenhouse gas production). Therefore, accurate representation of these processes and their dependencies on environmental variables (constrained, for example, by empirical measurements) will be critical in accurately simulating soil carbon cycling in Arctic soils on a seasonal basis.

3.2.2. Traits of soil microorganisms

The traits of soil microorganisms are typically described numerically by assigning various sets of parameters to each defined biomass pool (e.g. bacteria, fungi), and the values of these parameters can be adjusted to represent sensitivities to different environmental factors. Manzoni and Porporato (2007), for instance, demonstrated that bacteria and fungi respond differently to soil moisture availability: fungi are better suited to cope with water stress, while bacteria are more sensitive to changes in soil moisture. Other studies have shown that rates of methanogenesis are more sensitive to temperature ($Q_{10} \approx 4$) than methanotrophy ($Q_{10} \approx 2$) (Segers, 1998; Le Mer and Roger, 2001), although high-affinity methanotrophs in Arctic soils are strongly temperature sensitive at temperatures lower than 5.6 °C ($Q_{10} \approx 8$) (Lau et al., 2015). In models, these behaviors can be shaped by specific algorithms and parameter values that are prescribed to groups of biomass (which are often represented by distinct state-variables/groups/pools). Given the distinct physiological and functional responses of different microbial groups to the various compounding and fluctuating environmental stresses (low

temperatures, frequent freeze-thaw cycles, low or fluctuating water availability) experienced by Arctic soil microorganisms, representation of distinct microbial pools in soil models to simulate community composition might enable more accurate simulation of activity, resistance and function of Arctic soil microbes to seasonal and other environmental changes.

Similarly, a dual-microbial biomass pool structure – which would comprise of distinct groups of biomasses with different growth and adaptation strategies – may be used to capture observed seasonal changes in Arctic soil respiration rates. Models can explicitly resolve microbial community composition by representing bacterial and fungal pools that respond differently to the changing environment. Such approaches have been applied to describe the priming effect in soil – whereby soil organic matter turnover changes due to substrate addition in the short term (Kuzryakov et al., 2000), effectively enabling the simulation of two microbial groups with different traits that compete for substrate with different growth strategies: r-strategists that grow quicker using more easily accessible substrates, and k-strategists that grow slower using resources more efficiently (Fontaine and Barot, 2005; Pianka, 1970). Ultimately, considering the high degrees of seasonal change in Arctic soils, models should capture variability in the traits of organisms to seasonal changes in environmental conditions, and in the biomass stock of organisms with different traits across seasons.

3.2.3. Functional groups

Some soil biogeochemical models distinguish distinct microbial functional groups. In such a ‘functionally-explicit’ model, microorganisms are classified into distinct biomass pools (represented numerically by separate state-variables) based on their metabolic function.

A common example among soil biogeochemical models is the distinction between CO₂ producing aerobic heterotrophs, and CH₄ generating anaerobes (i.e. methanogens). Methane dynamics are simulated in some soil biogeochemical models – in particular for wetlands (Cao et al., 1995; Christensen et al., 1996; Fan et al., 2013; Grant, 1998; Riley et al., 2011; Treat et al., 2018; Wania et al., 2010). Some numerical soil biogeochemical models treat CH₄ implicitly – where processes such as methane production are simulated as a portion of total heterotrophic respiration (Xu et al., 2016; Wania et al., 2010), or as a proportion of litter/SOC decomposition (Cao et al., 1995; Zhuang et al., 2004). Alternatively, some models treat methane dynamics explicitly, for example, in Segers’ model (Segers and Kengen, 1998) and the ‘ecosys’ model (Grant, 1998), where CH₄ production is captured as anaerobic carbon mineralization of acetate (in saturated environments). Methane production in anaerobic soils is generally simulated as a function of an ecosystem-specific maximum potential production rate and limited by substrate availability, soil temperature, pH, and soil redox state (Xu et al., 2016). CH₄ produced in soil may be oxidized as it diffuses through the soil profile (Zhuang et al., 2004; Oh et al., 2020). This process can be captured in models using specific algorithms for aerobic methanotrophy which generally fall into two categories (both of which use dual Monod Michaelis-Menten-like equations): (1) CH₄ oxidation as a function of soil CH₄ concentration, temperature, and soil moisture (Tian et al., 2010; Xu and Tian, 2012; Zhu et al., 2014); and (2) CH₄ oxidation as a function of soil CH₄ and O₂ concentration, temperature, and soil moisture (Arah and Stephen, 1998; Riley et al., 2011). Models that explicitly simulate methane dynamics often show large discrepancies between model predictions and empirical data (Bohn et al., 2015; Melton et al., 2013; Treat et al., 2018).

It is well established that the hydrological regime of Arctic soils, including the prevalence of water-logged conditions in thawed soil layers, varies on a seasonal basis (Hodson et al., 2019; Joabsson and Christensen, 2001), with implications for soil redox status and dominant carbon degradation pathways including CO₂ and CH₄ production (see Section 3.3.3). A study in a seasonal lagoon in Svalbard showed that methane concentrations vary from over 10 mg/L in the transitional spring period to below 0.005 mg/L in summer (Hodson et al., 2019).

Another study in a seasonal wetland in Greenland showed that CO₂ and CH₄ fluxes change significantly during the summer thawed period (Joabsson and Christensen, 2001). For certain modelling applications, therefore, it is important that soil redox states (including transitions between oxic and anoxic conditions) and the corresponding dominance of microbial groups (e.g. aerobic versus anaerobic heterotrophs) and decomposition products (e.g. CO₂ versus CH₄) are captured by soil biogeochemical models - potentially necessitating the explicit consideration of different functional groups in soil biogeochemical models.

The majority of microbially explicit soil models do not explicitly simulate nutrient cycling (Allison et al., 2010; Tang and Riley, 2015; Wieder et al., 2015), although N and P cycling processes are included in some more recent model applications (Huang et al., 2021; Yu et al., 2020), and nutrient availability may be important to capture year-round considering the role of seasonal changes in nutrients (arising from litter inputs, precipitation and hydrological pathways, weathering, and other factors) on microbial activity and carbon cycling.

3.2.4. Microbial Dormancy

Dormancy – despite its ecological significance, is often overlooked in soil modelling studies (Lennon and Jones, 2011). Most microbially-explicit models assume a single pool representing total microbial biomass, and sensitivities to environmental conditions are prescribed by variations in the growth and mortality terms. Typical soil models therefore consider microbial cells to be either alive and active, or dead. However, as has been described in Section 2.1, a large fraction of the microorganisms in natural soils may be in a metabolically inactive state, and the proportion of active versus inactive (i.e. dormant) microorganisms is subject to change depending on environmental conditions. Thus, incorporating dormancy into microbial models may lead to more accurate and realistic simulations of microbial activity, respiration, and carbon and nutrient cycling – especially in settings where environmental harshness fluctuates, such as with seasonality in Arctic soils and other high latitude systems.

There are two general approaches for capturing the different physiological states of microbial biomass in numerical models.

The first approach is to regard the ‘fraction of active biomass’ (i.e. the ratio of active biomass to total living biomass, Fig. 2C) as a state variable (e.g., Panikov, 1995; Blagodatsky and Richter, 1998; Ingwersen et al., 2008). The value of this state variable can be influenced by environmental conditions – enabling a dynamic response of the active/dormant fraction of microbial biomass to factors such as substrate limitation and temperature.

The second approach is to separate the total live biomass into two (or more) explicit pools, considering active biomass and dormant biomass as distinct state variables (e.g. Bär et al., 2002; He et al., 2015; Stolpovsky et al., 2011). Like the first approach, this second approach also enables microbial biomass to respond dynamically to changes in the environment, but it requires more parameters to describe it. He et al. (2015) explicitly simulated active and dormant microbial biomass pools, calculating the transition between the two states by introducing a parameter representing substrate availability to microbes, which is controlled by substrate diffusivity and soil moisture content. A general formulation of the second approach, i.e. the explicit representation of active and dormant microbial pools, is:

$$\frac{dB_A}{dt} = F_D - R_G - R_M + R_{DA} - R_{AD} - M_A \quad (3)$$

$$\frac{dB_D}{dt} = R_{AD} - R_{DA} - M_D \quad (4)$$

where B_A and B_D represent active and dormant microbial biomass, respectively, F_D is monomer uptake, R_G and R_M are growth and maintenance rates, M_A and M_D represent cell mortality, and R_{AD} and R_{DA} represent the transitions between active and dormant microbial pools. Here, only the active microbes take up monomers from litter inputs and

allocate carbon for growth, maintenance, and enzyme production. Dormant microorganisms take up and also allocate carbon towards maintenance activities. Studies suggest that the maintenance coefficient of dormant microbes can be two to three orders of magnitude lower than that of the active fraction (Anderson and Domsch, 1985a, 1985b). In most models that explicitly resolve active and dormant microbial biomass, both biomass pools (active and dormant), as well as enzymes (if they are explicitly resolved), decay due to mortality and natural breakdown processes, returning carbon to monomer and polymer pools (Fig. 2). In the ‘explicit pools’ approach, rates for the reversible processes of activation and deactivation can be prescribed (Locey, 2010; Ayati, 2012), or calculated based on sensitivity to environmental factors – a phenomenon known as ‘responsive’ switching (Lennon and Jones, 2011; Bär et al., 2002). The model of Manzoni et al. (2014) prescribed a dormancy-dependance based on osmoregulation – suggesting that increases in osmolyte concentration (that are described numerically by a water potential parameter) trigger a transition to dormancy (Manzoni et al., 2014). As well as ‘responsive switching’, microorganisms may also transition into and out of dormancy by ‘spontaneous’ switching, even under stable environments (Kussell and Leibler, 2005; Lennon and Jones, 2011; Piggot and Hilbert, 2004).

For model applications to high latitude Arctic soils, the ‘responsive’ dormancy strategy might be most suitable to realistically capture the variability in microbial activity and associated biogeochemical processes on a seasonal basis – considering the fluctuating environmental conditions (i.e. moisture, temperature, light, substrate inputs) encountered by soil microorganisms across seasonal (as well as shorter and longer) timescales. The freezing of soils not only produces ‘harsh’ or stressful conditions to soil microorganisms, but also limits the physical transport of substrate (and thus energy sources to microorganisms) into the soil. Seasonal patterns in microbial state-switching may be critical in shaping year-round soil carbon decomposition processes and thus the overall carbon budget of the Arctic. Resolving the effects of environmental factors (such as temperature and soil moisture) on triggering the transitions of soil microorganisms between active and dormant states, as well as imposing physical limitations to substrate transport, likely will lead to improvements in the general understanding and simulation of year-round microbial processes in Arctic soils.

3.2.5. Extracellular enzymes

Enzyme production is explicitly calculated in some microbially-explicit models as being proportional to the size of the microbial biomass pool. Enzyme production rates can be prescribed as a function of the magnitude of the total or active microbial biomass pools. Enzyme loss rates can similarly be prescribed as a turnover rate (He et al., 2015; Zha and Zhuang, 2018). Enzyme-catalyzed SOC decomposition can be viewed from the perspective of Langmuir adsorption, and reaction rates can be described as a maximum reaction rate scaled by temperature dependence, such as an Arrhenius equation (Davidson et al., 2012). This approach results in enzyme-catalyzed SOC decomposition rates that are more sensitive to environmental factors than the fixed Q_{10} -equation approach used by models assuming first-order kinetics. Prescribing sensitivities of enzyme production and loss to seasonally-variable environmental factors will be important to accurately capture year-round dynamics in enzyme-explicit models.

3.3. Incorporating physical properties and processes into soil biogeochemical models

The dynamics of the active layer and therefore, the activity of soil microorganisms and associated carbon transformations in permafrost are strongly shaped by soil physical properties, and influenced by snowpack and vegetation dynamics, which, in Arctic systems in particular, vary considerably on a seasonal basis (Rasmussen et al., 2018; Wei et al., 2021; Jin et al., 2022; Keuschnig et al., 2022). Here, we describe some of the most important physical characteristics that can be

incorporated into soil biogeochemical models to enable year-round simulations of microbial and biogeochemical processes in the Arctic. The long-term goal is to use continuous soil- and geophysical observations as input data to inform more realistic soil biogeochemical modelling.

3.3.1. Soil temperature distribution and effects on microbial activity

Thermal conductivity in permafrost determines the speed of heat transfer through the soil, and thus, governs temperature distribution with depth and the rate of thawing. For numerical models to capture the thermal conductivity of frozen soils, they need to account for:

- (i) Soil porosity, bulk density and compaction. Soils with higher porosity tend to have lower thermal conductivity because the pores act as insulating voids, hindering heat flow. Hence, soil compaction with depth generally leads to increased thermal conductivity (Angelopoulos et al., 2020).
- (ii) Particle size and distribution (i.e. soil texture), and mineral composition: coarse-grained soils, such as sand and gravel, generally have higher thermal conductivity due to better heat conduction between larger particles compared to fine-grained soils. Additionally, different minerals have different thermal conductivities and heat capacities. For example, soils with high quartz content generally have higher thermal conductivity than those with higher clay content (Ye et al., 2022).
- (iii) Gas and liquid content: dry or water un-saturated soils have lower thermal conductivity than water-saturated ones because of the higher thermal conductivity of water compare to air (Dong et al., 2015).
- (iv) Salinity: pure water has relatively low thermal conductivity compared to saltwater.
- (v) Ice and organic matter content: increasing ice and organic matter content within the soil structure have opposite effects on its thermal conductivity. While ice content increases the thermal conductivity of the composite soil (Chuvilin et al., 2021), organic matter acts as an insulating material (Bruin et al., 2023). This thermal coupling is thought to be critical for projecting future permafrost dynamics, since the response of soil temperature to rising air temperature would be significantly impacted by changes in organic matter content related to microbial activity (Zhu et al., 2019).

Soil temperature has long been recognized to exert a significant control on microbial activity and carbon cycling – including affecting microbial enzyme activity, organic matter decomposition rates, growth rates, maintenance costs, and in fact, most biogeochemical reactions in soil ecosystems (Burns et al., 2013; Davidson and Janssens, 2006; Mu et al., 2016). Considering the limitation that temperature imposes on Arctic soil microbial activities, including the impact of freeze-thaw processes and substrate transport, a realistic representation of temperature sensitivity is especially important for achieving robust and accurate estimates of seasonal carbon dynamics in Arctic soils.

One of the simplest formulations used to explain the temperature dependence of biochemical reactions in numerical modelling is a linear relationship with parameters extracted from linear regression on temperatures, as used in DAYCENT and LPJ models (Del Grosso et al., 2002; Wania, 2007). Another earlier formulation used is the Q_{10} temperature coefficient. This method applies an empirical coefficient to prescribe an exponential relationship between reaction rates and temperature. Q_{10} values typically vary between 1.3 and 3.3 – effectively prescribing an increase in reaction rates by a factor of 1.3 to 3.3 per 10 °C rise in temperature (Bekku et al., 2003; Raich and Schlesinger, 1992). A later commonly used metric describing the sensitivity of microbial and enzyme activity to temperature is the Arrhenius equation. Whilst the Q_{10} method is based on a single parameter that prescribes the slope between rates and temperature, the Arrhenius equation requires both an

activation energy and an exponential factor (Arrhenius, 1889). Both temperature sensitivity metrics are, in effect, first derivatives of the temperature (Singh and Gupta, 1977), and assume that the activation energy needed for biochemical reactions is constant within a given temperature range (Allison et al., 2018). In fact, in the Michaelis-Menten formulation of decomposition, temperature sensitivity is already established in rate constants: V_{max} depends on temperature following the Arrhenius equation, and K_S, K_E depend on temperature following a linear relationship (Todd-Brown et al., 2012). This assumption has been questioned by recent studies providing evidence that enzymatic heat capacity might change with temperature (Hobbs et al., 2013). New theories such as macromolecular rate theory (MMRT) have been proposed to explain the varying activation energy with temperature (Alster et al., 2016; Hobbs et al., 2013). Nevertheless, it has been suggested that when temperatures are close to zero, both Arrhenius theory and MMRT yield very similar Q_{10} values (Allison et al., 2018). Therefore, it may be more important to consider applying the more complex MMRT to ecosystems with higher and more variable temperature ranges. These theories can be formulated mathematically as:

$$\text{Linear : } r = r_0 \cdot (aT + b) \quad (5)$$

$$Q_{10} : r = r_0 \cdot Q_{10}^{(T-T_0)/10} \quad (6)$$

$$\text{Arrhenius : } r = Ae^{-E_a/R \cdot T} \quad (7)$$

$$\begin{aligned} \text{MMRT : } \ln(r) \\ = \ln\left(\frac{k_B \cdot T}{h}\right) - \frac{\Delta H_{T_0} + \Delta C_p(T - T_0)}{R \cdot T} + \frac{\Delta S_{T_0} + \Delta C_p(\ln T - \ln T_0)}{R} \end{aligned} \quad (8)$$

where r is the reaction rate, T denotes the temperature, T_0 is the reference temperature, E_a represents the activation energy, k_B is Boltzmann's constant, h is Planck's constant, R is the universal gas constant, H denotes enthalpy, and S stands for entropy.

3.3.2. Hydraulic conductivity and soil moisture effects on microbial activity

Hydraulic conductivity in permafrost refers to how easily fluids (i.e., liquid and gas/air) can flow through the porous media and existing fractures, thus, governing water infiltration, soil moisture content, and gas diffusion and transport (Huang et al., 2016; Heinze, 2021; Jiang et al., 2021). To capture the hydraulic conductivity of frozen soils it is important that models account for soil properties, such as pore connectivity, bulk density, organic matter content, and soil texture, which exert first-order controls on the soil's ability to store and transport fluids (Huang et al., 2016). For certain applications, models may track variations in ice content through time, as well as the consequent geo-mechanical response of the soil. For instance, the presence of ice can partially or entirely fill the soil's pore spaces, significantly reducing its connectivity and restricting fluid flow and infiltration (Larsbo et al., 2019). In contrast, thawing increases the soil's hydraulic conductivity by removing ice from pores (e.g., Watanabe and Kugisaki, 2017; Jiang et al., 2021). Yet, such an effect can be masked by soil volumetric consolidation or collapse that is expected upon thawing (De La Fuente et al., 2020; Shastri et al., 2021). Moreover, models that account for the occurrence of fractures (induced either by pore-overpressure or freeze-thaw cycles and that might act as preferential permeable paths), the formation of taliks (which generally act as preferential flow paths for water), or the effects of vegetation on soil structure (the root systems of plants can create channels and pores in the soil, which can either enhance or reduce hydraulic conductivity, depending on the type of vegetation and the depth and density of the roots) and exposure to rain (plants may act as a natural mulch, reducing erosion and increasing water infiltration) are more suitable to capture preferential pathways for fluid flow, which can dramatically increase the hydraulic conductivity of the system (Watanabe and Kugisaki, 2017; Jin et al., 2022).

Soil moisture is also known to exert a significant control on microbial activity, respiration and carbon assimilation. This is firstly due to the intrinsic nature by which soil microbes and enzymes require moisture and therefore are affected by moisture stress, and secondly, the consideration that physical transport of substrates through the soil medium, for instance active and passive movement of chemical species through pores in the soil column, affects substrate availability by limiting supply via active transport (Harris, 1981; Manzoni et al., 2014).

Most numerical models that explicitly consider soil moisture prescribe a non-linear relationship between microbial activity and moisture availability. A widely used algorithm uses water potential, which can be calculated from volumetric water content, to prescribe the empirical relationship $A(\psi)$ (Andren and Paustian, 1987; Rodrigo et al., 1997):

$$A(\psi) = \frac{\log(-\psi) - \log(-\psi_{min})}{\log(-\psi_{opt}) - \log(-\psi_{min})} \quad (9)$$

where ψ is water potential, ψ_{opt} is the water potential that yields the optimal decomposition rate and ψ_{min} represents the water potential at which microbial activities cease. A similar equation can be written as (Manzoni et al., 2012):

$$A(\psi) = 1 - \left(\frac{\log(\psi/\psi_{fc})}{\log(\psi_{th}/\psi_{fc})} \right)^\alpha \quad (10)$$

where α represents a shape parameter, ψ_{th} denotes a water stress threshold and ψ_{fc} is a reference water potential corresponds to the soil field capacity. The equation allows respiration to decrease gradually as water potential becomes increasingly negative, and empirical values of ψ_{th} and α determine the sensitivity of microbial activity to changes in soil moisture.

Some models use water content instead of water potential to describe the relationship between microbial activities and soil moisture, as shown below (Myers et al., 1982; Rodrigo et al., 1997):

$$g(\theta) = m \frac{(\theta - \theta_{ref})}{(\theta_{opt} - \theta_{ref})} + c \quad (11)$$

where θ is volumetric water content, $g(\theta)$ describes microbial sensitivity in response to soil moisture, θ_{opt} is the optimal water content for microbial activities, θ_{ref} is a reference parameter, and m and c are empirical coefficients.

Few soil models also account for the effect of water limitation on substrate diffusion and binding. However, those that do generally prescribe an effective affinity parameter to match Michaelis-Menten-type substrate uptake by microbes (Tang and Riley, 2019):

$$K_{s,w} = K_{s,w,0} \left(\frac{\nu_m \cdot \delta}{4 \cdot \pi \cdot D_{s,w,0} \cdot r_m (r_m + \delta)} + \frac{\nu_m}{4 \cdot \pi \cdot D_s (r_m + \delta)} \right) k_{s,w,1} \frac{N_{cell}}{N_A \cdot \nu_m} \quad (12)$$

where $K_{s,w,0}$ is the reference substrate affinity parameter, $k_{s,w,1}$ denotes the unicellular substrate interception coefficient, δ represents the thickness of the water film formed over the microsite of a cluster of particles, microbes and substrates, ν_m is the mean microsite volume size, N_{cell} is the number of cells per microsite, r_m represents the microbial cell size, D_s is bulk diffusivity for substrate, $D_{s,w,0}$ is aqueous diffusivity for substrate and N_A is Avogadro's number.

3.3.3. Freeze-thaw dynamics

Determining the freezing point in soils is vital to comprehend permafrost dynamics (freeze-thaw behavior) and quantify ice and unfrozen water content within the porous media (Thomas et al., 2009; Yokohata et al., 2020). Experimental tests and field observations have evidenced the effects of pore capillary pressure in narrow pores on inhibiting ice formation (De La Fuente et al., 2021). Water in frozen soils is proven to remain unfrozen at temperatures well below 0 °C due to such capillary forces, which lower the activity of the water, and,

consequently, its freezing point (Zhou et al., 2018). This is important not only because of the impact of the pore-filling species on soil's thermal and hydraulic properties, but because in permafrost, access to nutrients and the ability to eliminate waste materials are limited by the thickness of the unfrozen water films (Rivkina et al., 2000). Thus, models should not assume that soils are fully frozen below 0 °C. Instead, they should account for the dependency of the water freezing point on the soil's pore size distribution and solute content of the liquid phase (Kozlowski, 2009).

The freeze-thaw dynamics of permafrost are also pivotal in determining the thickness of both saturated and unsaturated sediment layers, thereby influencing the penetration of oxygen vertically within the soil. This dynamic control establishes a depth-equilibrium between aerobic and anaerobic oxidation processes, shaping the redox conditions and biogeochemical processes occurring within the soil (Patrick et al., 1996; Fiedler et al., 2007; Vepraskas and Faulkner, 2001). The resultant redox conditions directly impact soil carbon storage by influencing the rates and pathways of organic matter decomposition. Additionally, soil redox conditions exert an indirect effect on organic matter remineralization by affecting the bioavailability of organic molecules and nutrients in the soil (Herndon et al., 2020).

3.3.4. Snowpack dynamics

Observations indicate that the timing of snow onset and changes in snowpack depth may result in prolonged periods of unfrozen soil despite soil temperatures close to or slightly below 0 °C – thus maintaining an active un-frozen layer in which microbial activity can continue (Box et al., 2019; Outcalt et al., 1990; Zona et al., 2016). Snow has a dual effect on the soil beneath it: a short period of cooling during early onset of snow accumulation, and a long period of warming as the snow provides insulation from dropping air temperatures. The insulating properties of the snowpack may bring about different effects on soil thermal and carbon dynamics in the short-term (depending on temperature and corresponding snowpack depth), but on a longer annual basis, the warming effect due to snowpack insulation is dominant (Zhang, 2005).

Snowpacks also act as a buffer between the air and the soil - altering the soil thermal conditions and contributing to attenuated soil temperature variation as compared with near-surface air temperature (Lawrence and Slater, 2010; Osterkamp, 2007; Stieglitz et al., 2003; Rixen et al., 2022). Lawrence and Slater (2010) estimated that during the latter half of the last century, >50% of the alteration to Arctic soil temperatures may be attributed to impacts from snow variability. Multiple characteristics of the snowpack affect insulation effects and therefore soil microbial activities, including: the timing and duration of snowfall, the density, thickness, and structure of the snowpack, as well as local topography and wind conditions (Zhang, 2005; Rixen et al., 2022). In the Terrestrial Ecosystem Model (TEM), snowpack thickness is changing over the simulation time and is estimated from snow water equivalent data and empirical snow density. The thermal heat flux within snow and soil layers were calculated and iterated until the heat flux from the snowpack side balances that from the ground side (Lyu and Zhuang, 2018). This insulation effect resulted in 6.4 °C warmer soil (top 5 cm) during the non-growing season in the Arctic, and a slower transition of freeze-thaw in early spring and later fall, as compared to the model where insulation is not simulated.

Although the effect of the snowpack as an insulator to soils is considered in many land-surface models, the dynamic development of the snowpack over the cold season is often neglected (Bormann et al., 2013; Chadburn et al., 2017; De Michele et al., 2013; Ekici et al., 2015; Lyu and Zhuang, 2018). Even studies focused on the improvement of snow and soil thermal dynamics schemes seldom assess the effect of snowpack dynamics on heterotrophic respiration or microbial activity (e.g., Chadburn et al., 2015). The accumulation and densification of a snowpack alters its thermal conductivity, and together with environmental factors such as wind, could be an important control on soil microbial activity and associated biogeochemical cycles. Incorporating the

effect of a dynamic snowpack in Arctic microbially-explicit models thus may enable improved simulations of seasonal patterns of microbial activity and carbon emissions in high latitude soils – particular during the early winter and spring thawing, when the typical burst emissions are observed.

3.3.5. Latent heat and zero-curtain phenomena

The zero-curtain effect refers to the phenomenon of soils maintaining temperatures near 0 °C over extended periods of freezing or thawing due to the release and uptake of latent heat by water. This phenomenon can be observed in the active layer of permafrost soils during transition seasons (Fig. 1). During the fall transitional to winter, when soil liquid water is about to freeze, latent heat released during the freezing process may offset heat loss by conduction towards the surface layer and keep the subsurface temperature around 0 °C for some extended period of time. Latent heat also plays an important role during the spring snowmelt period, when energy from solar radiation and rising ambient temperatures is absorbed by the snowpack. Early snowmelt may keep the soil surface temperature lower than the air temperature due to latent heat energy lost from the soil to snowmelt. As the snowmelt continues and snowpack temperature stays around the 0 °C isotherm, refreezing of the meltwater (from snowmelt) releases latent heat that subsequently warms the snowpack and the soil (Zhang, 2005). The overall effect of this process on soil temperature and microbial activity is complex and still requires further investigation.

During the zero-curtain period, water contained within freezing soils may be forced to the top and bottom of the freezing fronts forming segregated ice (Fu et al., 2022), meanwhile leaving air pockets in the active layer (Arndt et al., 2020). During this period, CO₂ production may remain relatively high and can account for a substantial portion of the annual carbon balance: fall time net carbon balance can be $15.9 \pm 0.7 \text{ g C m}^{-2}$, contributing about 42% of the total annual non-summer net carbon emission (Commane et al., 2017; Oechel et al., 2014). Throughout the zero-curtain period, microbial activity may continue in unfrozen soils whilst the diffusion and advection of the produced CO₂ may be obstructed by the freezing fronts and overlying snowpack, which act as barriers. This can result in periodic bursts of carbon released as CO₂ after the spring thaw eliminates these barriers (Elberling and Brandt, 2003; Koponen and Martikainen, 2004). Importantly, zero-curtain phenomena (1) allow microbes to remain active in soils during winter, (2) change the diffusion, advection and microbial status in response to seasonal transitions, and (3) may de-couple the timing of CO₂ that is produced in soils from CO₂ fluxes measured above ground.

In order to simulate the zero-curtain phenomena observed in the Arctic, accurate representation of soil physical parameters, especially temperature, moisture, and latent heat, is likely to be necessary. Currently, many land surface models tend to underestimate soil temperature during the freezing period due to possible inadequate estimation of the effect of snow cover insulation and latent heat, which subsequently introduces freezing of soil water and inhibition of microbial activity and respiration in the model simulations that are not representative of the natural system (which may remain at least partially thawed) (Dankers et al., 2011; Nicolsky et al., 2017; Yang et al., 2018). Accurate representation of the zero-curtain phenomenon is therefore critical for improving simulations of seasonal changes in Arctic soil carbon cycling – especially since inaccurate estimates of soil temperature and freeze-thaw processes may lead to an underestimation of seasonal CO₂ emissions and inaccuracies in the overall carbon balance. A phase-change scheme incorporating latent heat could help improve the simulation of freeze-thaw processes in the cold season and year-round.

3.3.6. Multiphase transport

The permafrost is a dynamic multiphase environment, encompassing both solid (i.e., soil, organic matter, and ice) and mobile fluid phases (i.e., gas and liquid). As indicated in previous sections, tracking the evolution of liquid-gas content within the active layer may be crucial for

certain model applications. Firstly, gas within pores serves as an insulator that contributes to the long-term preservation of ice for instance warming such as during spring/summer transitions (see section 3.3.1). Secondly, its occurrence in pores is linked to the soil water content, thus influencing microbial activity, and potentially triggering sediment fractures (see section 3.3.2) Thirdly, gas content and soil permeability (which can be affected seasonally by freezing and thawing cycles) affects gas exchange between soil and atmosphere, thus, having important implications on carbon cycling (Lange et al., 2016).

Models that incorporate multiphase transport may be particularly suitable for assessing the fate of microbially-produced CO₂ and CH₄ and, thus their ultimate impact on carbon cycling and climate (De La Fuente et al., 2022). Multiphase transport models are also capable to account for additional sources of gases to the system, which may include injection via filtration along faults from depth (Khimenkov and Stanilovskaya, 2022) or due to destabilization of permafrost-associated hydrates (Ruppel and Kessler, 2017).

3.4. Model implementation

The application of soil biogeochemical models to the simulation of year-round dynamics of Arctic soils will require certain implementation considerations, including the spatial and temporal dimensions to be covered, the resolution to be achieved, and the types and characteristics of data that are used to initialize and validate model simulations. Here, we briefly discuss implementation issues to consider specifically for year-round applications of soil biogeochemical models.

3.4.1. Temporal resolution and time step

The temporal resolution of model output can generally be adjusted depending on the timescale of the particular processes of interest – down to a minimum time step that is commensurate with the numerical implementation of the mathematical solver used to execute the model. Most existing soil biogeochemical models are run at hourly or daily time steps, and provide output of simulated fluxes at hourly, daily or monthly intervals (Allison et al., 2010; Tang and Riley, 2015; Sulman et al., 2014; Wieder et al., 2015; Zhuang et al., 2001).

A timescale of hours or days is likely suitable to allow model simulations to capture responses to diurnal and seasonal changes, including during the zero-curtain period, when soil conditions transition between frozen and thawed states in a matter of days (Belokopytova et al., 2021; Boike et al., 2018; Hinkel et al., 2001). Data collected at the Bayelva permafrost monitoring site in western Svalbard indicates that variability in surface soil temperature largely correspond to air temperature. At this site, temperature variation in near-surface soils is relatively minor during the summer (constant light) and winter (constant dark) periods, but hourly variability and diurnal cycles are pronounced during transitional seasons (Boike et al., 2018), where the temperature of uppermost soil layers can fluctuate above and below freezing during day and night respectively (Boike et al., 2018). Thus, models with hourly temporal resolution may capture the rapid freezing and thawing and accordingly trigger microbial growth, turnover and dormancy responses. Coarser time-steps (such as with daily or monthly averages) would not capture fluctuations in environmental conditions (such as freeze-thaw) occurring on shorter timescales, and therefore responses in the microbial activity and associated carbon transformations may be inaccurate. Thus, an hourly temporal resolution may suit the model equilibrium assumption whilst also allowing for the model to capture short and longer-term variability in the carbon flux observed at various monitoring sites. Daily time intervals, whilst not fully capturing diurnal oscillations in temperature or associated soil carbon fluxes (in particular during transition seasons) may nevertheless be suitable to capture major seasonal patterns.

The temporal resolution of available forcing data in the study area is also an important factor to consider. When finer-resolution temporal forcing data are not available, it is common practice to interpolate

coarser-resolution data as forcing data to drive a model. For example, linear interpolation methods were used to transform the monthly CRU climate dataset into daily intervals to study the carbon dynamics of Alaska wetlands (Lyu et al., 2018). The Arctic region generally lacks year-round field observations, in particular due to logistical and practical difficulties in making continuous measurements during the snow-covered/winter period. Interpolated meteorological data may be the only available data for the cold season, and importantly, interpolations may not capture rapid responses to abrupt environmental changes. Year-round meteorological data (with hourly to daily resolution) are now increasingly available at high-latitude sites (Belokopytova et al., 2021; Boike et al., 2018, 2019; Hartl et al., 2020), providing suitable forcing data to models. This improved forcing data, combined with suitably fine temporal resolution in models, will likely bring about improvements to modelling seasonal changes in carbon and microbial activity.

As a word of caution, it is important to note that imposing a higher temporal resolution in modelling incurs a higher computational cost, and thus the trade-offs between model resolution and computational cost must also be considered.

3.4.2. Spatial dimensions: Vertical

Many soil-biogeochemical models are 0-D (i.e. not spatially resolved), with a box-model structure that does not discern any vertical structure or differences (Koven et al., 2013). Some simplified and earlier soil biogeochemical models consider only the top organic layer as a single fixed SOC profile and assume that the deeper mineral layer does not significantly impact biogeochemical reactions (Baudena et al., 2007; Brandes and Den Hollander, 1996; Klepper and den Hollander, 1999). Beyond single layer box models, discrete layers of soil may be represented in multi-layer box models, where each vertically-stacked box contains prescribed initial conditions and the reactions unfold without any vertical exchange of carbon (Kuc, 2005). Vertical heterogeneity in soil characteristics and resulting carbon cycling is considered in the Community Land Model (CLM, Koven et al., 2013; Lawrence et al., 2011) and RothC model (Coleman and Jenkinson, 1996; Jenkinson and Coleman, 2008), capturing vertical advection and diffusion to at least 3 m depth (Koven et al., 2013; Tang and Riley, 2015). Currently, vertical SOC distribution is generally defined according to the root distribution, and trade-offs between the vertical allocation and root nutrient uptake are not considered (Fisher et al., 2012; Koven et al., 2013).

The vertical characteristics of Arctic soils are physically, chemically and biologically heterogeneous, both at discrete timepoints and also considering seasonal and long-term change. For example, deeper layers of some permafrost soils may contain vast quantities of SOC which are largely buffered from seasonal temperature changes, but may become more dynamic and play an active role in soil biogeochemical cycling under future climate warming as permafrost thaw depth increases (Koarashi et al., 2012; Tarnocai et al., 2009). Past vertically resolved applications of microbially-implicit soil models to the Arctic have assessed how changes in simulated soil temperature affected the active-layer depth, and have predicted variations in overall soil carbon stores and fluxes as a result of built-in SOC-temperature sensitivities (Genet et al., 2013; Burke et al., 2017; Lyu et al., 2018; Lyu and Zhuang, 2018).

Vertically-resolved model schemes will enable seasonal changes in the depth of the active layer, the thawing and/or freezing fronts, and subsequent changes to microbial activity in the soil depth-profile to be captured. Since the upper-most layers of soil profiles are typically the most biologically active (due to the abundant microorganisms and substrates) and responsive to environmental variation, it might be advantageous to apply a finer vertical resolution in the uppermost layers and a relatively coarser vertical resolution beneath these layers to the bottom mineral layer, in order to reduce computational costs.

3.4.3. Spatial dimensions: Horizontal

Heterogeneity in soil biological, physical and chemical characteristics occurs across a landscape (i.e. laterally/horizontally) across a range

of scales from individual soil particles to plots and landscapes. These heterogeneities arise due to numerous interacting factors including geomorphological, climatological, hydrological, chemical and biological processes, often mediated by the soil, that have occurred in the past and are presently shaping the environment. Seasonal processes affecting Arctic soils may also manifest differently across spatial scales, for instance due to (micro-)topographic features, local hydrological processes including variations in the depth of the water table, and snow cover. For instance, hydrological heterogeneities are highly pronounced at the Samoylov research site in the Lena River delta, northeast Siberia, where 58% of the land area is classified as dry tundra and 17% is classified as wet tundra (Boike et al., 2019; Nitzbon et al., 2019). Seasonal permafrost thawing further contributes to horizontal hydrological heterogeneities through its dual impact on ground permeability. The increased permeability resulting from permafrost thaw facilitates lateral water movement across the landscape, influencing lateral flow patterns and fostering the creation of features such as thaw ponds and channels (Bense et al., 2012). Changes in lateral water movement impacts hydrological connectivity, enhancing the transport of suspended sediment and particulate organic matter (Lafrenière and Lamoureux, 2019; Walvoord and Striegl, 2021). On the other hand, water released from thawing can strengthen surface or deep runoff rather than storage, causing alterations to seasonal runoff patterns (Bring et al., 2016). Both of these processes can ultimately affect microbial community dynamics and biogeochemical processes (Blaud et al., 2015).

When applying a model to a regional scale and beyond, the spatial resolution of the model simulation may be limited to a large extent by the spatial coverage and resolution of the available forcing data, particularly in regions which are highly heterogeneous in soil characteristics, hydrology, and topography. Meteorological and geohydrological data are usually available either through reanalysis or remote sensing data collected by airborne platforms or satellites. These data are generally gridded at 1.0 km or 0.5° resolution, with some products available at a finer resolution of 250 m (Justice et al., 1998; Shao et al., 2011). Therefore, these are also the most common spatial resolutions used by regional soil biogeochemical models. Nevertheless, considering the horizontal scale and resolution that is resolved in soil biogeochemical models is especially important when extrapolating parameters that are calibrated at the site-level to regional scales and beyond.

Differences in the spatial resolution of soil biogeochemical models can result in different output characteristics. For example, 11 different Earth system models were driven with the same set of meteorological data and run to compare the SOC change between the end and the beginning of the 21st century in the Coupled Model Intercomparison Project Phase 5 (CMIP5) (Todd-Brown et al., 2013, 2014). These models are originally designed with different modelling grid resolution for different geographies and ecosystems, and were extrapolated to be applied to the global scale. The different models produced a wide range of simulated SOC changes ranging from carbon loss of 72 Pg C over a century, to carbon gain of 253 Pg C. In particular, high-latitude regions were subject to large discrepancies between model outputs, from carbon loss of 37 Pg C to carbon gain of 146 Pg C (Todd-Brown et al., 2014). This indicates that many models especially lack accurate simulation of high-latitude soil processes, and poorly represent heterogeneous permafrost dynamics across the northernmost land masses.

3.4.4. Validation data

The application of land surface models is critically reliant on data: used both as forcing data for model initialization and utilization, and for calibration and validation. The quantity and quality of data that is available may greatly influence the design, behavior, and performance of numerical models, and the confidence entrusted in their outputs. Both in situ and remotely sensed data may be used for forcing, calibration and validation purposes (Liu et al., 2016; Lyu and Zhuang, 2018; Tan et al., 2016; Chadburn et al., 2017). For instance, climatic data collected on-

site was combined with remotely-sensed data on snow prevalence (of the same temporal and spatial resolution) to run a biogeochemical model to quantify northern high latitude region carbon responses to soil thermal changes (Lyu and Zhuang, 2018). In NASA's ABoVe project, an array of 18 land surface models guided the identification of target variables, as well as the spatial and temporal resolution of monitoring through a suite of field, airborne and satellite measurements (Fisher et al., 2018). Interpolation or downscaling methods can also be used to accommodate the integration of multiple data types of different resolutions for model applications (Lyu et al., 2018; McGuire et al., 2018b).

Future applications of soil biogeochemical models must, in particular, consider how data can be effectively collected and used year-round to calibrate model simulations. Consideration must be given not only to which variables might be critical to measure, but also to the timespan and temporal resolution of this data. For model applications to Arctic soils, continuous vertically-resolved data on soil temperature and water content will be especially important in order to predict soil microbial activity and carbon degradation year-round. In addition, records of snow depth and snow characteristics would benefit the representation of both soil temperature and emission timing. When incorporating meteorological data, consideration should be given to seasonal variability but also inter-annual variability, including extreme weather conditions such as periodic winter-warming, or summer drought (Zhu and Zhuang, 2013). For Arctic regions, extreme events could greatly affect the soil thermal and hydrological conditions, driving subsequent changes to soil microbial activities and carbon fluxes. The use of forcing and validation datasets of past extreme events may enable a better constraint of model parameters and sensitivity.

4. Conclusions and recommendations

A better understanding of longer-term interactions between Arctic soils and permafrost, climate, microorganisms, vegetation, soil organic matter and nutrients, and snow cover, alongside a refined model depiction of subsurface conditions such as ground ice, will contribute significantly to improved modelling of soil biogeochemical processes. Here we describe the seasonal changes in Arctic soil microbiology and biogeochemistry, and the various factors that should be considered for the development and application of soil biogeochemical models year-round in Arctic settings to improve understanding of processes in Arctic soils and permafrost, and reduce uncertainty regarding the thermo-hydraulic state of permafrost and its future climate feedbacks. Ultimately, model complexity and structure should be guided by available knowledge, data, and the overarching research questions. Therefore, in practice, model development may need to be largely bespoke to each specific application. The most complex model, whilst potentially more closely resembling the complexities of the natural environment, is not necessarily the most useful. In general, a model should be as simple as possible, but not any simpler than that – a simple model may produce equally beneficial and constructive results (e.g. Manzoni and Porporato, 2009). Simple mechanistic models applied to different locations with high temporal and spatial resolution may help in the identification of the most relevant processes to be implemented and tested in global models (e.g. Todd-Brown et al., 2012), as well as explore the importance of representing interspecies and interkingdom (plant-microbe) interactions (Naylor et al., 2020). These simple mechanistic models are promising for bridging the gap between data-driven models aimed at identifying the drivers of carbon dynamics in field studies, and complex land surface models, which often fail at accurately representing practical observations despite being process-oriented. With a simpler model, the opportunity to learn about microbiome-climate feedbacks and their role on shaping seasonal dynamics may be limited. The application of any model should be viewed as a work in progress, to be constantly re-evaluated and tested in the context of the evolving data availability and mechanistic understanding. Over time, these tools will be incrementally improved and better constrained such that they become more

useful in analytically testing hypotheses, extrapolating, interpolating and constructing budgets of processes and rates, and enabling predictive capabilities.

The emergence of year-round data and the integration of this data and understanding of seasonal processes into soil biogeochemical models will be critical to understand and predict the year-round functioning and fate of Arctic soil biology and biogeochemistry, particularly under a changing climate. Year-round measurements of microbial activity rates, organic carbon uptake and remineralization, carbon dioxide and methane production and consumption, and soil thermal and hydrological characteristics will particularly help to constrain the understanding of seasonal processes and simulations of microbially-explicit models. Moreover, improvements to models and understanding rely especially on consideration of how the carbon pool magnitude, fate and cycling dynamics varies with the microbial community's taxonomic and functional trait composition, and ways in which the carbon dynamics depart from idealized models due to spatial and temporal heterogeneity of biological, geological, physical and geochemical factors. Incorporating this complexity into models, over temporal and spatial scales that are appropriate to resolve potentially important abiotic constraints on biological processes (such as the zero-curtain effect and the progression of thawing and freezing fronts with soil depth) is a formidable but vital challenge, which, if overcome, will yield vast improvements in the understanding and predictive capabilities of the year-round functioning of Arctic soils and carbon stocks.

CRedit authorship contribution statement

Zhou Lyu: Investigation, Data curation, Writing – original draft, Writing – review & editing, Visualization. **Pacifica Sommers:** Writing – review & editing, Funding acquisition. **Steven K. Schmidt:** Writing – review & editing, Funding acquisition. **Marta Magnani:** Writing – review & editing. **Mihai Cimpoiasu:** Writing – review & editing. **Oliver Kuras:** Writing – review & editing, Funding acquisition. **Qianlai Zhuang:** Writing – review & editing. **Youmi Oh:** Writing – review & editing. **Maria De La Fuente:** Writing – review & editing. **Margaret Cramm:** Writing – review & editing. **James A. Bradley:** Conceptualization, Investigation, Data curation, Writing – original draft, Writing – review & editing, Visualization, Supervision, Project administration, Funding acquisition.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

No data was used for the research described in the article.

Acknowledgements

This work was funded by the NSF-NERC (UKRI) thematic program "Signals in the Soil" (NERC grants: NE/T010967/1, NE/T010568/1; NSF grant: 1935689). JB also acknowledges support from the CNRS Chaires de Professeur Junior (CPJ) and NERC grant NE/V012991/1 during the writing of this manuscript. QZ acknowledges support from NSF grant: 1802832. MM acknowledges the Italian National Biodiversity Future Center (NBFC): National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.4 of the Ministry of University and Research; funded by the European Union – NextGenerationEU (Project code CN00000033).

References

- Aanderud, Z.T., Jones, S.E., Schoolmaster Jr., D.R., Fierer, N., Lennon, J.T., 2013. Sensitivity of soil respiration and microbial communities to altered snowfall. *Soil Biol. Biochem.* 57, 217–227.
- Allison, S.D., 2012. A trait-based approach for modelling microbial litter decomposition. *Ecol. Lett.* 15 (9), 1058–1070.
- Allison, S.D., Wallenstein, M.D., Bradford, M.A., 2010. Soil-carbon response to warming dependent on microbial physiology. *Nat. Geosci.* 3 (5), 336–340.
- Allison, S.D., Romero-Olivares, A.L., Lu, Y., Taylor, J.W., Treseder, K.K., 2018. Temperature sensitivities of extracellular enzyme Vmax and Km across thermal environments. *Glob. Chang. Biol.* 24 (7), 2884–2897.
- Alster, C.J., Koyama, A., Johnson, N.G., Wallenstein, M.D., von Fischer, J.C., 2016. Temperature sensitivity of soil microbial communities: an application of macromolecular rate theory to microbial respiration. *J. Geophys. Res. Biogeosci.* 121 (6), 1420–1433.
- Anderson, T.H., Domsch, K.H., 1985a. Determination of ecophysiological maintenance carbon requirements of soil microorganisms in a dormant state. *Biol. Fertil. Soils* 1 (2), 81–89.
- Anderson, T.H., Domsch, K.H., 1985b. Maintenance carbon requirements of actively-metabolizing microbial populations under in situ conditions. *Soil Biol. Biochem.* 17 (2), 197–203.
- Andren, O., Paustian, K., 1987. Barley straw decomposition in the field: a comparison of models. *Ecology* 68 (5), 1190–1200.
- Angelopoulos, M., Overduin, P.P., Miesner, F., Grigoriev, M.N., Vasiliev, A.A., 2020. Recent advances in the study of Arctic submarine permafrost. *Permafrost. Periglac. Process.* 31, 442–453.
- Arah, J.R.M., Stephen, K.D., 1998. A model of the processes leading to methane emission from peatland. *Atmos. Environ.* 32 (19), 3257–3264.
- Arndt, K.A., Lipson, D.A., Hashemi, J., Oechel, W.C., Zona, D., 2020. Snow melt stimulates ecosystem respiration in Arctic ecosystems. *Glob. Chang. Biol.* 26 (9), 5042–5051.
- Arrhenius, S., 1889. Über die Reaktionsgeschwindigkeit bei der Inversion von Rohrzucker durch Säuren. *Z. Phys. Chem.* 4 (1), 226–248.
- Ayati, B.P., 2012. Microbial dormancy in batch cultures as a function of substrate-dependent mortality. *J. Theor. Biol.* 293, 34–40.
- Bär, M., Hardenberg, J., Meron, E., Provenzale, A., 2002. Modelling the survival of bacteria in drylands: the advantage of being dormant. *Proc. R. Soc. Lond. Ser. B Biol. Sci.* 269 (1494), 937–942.
- Bardgett, R.D., Bowman, W.D., Kaufmann, R., Schmidt, S.K., 2005. A temporal approach to linking aboveground and belowground ecology. *Trends Ecol. Evol.* 20 (11), 634–641.
- Batjes, N.H., 2014. Total carbon and nitrogen in the soils of the world. *Eur. J. Soil Sci.* 65 (1), 10–21.
- Baudena, M., Boni, G., Ferraris, L., Von Hardenberg, J., Provenzale, A., 2007. Vegetation response to rainfall intermittency in drylands: results from a simple ecohydrological box model. *Adv. Water Resour.* 30 (5), 1320–1328.
- Bekku, Y.S., Nakatsubo, T., Kume, A., Adachi, M., Koizumi, H., 2003. Effect of warming on the temperature dependence of soil respiration rate in arctic, temperate and tropical soils. *Appl. Soil Ecol.* 22 (3), 205–210.
- Belokopytova, L.V., Zhirmova, D.F., Meko, D.M., Babushkina, E.A., Vaganov, E.A., Krutovskiy, K.V., 2021. Tree rings reveal the impact of soil temperature on larch growth in the forest-steppe of Siberia. *Forests* 12 (12), 1765.
- Belshe, E.F., Schuur, E.A.G., Bolker, B.M., 2013. Tundra ecosystems observed to be CO₂ sources due to differential amplification of the carbon cycle. *Ecol. Lett.* 16 (10), 1307–1315.
- Bense, V.F., Kooi, H., Ferguson, G., Read, T., 2012. Permafrost degradation as a control on hydrogeological regime shifts in a warming climate. *J. Geophys. Res.* 117, F03036. <https://doi.org/10.1029/2011JF002143>.
- Blagodatsky, S.A., Richter, O., 1998. Microbial growth in soil and nitrogen turnover: a theoretical model considering the activity state of microorganisms. *Soil Biol. Biochem.* 30 (13), 1743–1755.
- Blaud, A., Lerch, T.Z., Phoenix, G.K., Osborn, A.M., 2015. Arctic soil microbial diversity in a changing world. *Res. Microbiol.* 166 (10), 796–813. ISSN 0923-2508, <https://doi.org/10.1016/j.resmic.2015.07.013>.
- Bohn, T.J., Melton, J.R., Ito, A., Kleinen, T., Spahni, R., Stocker, B.D., Zhang, B., Zhu, X., Schroeder, R., Glagolev, M.V., Maksyutov, S., 2015. WETCHIMP-WSL: intercomparison of wetland methane emissions models over West Siberia. *Biogeosciences* 12 (11), 3321–3349.
- Boike, J., Juszak, I., Lange, S., Chadburn, S., Burke, E., Overduin, P.P., Roth, K., Ippisch, O., Bornemann, N., Stern, L., Gouttevin, I., 2018. A 20-year record (1998–2017) of permafrost, active layer and meteorological conditions at a high Arctic permafrost research site (Bayelva, Spitsbergen). *Earth Syst. Sci. Data* 10 (1), 355–390.
- Boike, J., Nitzbon, J., Anders, K., Grigoriev, M., Bolshiyarov, D., Langer, M., Lange, S., Bornemann, N., Morgenstern, A., Schreiber, P., Wille, C., 2019. A 16-year record (2002–2017) of permafrost, active-layer, and meteorological conditions at the Samoylov Island Arctic permafrost research site, Lena River delta, northern Siberia: an opportunity to validate remote-sensing data and land surface, snow, and permafrost models. *Earth Syst. Sci. Data* 11 (1), 261–299.
- Bormann, K.J., Westra, S., Evans, J.P., McCabe, M.F., 2013. Spatial and temporal variability in seasonal snow density. *J. Hydrol.* 484, 63–73.
- Bormann, K.J., Brown, R.D., Derksen, C., Painter, T.H., 2018. Estimating snow-cover trends from space. *Nat. Clim. Chang.* 8 (11), 924–928.
- Box, J.E., Colgan, W.T., Christensen, T.R., Schmidt, N.M., Lund, M., Parmentier, F.J.W., Brown, R., Bhatt, U.S., Euskirchen, E.S., Romanovsky, V.E., Walsh, J.E., 2019. Key indicators of Arctic climate change: 1971–2017. *Environ. Res. Lett.* 14 (4), 045010.
- Bradley, J.A., Singarayer, J.S., Anesio, A.M., 2014. Microbial community dynamics in the forefield of glaciers. *Proc. R. Soc. B Biol. Sci.* 281 (1795), 20140882.
- Bradley, J.A., Anesio, A., Singarayer, J., Heath, M., Arndt, S., 2015. SHIMMER (1.0): a novel mathematical model for microbial and biogeochemical dynamics in glacier forefield ecosystems. *Geosci. Model Dev.* 8, 3441–3470. <https://doi.org/10.5194/gmd-8-3441-2015>.
- Bradley, J.A., Anesio, A., Arndt, S., 2016a. Bridging the divide: a model-data approach to Polar & Alpine Microbiology. *FEMS Microbiol. Ecol.* 92.
- Bradley, J.A., Arndt, S., Šabacká, M., Benning, L.G., Barker, G.L., Blacker, J.J., Yallop, M. L., Wright, K.E., Bellas, C.M., Telling, J., Tranter, M., 2016b. Microbial dynamics in a High Arctic glacier forefield: a combined field, laboratory, and modelling approach. *Biogeosciences* 13 (19), 5677–5696.
- Brandes, L.J., Den Hollander, H., 1996. SimpleBox 2.0: a nested multimedia fate model for evaluating the environmental fate of chemicals.
- Bring, A., Fedorova, I., Dibike, Y., Hinzman, L., Mård, J., Mernild, S.H., Prowse, T., Semenova, O., Stuefer, S.L., Woo, M.-K., 2016. Arctic terrestrial hydrology: a synthesis of processes, regional effects, and research challenges. *Eur. J. Vasc. Endovasc. Surg.* 121, 621–649. <https://doi.org/10.1002/2015JG003131>.
- Brooks, P.D., Schmidt, S.K., Williams, M.W., 1997. Winter production of CO₂ and N₂O from alpine tundra: environmental controls and relationship to inter-system C and N fluxes. *Oecologia* 110 (3), 403–413.
- Brown, R., Barrette, C., Brown, L., Chaumont, D., Grenier, P., Howell, S., Sharp, M., 2018. Climate variability, trends and projected change. In: *From Science to Policy in the Eastern Canadian Arctic: An Integrated Regional Impact Study (IRIS) of Climate Change and Modernization*. ArcticNet, Quebec City, 560 pp, p.55.
- Bruhwiller, L., Parmentier, F.J.W., Crill, P., Leonard, M., Palmer, P.L., 2021. The Arctic carbon cycle and its response to changing climate. *Curr. Clim. Chang. Rep.* 7, 14–34.
- Bruin, J., Bense, V., Ploeg, M., 2023. Soil thermal properties during freeze–thaw dynamics as function of variable organic carbon and grain size distribution. *Vadose Zone J.* 22.
- Buckeridge, K.M., Grogan, P., 2008. Deepened snow alters soil microbial nutrient limitations in arctic birch hummock tundra. *Appl. Soil Ecol.* 39 (2), 210–222.
- Buckeridge, K.M., Banerjee, S., Siciliano, S.D., Grogan, P., 2013. The seasonal pattern of soil microbial community structure in Mesic low arctic tundra. *Soil Biol. Biochem.* 65, 338–347.
- Bulygina, O.N., Razuvaev, V.N., Korshunova, N.N., 2009. Changes in snow cover over Northern Eurasia in the last few decades. *Environ. Res. Lett.* 4 (4), 045026.
- Burke, E.J., Ekici, A., Huang, Y., Chadburn, S.E., Huntingford, C., Ciais, P., Krinner, G., 2017. Quantifying uncertainties of permafrost carbon–climate feedbacks. *Biogeosciences* 14 (12), 3051–3066.
- Burns, R.G., DeForest, J.L., Marxsen, J., Sinsabaugh, R.L., Stromberger, M.E., Wallenstein, M.D., Weintraub, M.N., Zoppini, A., 2013. Soil enzymes in a changing environment: current knowledge and future directions. *Soil Biol. Biochem.* 58, 216–234.
- Cannone, N., Augusti, A., Malfasi, F., Pallozzi, E., Calfapietra, C., Brugnoli, E., 2016. The interaction of biotic and abiotic factors at multiple spatial scales affects the variability of CO₂ fluxes in polar environments. *Polar Biol.* 39, 1581–1596.
- Cannone, N., Ponti, S., Christiansen, H.H., Christensen, T.R., Pirk, N., Guglielmin, M., 2019. Effects of active layer seasonal dynamics and plant phenology on CO₂ land-atmosphere fluxes at polygonal tundra in the High Arctic, Svalbard. *Catena* 174, 142–153.
- Cao, M., Dent, J.B., Heal, O.W., 1995. Modeling methane emissions from rice paddies. *Glob. Biogeochem. Cycles* 9 (2), 183–195.
- Chadburn, S., Burke, E., Essery, R., Boike, J., Langer, M., Heikenfeld, M., Cox, P., Friedlingstein, P., 2015. An improved representation of physical permafrost dynamics in the JULES land-surface model. *Geosci. Model Dev.* 8 (5), 1493–1508.
- Chadburn, S.E., Krinner, G., Porada, P., Bartsch, A., Beer, C., Beletti Marchesini, L., Boike, J., Ekici, A., Elberling, B., Friborg, T., Hugelius, G., 2017. Carbon stocks and fluxes in the high latitudes: using site-level data to evaluate Earth system models. *Biogeosciences* 14 (22), 5143–5169.
- Chandel, A.K., Jiang, L., Luo, Y., 2023. Microbial models for simulating soil carbon dynamics: a review. *J. Geophys. Res. Biogeosci.* 128 e2023JG007436.
- Chapin III, F.S., Sturm, M., Serreze, M.C., McFadden, J.P., Key, J.R., Lloyd, A.H., McGuire, A.D., Rupp, T.S., Lynch, A.H., Schimel, J.P., Beringer, J., 2005. Role of land-surface changes in Arctic summer warming. *Science* 310 (5748), 657–660.
- Christensen, T.R., Prentice, I.C., Kaplan, J., Haxeltine, A., Sitch, S., 1996. Methane flux from northern wetlands and tundra. *Tellus B* 48 (5), 652–661.
- Chuvpilo, E., Bukhanov, B., Grebenkin, S., Tumsokoy, V., Shakhova, N., Dudarev, O., Semiletov, I., Spasennykh, M., 2021. Thermal properties of sediments in the east Siberian Arctic seas: a case study in the Buor-Khaya Bay. *Mar. Pet. Geol.* 123, 104672.
- Clein, J.S., Schimel, J.P., 1995. Microbial activity of tundra and taiga soils at sub-zero temperatures. *Soil Biol. Biochem.* 27 (9), 1231–1234.
- Coleman, K., Jenkinson, D.S., 1996. RothC-26.3: A Model for the turnover of carbon in soil. In: *Evaluation of Soil Organic Matter Models: Using Existing Long-Term Datasets*. Springer, Berlin Heidelberg, pp. 237–246.
- Commane, R., Lindaas, J., Benmergui, J., Luus, K.A., Chang, R.Y.W., Daube, B.C., Euskirchen, E.S., Henderson, J.M., Karion, A., Miller, J.B., Miller, S.M., 2017. Carbon dioxide sources from Alaska driven by increasing early winter respiration from Arctic tundra. *Proc. Natl. Acad. Sci.* 114 (21), 5361–5366.
- Comyn-Platt, E., et al., 2018. Carbon budgets for 1.5 and 2 °C targets lowered by natural wetland and permafrost feedbacks. *Nat. Geosci.* 11, 568–573.

- Conant, R.T., Ryan, M.G., Ågren, G.I., Birge, H.E., Davidson, E.A., Eliasson, P.E., Evans, S.E., Frey, S.D., Giardina, C.P., Hopkins, F.M., Hyvönen, R., 2011. Temperature and soil organic matter decomposition rates—synthesis of current knowledge and a way forward. *Glob. Chang. Biol.* 17 (11), 3392–3404.
- Cotrufo, M.F., Wallenstein, M.D., Boot, C.M., Deneff, K., Paul, E., 2013. The Microbial Efficiency-Matrix Stabilization (MEMS) framework integrates plant litter decomposition with soil organic matter stabilization: do labile plant inputs form stable soil organic matter? *Glob. Chang. Biol.* 19 (4), 988–995.
- Dankers, R., Burke, E.J., Price, J., 2011. Simulation of permafrost and seasonal thaw depth in the JULES land surface scheme. *Cryosphere* 5 (3), 773–790.
- Datta, R., Kelkar, A., Baraniya, D., Molaei, A., Moullick, A., Meena, R.S., Formanek, P., 2017. Enzymatic degradation of lignin in soil: a review. *Sustainability* 9 (7), 1163.
- Davidson, E.A., Janssens, I.A., 2006. Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature* 440 (7081), 165–173.
- Davidson, E.A., Samanta, S., Caramori, S.S., Savage, K., 2012. The Dual Arrhenius and Michaelis-Menten kinetics model for decomposition of soil organic matter at hourly to seasonal time scales. *Glob. Chang. Biol.* 18 (1), 371–384.
- De La Fuente, M., Vaunat, J., Marín-Moreno, H., 2020. A densification mechanism to model the mechanical effect of methane hydrates in sandy sediments. *Int. J. Numer. Anal. Methods Geomech.* 44, 782–802.
- De La Fuente, M., Vaunat, J., Marín-Moreno, H., 2021. Modelling methane Hydrate Saturation in Pores: Capillary Inhibition Effects. *Energies* 15, 5627.
- De La Fuente, M., Arndt, S., Marín-Moreno, H., Minshull, T.A., 2022. Assessing the Benthic Response to Climate-Driven methane Hydrate Destabilisation: State of the Art and Future Modelling Perspectives. *Energies* 15, 3307.
- De Michele, C., Avanzi, F., Ghezzi, A., Jommi, C., 2013. Investigating the dynamics of bulk snow density in dry and wet conditions using a one-dimensional model. *Cryosphere* 7 (2), 433–444.
- Del Grosso, S., Ojima, D., Parton, W., Mosier, A., Peterson, G., Schimel, D., 2002. Simulated effects of dryland cropping intensification on soil organic matter and greenhouse gas exchanges using the DAYCENT ecosystem model. *Environ. Pollut.* 116, S75–S83.
- D'Imperio, L., Nielsen, C.S., Westergaard-Nielsen, A., Michelsen, A., Elberling, B., 2017. Methane oxidation in contrasting soil types: responses to experimental warming with implication for landscape-integrated CH₄ budget. *Glob. Chang. Biol.* 23 (2), 966–976.
- Dong, Y., McCartney, J.S., Lu, N., 2015. Critical Review of thermal Conductivity Models for Unsaturated Soils. *Geotech. Geol. Eng.* 33, 207–221.
- Drake, J.E., Darby, B.A., Giasson, M.A., Kramer, M.A., Phillips, R.P., Finzi, A.C., 2013. Stoichiometry constrains microbial response to root exudation—insights from a model and a field experiment in a temperate forest. *Biogeosciences* 10 (2), 821–838.
- Dunfield, P., Dumont, R., Moore, T.R., 1993. Methane production and consumption in temperate and subarctic peat soils: response to temperature and pH. *Soil Biol. Biochem.* 25 (3), 321–326.
- Ekici, A., Chadburn, S., Chaudhry, N., Hajdu, L.H., Marmy, A., Peng, S., Boike, J., Burke, E., Friend, A.D., Hauck, C., Krinner, G., 2015. Site-level model intercomparison of high latitude and high altitude soil thermal dynamics in tundra and barren landscapes. *Cryosphere* 9 (4), 1343–1361.
- Elberling, B., Brandt, K.K., 2003. Uncoupling of microbial CO₂ production and release in frozen soil and its implications for field studies of arctic C cycling. *Soil Biol. Biochem.* 35 (2), 263–272.
- Elmendorf, S.C., Henry, G.H., Hollister, R.D., Björk, R.G., Boulanger-Lapointe, N., Cooper, E.J., Cornelissen, J.H., Day, T.A., Dorrepaal, E., Elumeeva, T.G., Gill, M., 2012. Plot-scale evidence of tundra vegetation change and links to recent summer warming. *Nat. Clim. Chang.* 2 (6), 453–457.
- Euskirchen, E.S., Bret-Harte, M.S., Shaver, G.R., Edgar, C.W., Romanovsky, V.E., 2017. Long-term release of carbon dioxide from arctic tundra ecosystems in Alaska. *Ecosystems* 20 (5), 960–974.
- Fahnestock, J.T., Jones, M.H., Welker, J.M., 1999. Wintertime CO₂ efflux from arctic soils: Implications for annual carbon budgets. *Global Biogeochem. Cy.* 13, 775–779.
- Fan, Z., David McGuire, A., Turetsky, M.R., Harden, J.W., Michael Waddington, J., Kane, E.S., 2013. The response of soil organic carbon of a rich fen peatland in interior Alaska to projected climate change. *Glob. Chang. Biol.* 19 (2), 604–620.
- Farquharson, L.M., Romanovsky, V.E., Kholodov, A., Nicolsky, D., 2022. Sub-aerial talik formation observed across the discontinuous permafrost zone of Alaska. *Nat. Geosci.* 15, 475–481.
- Fiedler, S., Vepraskas, M.J., Richardson, J.L., 2007. Soil redox potential: importance, field measurements, and observations. In: Sparks, D.L. (Ed.), *Advances in Agronomy*, 94. Academic Press, pp. 1–54. ISSN 0065-2113, ISBN 9780123741073. [https://doi.org/10.1016/S0065-2113\(06\)94001-2](https://doi.org/10.1016/S0065-2113(06)94001-2).
- Finestone, J., Templer, P.H., Bhatnagar, J.M., 2022. Soil fungi exposed to warming temperatures and shrinking snowpack in a northern forest have lower capacity for growth and nutrient cycling. *Front. Global Change* 5. <https://doi.org/10.3389/fgc.2022.800335>.
- Fisher, J.B., Badgley, G., Blyth, E., 2012. Global nutrient limitation in terrestrial vegetation. *Glob. Biogeochem. Cycles* 26 (3).
- Fisher, J.B., Sikka, M., Oechel, W.C., Huntzinger, D.N., Melton, J.R., Koven, C.D., Ahlström, A., Arain, M.A., Baker, I., Chen, J.M., Ciais, P., 2014. Carbon cycle uncertainty in the Alaskan Arctic. *Biogeosciences* 11 (15), 4271–4288.
- Fisher, J.B., Hayes, D.J., Schwalm, C.R., Huntzinger, D.N., Stofferahn, E., Schaefer, K., Luo, Y., Wullschlegel, S.D., Goetz, S., Miller, C.E., Griffith, P., 2018. Missing pieces to modeling the Arctic-Boreal puzzle. *Environ. Res. Lett.* 13 (2), 020202.
- Flessa, H., Rodionov, A., Guggenberger, G., Fuchs, H., Magdon, P., Shibistova, O., Zrazhevskaya, G., Mikheyeva, N., Kasansky, O.A., Blodau, C., 2008. Landscape controls of CH₄ fluxes in a catchment of the forest tundra ecotone in northern Siberia. *Glob. Chang. Biol.* 14 (9), 2040–2056.
- Fontaine, S., Barot, S., 2005. Size and functional diversity of microbe populations control plant persistence and long-term soil carbon accumulation. *Ecol. Lett.* 8 (10), 1075–1087.
- Frey, S.D., Drijber, R., Smith, H., Melillo, J., 2008. Microbial biomass, functional capacity, and community structure after 12 years of soil warming. *Soil Biol. Biochem.* 40 (11), 2904–2907.
- Fu, Z., Wu, Q., Zhang, W., He, H., Wang, L., 2022. Water migration and segregated ice formation in frozen ground: current advances and future perspectives. *Front. Earth Sci.* 10, 826961.
- Geml, J., Morgado, L.N., Semenova-Nelsen, T.A., 2021. Tundra Type Drives Distinct Trajectories of Functional and Taxonomic Composition of Arctic Fungal Communities in Response to climate Change—results from Long-Term Experimental Summer Warming and increased Snow Depth. *Front. Microbiol.* 12, 628746.
- Genet, H., McGuire, A.D., Barrett, K., Breen, A., Euskirchen, E.S., Johnstone, J.F., Kasischke, E.S., Melvin, A.M., Bennett, A., Mack, M.C., Rupp, T.S., 2013. Modeling the effects of fire severity and climate warming on active layer thickness and soil carbon storage of black spruce forests across the landscape in interior Alaska. *Environ. Res. Lett.* 8 (4), 045016.
- Goulden, M.L., Wofsy, S.C., Harden, J.W., Trumbore, S.E., Crill, P.M., Gower, S.T., Fries, T., Daube, B.C., Fan, S.M., Sutton, D.J., Bazzaz, A., 1998. Sensitivity of boreal forest carbon balance to soil thaw. *Science* 279 (5348), 214–217.
- Graham, R.M., Cohen, L., Petty, A.A., Boisvert, L.N., Rinke, C.A., Hudson, S.R., Nicolaus, M., Granskog, M.A., 2017. Increasing frequency and duration of Arctic winter warming events. *Geophys. Res. Lett.* 44 (13), 6974–6983.
- Grant, R.F., 1998. Simulation of methanogenesis in the mathematical model ecosys. *Soil Biol. Biochem.* 30 (7), 883–896.
- Grant, R.F., Juma, N.G., Robertson, J.A., Izaurre, R.C., McGill, W.B., 2001. Long-term changes in soil carbon under different fertilizer, manure, and rotation: Testing the mathematical model ecosys with data from the Breton plots. *Soil Sci. Soc. Am. J.* 65 (1), 205–214.
- Guo, X., Gao, Q., Yuan, M., Wang, G., Zhou, X., Feng, J., Shi, Z., Hale, L., Wu, L., Zhou, A., Tian, R., 2020. Gene-informed decomposition model predicts lower soil carbon loss due to persistent microbial adaptation to warming. *Nat. Commun.* 11 (1), 1–12.
- Harris, R.F., 1981. Effect of water potential on microbial growth and activity. *Water Potent. Relat. Soil Microbiol.* 9, 23–95.
- Hartl, L., Stuefer, M., Saito, T., Okura, Y., 2020. History and Data Records of the Automatic Weather Station on Denali Pass (5715 m), 1990–2007. *J. Appl. Meteorol. Climatol.* 59 (12), 2113–2127.
- He, Y., Zhuang, Q., Harden, J.W., McGuire, A.D., Fan, Z., Liu, Y., Wickland, K.P., 2014. The implications of microbial and substrate limitation for the fates of carbon in different organic soil horizon types of boreal forest ecosystems: a mechanistically based model analysis. *Biogeosciences* 11 (16), 4477–4491.
- He, Y., Yang, J., Zhuang, Q., Harden, J.W., McGuire, A.D., Liu, Y., Wang, G., Gu, L., 2015. Incorporating microbial dormancy dynamics into soil decomposition models to improve quantification of soil carbon dynamics of northern temperate forests. *J. Geophys. Res. Biogeosci.* 120 (12), 2596–2611.
- He, L., Lai, C.T., Mayes, M.A., Murayama, S., Xu, X., 2021. Microbial seasonality promotes soil respiratory carbon emission in natural ecosystems: a modeling study. *Glob. Chang. Biol.* 27 (13), 3035–3051.
- Heinze, T., 2021. A multi-phase heat transfer model for water infiltration into frozen soil. *Water Resour. Res.* 57, e2021WR030067.
- Hellweger, F.L., Bucci, V., 2009. A bunch of tiny individuals—individual-based modeling for microbes. *Ecol. Model.* 220, 8–22.
- Herbert, R.G., 1959. Maintenance in Nullity Actions—The BC Supreme Court Rules. *U. Brit. Colum. L. Rev.* 1, 581.
- Herndon, E., Kinsman-Costello, L., Godsey, S., 2020. Biogeochemical cycling of redox-sensitive elements in permafrost-affected ecosystems. In: Dontsova, K., Balogh-Brunstad, Z., Le Roux, G. (Eds.), *Biogeochemical Cycles*. <https://doi.org/10.1002/9781119413332.ch12>.
- Hinkel, K.M., Paetzold, F., Nelson, F.E., Bockheim, J.G., 2001. Patterns of soil temperature and moisture in the active layer and upper permafrost at Barrow, Alaska: 1993–1999. *Glob. Planet. Change* 29 (3–4), 293–309.
- Hobbs, J.K., Jiao, W., Easter, A.D., Parker, E.J., Schipper, L.A., Arcus, V.L., 2013. Change in heat capacity for enzyme catalysis determines temperature dependence of enzyme catalyzed rates. *ACS Chem. Biol.* 8 (11), 2388–2393.
- Hodson, A.J., Nowak, A., Redeker, K.R., Holmlund, E.S., Christiansen, H.H., Turchyn, A.V., 2019. Seasonal Dynamics of Methane and Carbon Dioxide Evasion from an Open System Pingo: Lagoon Pingo. *Frontiers in Earth Science*, Svalbard, p. 30.
- Holl, D., Wille, C., Sachs, T., Schreiber, P., Runkle, B.R., Beckebanze, L., Langer, M., Boike, J., Pfeiffer, E.M., Fedorova, I., Bolshinov, D.Y., 2019. A long-term (2002 to 2017) record of closed-path and open-path eddy covariance CO₂ net ecosystem exchange fluxes from the Siberian Arctic. *Earth Syst. Sci. Data* 11 (1), 221–240.
- Hopple, A.M., Wilson, R.M., Kolton, M., Zalman, C.A., Chanton, J.P., Kostka, J., Hanson, P.J., Keller, J.K., Bridgman, S.D., 2020. Massive peatland carbon banks vulnerable to rising temperatures. *Nat. Commun.* 11 (1), 1–7.
- Huang, X., Shi, Z., Zhu, H., et al., 2016. Soil moisture dynamics within soil profiles and associated environmental controls. *Catena* 136, 189–196.
- Huang, Y., Guenet, B., Wang, Y.L., Ciais, P., 2021. Global simulation and evaluation of soil organic matter and microbial carbon and nitrogen stocks using the microbial decomposition model ORCHIMIC v2.0. *Glob. Biogeochem. Cycles* 35 (5), e2020GB006836.
- Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J.W., Schuur, E.A.G., Ping, C.L., Schirmer, L., Grosse, G., Michaelson, G.J., Koven, C.D., O'Donnell, J.A., 2014. Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps. *Biogeosciences* 11 (23), 6573–6593.

- Ingwersen, J., Poll, C., Streck, T., Kandeler, E., 2008. Micro-scale modelling of carbon turnover driven by microbial succession at a biogeochemical interface. *Soil Biol. Biochem.* 40 (4), 864–878.
- Ito, A., Li, T., Qin, Z., Melton, J.R., Tian, H., Kleinen, T., Zhang, W., Zhang, Z., Joos, F., Ciais, P., Hopcroft, P.O., 2023. Cold-season methane fluxes simulated by GCP-CH4 Models. *Geophys. Res. Lett.* 50 (14) p.e2023GL103037.
- Jenkinson, D.S., 1990. The turnover of organic carbon and nitrogen in soil. *Philos. Trans. R. Soc. Lond. Ser. B Biol. Sci.* 329 (1255), 361–368.
- Jenkinson, D.S., Coleman, K., 2008. The turnover of organic carbon in subsoils. Part 2. Modelling carbon turnover. *Eur. J. Soil Sci.* 59 (2), 400–413.
- Jenkinson, D.S., Hart, P.B.S., Rayner, J.H., Parry, L.C., 1987. Modelling the Turnover of Organic Matter in Long-Term Experiments at Rothamsted.
- Jentzsch, K., Schulz, A., Pirk, N., Foken, T., Crewell, S., Boike, J., 2021. High Levels of CO₂ Exchange during Synoptic-Scale events Introduce large uncertainty into the Arctic Carbon Budget. *Geophys. Res. Lett.* 48 (9) e2020GL092256.
- Jiang, R., Li, T., Liu, D., Fu, Q., Hou, R., Li, Q., Cui, S., Li, M., 2021. Soil infiltration characteristics and pore distribution under freezing–thawing conditions. *Cryosphere* 15, 2133–2146.
- Jin, H., Huang, Y., Bense, V.F., Ma, Q., Marchenko, S.S., Shepelev, V.V., Hu, Y., Liang, S., Spektor, V.V., Jin, X., et al., 2022. Permafrost Degradation and its Hydrogeological Impacts. *Water* 14, 372.
- Joabsson, A., Christensen, T.R., 2001. Methane emissions from wetlands and their relationship with vascular plants: an Arctic example. *Glob. Chang. Biol.* 7 (8), 919–932.
- Jobbágy, E.G., Jackson, R.B., 2000. The vertical distribution of soil organic carbon and its relation to climate and vegetation. *Ecol. Appl.* 10 (2), 423–436.
- Juncher Jørgensen, C., Lund Johansen, K.M., Westergaard-Nielsen, A., Elberling, B., 2015. Net regional methane sink in High Arctic soils of Northeast Greenland. *Nat. Geosci.* 8 (1), 20–23.
- Justice, C.O., Vermote, E., Townshend, J.R., Defries, R., Roy, D.P., Hall, D.K., Salomonson, V.V., Privette, J.L., Riggs, G., Strahler, A., Lucht, W., 1998. The Moderate Resolution Imaging Spectroradiometer (MODIS): Land remote sensing for global change research. *IEEE Trans. Geosci. Remote Sens.* 36 (4), 1228–1249.
- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., 1996. The NCEP/NCAR 40-year reanalysis project. *Bull. Am. Meteorol. Soc.* 77 (3), 437–472.
- Keuschnig, C., et al., 2022. Reduced methane emissions in former permafrost soils driven by vegetation and microbial changes following drainage. *Glob. Chang. Biol.* <https://doi.org/10.1111/gcb.16137>.
- Khimenkov, A., Stanilovskaya, J., 2022. Explosive Processes in Permafrost as a result of the Development of Local Gas-Saturated Fluid-Dynamic Geosystems. *Gases* 2, 146–165.
- Klepper, O., den Hollander, H.A., 1999. A comparison of spatially explicit and box models for the fate of chemicals in water, air and soil in Europe. *Ecol. Model.* 116 (2–3), 183–202.
- Koarashi, J., Hockaday, W.C., Masiello, C.A., Trumbore, S.E., 2012. Dynamics of decadal cycling carbon in subsurface soils. *J. Geophys. Res. Biogeosci.* 117 (G3).
- Koponen, H.T., Martikainen, P.J., 2004. Soil water content and freezing temperature affect freeze–thaw related N₂O production in organic soil. *Nutr. Cycl. Agroecosyst.* 69 (3), 213–219.
- Koven, C.D., Riley, W.J., Subin, Z.M., Tang, J.Y., Torn, M.S., Collins, W.D., Bonan, G.B., Lawrence, D.M., Swenson, S.C., 2013. The effect of vertically resolved soil biogeochemistry and alternate soil C and N models on C dynamics of CLM4. *Biogeosciences* 10 (11), 7109–7131.
- Kozłowski, T., 2009. Some factors affecting supercooling and the equilibrium freezing point in soil–water systems. *Cold Reg. Sci. Technol.* 59 (1), 25–33.
- Kuc, T., 2005. A multi-layer box model of carbon dynamics in soil. *Nukleonika* 50 (2), 49–55.
- Kussell, E., Leibler, S., 2005. Phenotypic diversity, population growth, and information in fluctuating environments. *Science* 309 (5743), 2075–2078.
- Kuzyakov, Y., Friedel, J.K., Stahr, K., 2000. Review of mechanisms and quantification of priming effects. *Soil Biol. Biochem.* 32 (11–12), 1485–1498.
- Lafrenière, M.J., Lamoureaux, S.F., 2019. Effects of changing permafrost conditions on hydrological processes and fluvial fluxes. *Earth Sci. Rev.* 191, 212–223. ISSN 0012-8252. <https://doi.org/10.1016/j.earscirev.2019.02.018>.
- Lange, S.F., Allaire, S.E., Cuellar Castillo, M.A., Dutilleul, P., 2016. N₂O and CO₂ dynamics in a pasture soil across the frozen period. *Can. J. Soil Sci.* 97, 497–511.
- Larsbo, M., Holten, R., Stenrod, M., Eklo, O.M., Jarvis, N., 2019. A dual-per-meability approach for modeling soil water flow and heat transport during freezing and thawing. *Vadose Zone J.* 18, 190012.
- Lau, M.C., Stackhouse, B.T., Layton, A.C., Chauhan, A., Vishnivetskaya, T.A., Chourey, K., Ronholm, J., Mykityczuk, N.C.S., Bennett, P.C., Lamarque-Gagnon, G., Burton, N., 2015. An active atmospheric methane sink in high Arctic mineral cryosols. *ISME J.* 9 (8), 1880–1891.
- Lawrence, D.M., Slater, A.G., 2010. The contribution of snow condition trends to future ground climate. *Clim. Dyn.* 34 (7), 969–981.
- Lawrence, D.M., Oleson, K.W., Flanner, M.G., Thornton, P.E., Swenson, S.C., Lawrence, P.J., Zeng, X., Yang, Z.L., Levis, S., Sakaguchi, K., Bonan, G.B., 2011. Parameterization improvements and functional and structural advances in version 4 of the Community Land Model. *J. Adv. Model. Earth Syst.* 3 (1).
- Le Mer, J., Roger, P., 2001. Production, oxidation, emission and consumption of methane by soils: a review. *Eur. J. Soil Biol.* 37 (1), 25–50.
- Lemke, P., Ren, J., Alley, R.B., Allison, I., Carrasco, J., Flato, G., Fujii, Y., Kaser, G., Mote, P., Thomas, R.H., Zhang, T., 2007. Observations: Changes in Snow, Ice and Frozen Ground.
- Lennon, J.T., Jones, S.E., 2011. Microbial seed banks: the ecological and evolutionary implications of dormancy. *Nat. Rev. Microbiol.* 9 (2), 119–130.
- Li, C., Frolking, S., Frolking, T.A., 1992. A model of nitrous oxide evolution from soil driven by rainfall events: 2. Model applications. *J. Geophys. Res.-Atmos.* 97 (D9), 9777–9783.
- Li, C., Frolking, S., Harriss, R., 1994. Modeling carbon biogeochemistry in agricultural soils. *Glob. Biogeochem. Cycles* 8 (3), 237–254.
- Lipson, D.A., Schmidt, S.K., 2004. Seasonal changes in an alpine soil bacterial community in the Colorado Rocky Mountains. *Appl. Environ. Microbiol.* 70, 2867–2879.
- Lipson, D.A., Schadt, C.W., Schmidt, S.K., 2002. Changes in microbial community structure and function in dry meadow tundra following spring snow melt. *Microb. Ecol.* 43, 307–314.
- Lipson, D.A., Monson, R.K., Schmidt, S.K., Weintraub, M.N., 2009. The trade-off between growth rate and yield in microbial communities and the consequences for under-snow soil respiration in a high elevation coniferous forest. *Biogeochemistry* 95 (1), 23–35.
- Liptzin, D., Williams, M.W., Helmig, D., et al., 2009. Process-level controls on CO₂ fluxes from a seasonally snow-covered subalpine meadow soil, Niwot Ridge, Colorado. *Biogeochemistry* 95, 151–166.
- Liptzin, D., Helmig, D., et al., 2015. Winter gas exchange between the atmosphere and snow-covered soils on Niwot Ridge, Colorado, USA. *Plant Ecol. Diversity* 8 (5–6), 677–688.
- Liu, S., Zhuang, Q., He, Y., Noormets, A., Chen, J., Gu, L., 2016. Evaluating atmospheric CO₂ effects on gross primary productivity and net ecosystem exchanges of terrestrial ecosystems in the conterminous United States using the AmeriFlux data and an artificial neural network approach. *Agric. For. Meteorol.* 220, 38–49.
- Lloyd, C.R., 2001. The measurement and modelling of the carbon dioxide exchange at a high Arctic site in Svalbard. *Glob. Chang. Biol.* 7 (4), 405–426.
- Locey, K.J., 2010. Synthesizing traditional biogeography with microbial ecology: the importance of dormancy. *J. Biogeogr.* 37 (10), 1835–1841.
- López-Blanco, E., Lund, M., Williams, M., Tamstorf, M.P., Westergaard-Nielsen, A., Exbrayat, J.F., Hansen, B.U., Christensen, T.R., 2017. Exchange of CO₂ in Arctic tundra: impacts of meteorological variations and biological disturbance. *Biogeosciences* 14 (19), 4467–4483.
- Lüers, J., Westermann, S., Piel, K., Boike, J., 2014. Annual CO₂ budget and seasonal CO₂ exchange signals at a high Arctic permafrost site on Spitsbergen, Svalbard archipelago. *Biogeosciences* 11 (22), 6307–6322.
- Lund, M., 2018. Uncovering the unknown-climate interactions in a changing arctic tundra. *Environ. Res. Lett.* 13 (061001), 10–1088.
- Lund, M., Lafleur, P.M., Roulet, N.T., Lindroth, A., Christensen, T.R., Aurela, M., Chojnicki, B.H., Flanagan, L.B., Humphreys, E.R., Laurila, T., Oechel, W.C., 2010. Variability in exchange of CO₂ across 12 northern peatland and tundra sites. *Glob. Chang. Biol.* 16 (9), 2436–2448.
- Lund, M., Falk, J.M., Friborg, T., Mbufong, H.N., Sigsgaard, C., Soegaard, H., Tamstorf, M.P., 2012. Trends in CO₂ exchange in a high Arctic tundra heath, 2000–2010. *J. Geophys. Res. Biogeosci.* 117 (G2).
- Lund, M., Hansen, B.U., Pedersen, S.H., Stieglar, C., Tamstorf, M.P., 2014. Characteristics of summer-time energy exchange in a high Arctic tundra heath 2000–2010. *Tellus Ser. B Chem. Phys. Meteorol.* 66 (1), 21631.
- Lyu, Z., Zhuang, Q., 2018. Quantifying the effects of snowpack on soil thermal and carbon dynamics of the Arctic terrestrial ecosystems. *J. Geophys. Res. Biogeosci.* 123 (4), 1197–1212.
- Lyu, Z., Genet, H., He, Y., Zhuang, Q., McGuire, A.D., Bennett, A., Breen, A., Clein, J., Euskirchen, E.S., Johnson, K., Kurkowski, T., 2018. The role of environmental driving factors in historical and projected carbon dynamics of wetland ecosystems in Alaska. *Ecol. Appl.* 28 (6), 1377–1395.
- Magnani, M., Baneschi, I., Giamberini, M., Raco, B., Provenzale, A., 2022. Microscale drivers of summer CO₂ fluxes in the Svalbard High Arctic tundra. *Sci. Rep.* 12 (1), 1–14.
- Männistö, M.K., Häggblom, M.M., 2006. Characterization of psychrotolerant heterotrophic bacteria from Finnish Lapland. *Syst. Appl. Microbiol.* 29 (3), 229–243.
- Manzoni, S., Porporato, A., 2007. A theoretical analysis of nonlinearities and feedbacks in soil carbon and nitrogen cycles. *Soil Biol. Biochem.* 39 (7), 1542–1556.
- Manzoni, S., Porporato, A., 2009. Soil carbon and nitrogen mineralization: Theory and models across scales. *Soil Biol. Biochem.* 41 (7), 1355–1379.
- Manzoni, S., Schimel, J.P., Porporato, A., 2012. Responses of soil microbial communities to water stress: results from a meta-analysis. *Ecology* 93 (4), 930–938.
- Manzoni, S., Schaeffer, S.M., Katul, G., Porporato, A., Schimel, J.P., 2014. A theoretical analysis of microbial eco-physiological and diffusion limitations to carbon cycling in drying soils. *Soil Biol. Biochem.* 73, 69–83.
- Mazur, P., 1980. Limits to life at low temperatures and at reduced water contents and water activities. In: *Limits of Life: Proceedings of the fourth College Park Colloquium on Chemical Evolution*. University of Maryland, College Park, Maryland, USA. October 18th to 20th, 1978 (pp. 1–23). Springer Netherlands.
- McGuire, A.D., Christensen, T.R., Hayes, D., Heroult, A., Euskirchen, E., Kimball, J.S., Koven, C., Lafleur, P., Miller, P.A., Oechel, W., Peylin, P., 2012. An assessment of the carbon balance of Arctic tundra: comparisons among observations, process models, and atmospheric inversions. *Biogeosciences* 9 (8), 3185–3204.
- McGuire, A.D., Koven, C., Lawrence, D.M., Clein, J.S., Xia, J., Beer, C., Burke, E., Chen, G., Chen, X., Delire, C., Jafarov, E., 2016. Variability in the sensitivity among model simulations of permafrost and carbon dynamics in the permafrost region between 1960 and 2009. *Glob. Biogeochem. Cycles* 30 (7), 1015–1037.
- McGuire, A.D., Lawrence, D.M., Koven, C., Clein, J.S., Burke, E., Chen, G., Jafarov, E., MacDougall, A.H., Marchenko, S., Nicolosky, D., Peng, S., Rinke, A., Ciais, P., Gouttevin, I., Hayes, D.J., Ji, D., Krinner, G., Moore, J.C., Romanovsky, V.E.,

- Schädel, C., Schaefer, K., Schuur, E.A.G., Zhuang, Q., 2018a. The dependence of the evolution of carbon dynamics in the northern permafrost region on the trajectory of climate change. *Proc. Natl. Acad. Sci.* <https://doi.org/10.1073/pnas.1719903115>, 6 pages.
- McGuire, A.D., Genet, H., Lyu, Z., Pastick, N., Stackpole, S., Birdsey, R., D'Amore, D., He, Y., Rupp, T.S., Striegl, R., Wylie, B.K., 2018b. Assessing historical and projected carbon balance of Alaska: a synthesis of results and policy/management implications. *Ecol. Appl.* 28 (6), 1396–1412.
- Meentemeyer, V., Box, E.O., Thompson, R., 1982. World patterns and amounts of terrestrial plant litter production. *BioScience* 32 (2), 125–128.
- Melton, J.R., Wania, R., Hodson, E.L., Poulter, B., Ringeval, B., Spahni, R., Bohn, T., Avis, C.A., Beerling, D.J., Chen, G., Eliseev, A.V., 2013. Present state of global wetland extent and wetland methane modelling: conclusions from a model inter-comparison project (WETCHIMP). *Biogeosciences* 10 (2), 753–788.
- Meredith, M., Sommerkorn, M., Cassotta, S., Derksen, C., Ekaykin, A., Hollowed, A., Kofinas, G., Maelkne, A., Melbourne-Thomas, J., Muelbert, M.M.C., Ottersen, G., 2019. Polar regions. In: Chapter 3, IPCC Special Report on the Ocean and Cryosphere in a Changing Climate.
- Michaelis, L., Menten, M.L., 1913. The kinetics of the inversion effect. *Biochem. Z.* 49, 333–369.
- Monod, J., 1949. The growth of bacterial cultures. *Ann. Rev. Microbiol.* 3 (1), 371–394.
- Monson, R.K., Lipson, D.A., Burns, S., et al., 2006. Winter soil respiration controlled by climate and microbial community composition. *Nature* 439, 711–714.
- Moorhead, D.L., Sinsabaugh, R.L., 2006. A theoretical model of litter decay and microbial interaction. *Ecol. Monogr.* 76 (2), 151–174.
- Mu, C., Zhang, T., Zhang, X., Cao, B., Peng, X., 2016. Sensitivity of soil organic matter decomposition to temperature at different depths in permafrost regions on the northern Qinghai-Tibet plateau. *Eur. J. Soil Sci.* 67 (6), 773–781.
- Myers, R.J.K., Weier, K.L., Campbell, C.A., 1982. Quantitative relationship between net nitrogen mineralization and moisture content of soils. *Can. J. Soil Sci.* 62 (1), 111–124.
- Nakatsubo, T., Bekku, Y.S., Uchida, M., Muraoka, H., Kume, A., Ohtsuka, T., Masuzawa, T., Kanda, H., Koizumi, H., 2005. Ecosystem development and carbon cycle on a glacier foreland in the High Arctic, Ny-Ålesund, Svalbard. *J. Plant Res.* 118, 173–179.
- Natali, S.M., Watts, J.D., Rogers, B.M., Potter, S., Ludwig, S.M., Selbmann, A.K., Sullivan, P.F., Abbott, B.W., Arndt, K.A., Birch, L., Björkman, M.P., 2019. Large loss of CO₂ in winter observed across the northern permafrost region. *Nat. Clim. Chang.* 9 (11), 852–857.
- Naylor, D., Sadler, N., Bhattacharjee, A., Graham, E.B., Anderton, C.R., McClure, R., Lipton, M., Hofmocker, K.S., Jansson, J.K., 2020. Soil microbiomes under climate change and implications for carbon cycling. *Annu. Rev. Environ. Resour.* 45 (1), 29–59.
- Newsham, K.K., Danielsen, B.K., Biersma, E.M., Elberling, B., Hillyard, G., Kumari, P., Yamamoto, N., 2022. Rapid Response to Experimental Warming of a Microbial Community Inhabiting High Arctic Patterned Ground Soil. *Biology* 11 (12), 1819.
- Nicolovsky, D.J., Romanovsky, V.E., Panda, S.K., Marchenko, S.S., Muskett, R.R., 2017. Applicability of the ecosystem type approach to model permafrost dynamics across the Alaska North Slope. *J. Geophys. Res. Earth* 122 (1), 50–75.
- Nielsen, C.B., Groffman, P.M., Hamburg, S.P., Driscoll, C.T., Fahey, T.J., Hardy, J.P., 2001. Freezing effects on carbon and nitrogen cycling in northern hardwood forest soils. *Soil Sci. Soc. Am. J.* 65 (6), 1723–1730.
- Nitzbon, J., Langer, M., Westermann, S., Martin, L., Aas, K.S., Boike, J., 2019. Pathways of ice-wedge degradation in polygonal tundra under different hydrological conditions. *Cryosphere* 13 (4), 1089–1123.
- Oechel, W.C., Hastings, S.J., Vourlitis, G., Jenkins, M., Riechers, G., Grulke, N., 1993. Recent change of Arctic tundra ecosystems from a net carbon dioxide sink to a source. *Nature* 361 (6412), 520–523.
- Oechel, W.C., Laskowski, C.A., Burba, G., Gioli, B., Kalhori, A.A., 2014. Annual patterns and budget of CO₂ flux in an Arctic tussock tundra ecosystem. *J. Geophys. Res. Biogeosci.* 119 (3), 323–339.
- Oh, Y., Zhuang, Q., Liu, L., Welp, L.R., Lau, M.C., Onstott, T.C., Medvigy, D., Bruhwiler, L., Dlugokencky, E.J., Hugelius, G., D'Imperio, G., 2020. Reduced net methane emissions due to microbial methane oxidation in a warmer Arctic. *Nat. Clim. Chang.* 10 (4), 317–321.
- Osterkamp, T.E., 2007. Characteristics of the recent warming of permafrost in Alaska. *J. Geophys. Res. Earth* 112 (F2).
- Outcalt, S.I., Nelson, F.E., Hinkel, K.M., 1990. The zero-curtain effect: Heat and mass transfer across an isothermal region in freezing soil. *Water Resour. Res.* 26 (7), 1509–1516.
- Panikov, N.S., 1995. Microbial Growth Kinetics.
- Pansu, M., Bottner, P., Sarmiento, L., Metselaar, K., 2004. Comparison of five soil organic matter decomposition models using data from a 14C and 15N labeling field experiment. *Glob. Biogeochem. Cycles* 18 (4).
- Pansu, M., Sarmiento, L., Rujano, M.A., Ablan, M., Acevedo, D., Bottner, P., 2010. Modeling organic transformations by microorganisms of soils in six contrasting ecosystems: Validation of the MOMOS model. *Glob. Biogeochem. Cycles* 24 (1).
- Parazoo, N.C., Koven, C.D., Lawrence, D.M., Romanovsky, V., Miller, C.E., 2018. Detecting the permafrost carbon feedback: talik formation and increased cold-season respiration as precursors to sink-to-source transitions. *Cryosphere* 12, 123–144.
- Parmentier, F.J.W., Van Der Molen, M.K., Van Huisteden, J., Karsanaev, S.A., Kononov, A.V., Suzdalov, D.A., Maximov, T.C., Dolman, A.J., 2011. Longer growing seasons do not increase net carbon uptake in the northeastern Siberian tundra. *J. Geophys. Res. Biogeosci.* 116 (G4).
- Parton, W.J., Schimel, D.S., Cole, C.V., Ojima, D.S., 1987. Analysis of factors controlling soil organic matter levels in Great Plains grasslands. *Soil Sci. Soc. Am. J.* 51 (5), 1173–1179.
- Parton, W.J., Stewart, J.W., Cole, C.V., 1988. Dynamics of C, N, P and S in grassland soils: a model. *Biogeochemistry* 5 (1), 109–131.
- Patrick, W.H., Gambrell, R.P., Faulkner, S.P., 1996. Redox measurements of soils. In: Sparks, D.L., Page, A.L., Helmke, P.A., Loeppert, R.H., Soltanpour, P.N., Tabatabai, M.A., Johnston, C.T., Sumner, M.E. (Eds.), *Methods of Soil Analysis*. <https://doi.org/10.2136/sssabookser5.3.c42>.
- Pavlova, K.K., 1970. Phase composition of water and thermo-physical characteristics of frozen peat in the study of infiltration. *Soviet Hydrol. Selected Pap.* 4, 361–378.
- Pedron, S.A., Welker, J.M., Euskirchen, E.S., Klein, E.S., Walker, J.C., Xu, X., Czimczik, C. I., 2022. Closing the Winter Gap—Year-Round Measurements of Soil CO₂ Emission sources in Arctic Tundra. *Geophys. Res. Lett.* 49 (6) p.e2021GL097347.
- Petersen, B.M., Bernsten, J., Hansen, S., Jensen, L.S., 2005. CN-SIM—a model for the turnover of soil organic matter. I. Long-term carbon and radiocarbon development. *Soil Biol. Biochem.* 37 (2), 359–374.
- Pianka, E.R., 1970. On r-and K-selection. *Am. Nat.* 104 (940), 592–597.
- Piggot, P.J., Hilbert, D.W., 2004. Sporulation of *Bacillus subtilis*. *Curr. Opin. Microbiol.* 7 (6), 579–586.
- Pirk, N., Sievers, J., Mertes, J., Parmentier, F.J.W., Mastepanov, M., Christensen, T.R., 2017. Spatial variability of CO₂ uptake in polygonal tundra: assessing low-frequency disturbances in eddy covariance flux estimates. *Biogeosciences* 14 (12), 3157–3169.
- Pirt, S.J., 1965. The maintenance energy of bacteria in growing cultures. *Proc. R. Soc. Lond. Ser. B Biol. Sci.* 163 (991), 224–231.
- Pold, G., Schimel, J.P., Sistla, S.A., 2021. Soil bacterial communities vary more by season than with over two decades of experimental warming in Arctic tussock tundra. *Elementa Sci. Anthropol.* 9 (1).
- Poppeliers, S.W.M., Hefting, M., Dorrepaal, E., Weedon, J.T., 2022. Functional microbial ecology in arctic soils: the need for a year-round perspective. *FEMS Microbiol. Ecol.* 98, [fiac134](https://doi.org/10.1093/femsec/fiac134) <https://doi.org/10.1093/femsec/fiac134>.
- Post, E., Alley, R.B., Christensen, T.R., Macias-Fauria, M., Forbes, B.C., Gooseff, M.N., Iler, A., Kerby, J.T., Laidre, K.L., Mann, M.E., Olofsson, J., 2019. The polar regions in a 2°C warmer world. *Science Advances* 5 (12) p.eaaw9883.
- Raich, J.W., Schlesinger, W.H., 1992. The global carbon dioxide flux in soil respiration and its relationship to vegetation and climate. *Tellus B* 44 (2), 81–99.
- Rapaic, M., Brown, R., Markovic, M., Chaumont, D., 2015. An evaluation of temperature and precipitation surface-based and reanalysis datasets for the Canadian Arctic, 1950–2010. *Atmosphere-Ocean* 53 (3), 283–303.
- Rasmussen, L.H., Zhang, W., Hollesen, J., Cable, S., Christiansen, H.H., Jansson, P., Elberling, B., 2018. Modelling present and future permafrost thermal regimes in Northeast Greenland. *Cold Reg. Sci. Technol.* 146, 199–213.
- Ravn, N.R., Elberling, B., Michelsen, A., 2020. Arctic soil carbon turnover controlled by experimental snow addition, summer warming and shrub removal. *Soil Biol. Biochem.* 142, 107698.
- Rawlins, M.A., Steele, M., Holland, M.M., Adam, J.C., Cherry, J.E., Francis, J.A., Groisman, P.Y., Hinzman, L.D., Huntington, T.G., Kane, D.L., Kimball, J.S., 2010. Analysis of the Arctic system for freshwater cycle intensification: Observations and expectations. *J. Clim.* 23 (21), 5715–5737.
- Raz-Yaseef, N., Torn, M.S., Wu, Y., Billesbach, D.P., Liljedahl, A.K., Kneafsey, T.J., Romanovsky, V.E., Cook, D.R., Wulfschlegler, S.D., 2017. Large CO₂ and CH₄ emissions from polygonal tundra during spring thaw in northern Alaska. *Geophys. Res. Lett.* 44 (1), 504–513.
- Riley, W.J., Subin, Z.M., Lawrence, D.M., Swenson, S.C., Torn, M.S., Meng, L., Mahowald, N.M., Hess, P., 2011. Barriers to predicting changes in global terrestrial methane fluxes: analyses using CLM4Me, a methane biogeochemistry model integrated in CESM. *Biogeosciences* 8 (7), 1925–1953.
- Rivkina, E.M., Friedmann, E.I., McKay, C.P., Gilichinsky, D.A., 2000. Metabolic activity of Permafrost Bacteria below the Freezing Point. *Appl. Environ. Microbiol.* 66 (8), 3230–3233.
- Rixen, C., Høye, T.T., Macek, P., Aerts, R., Alatalo, J.M., Anderson, J.T., Arnold, P.A., Barrio, I.C., Bjerke, J.W., Björkman, M.P., Blok, D., 2022. Winters are changing: snow effects on Arctic and alpine tundra ecosystems. *Arctic Sci.* 8 (3), 572–608.
- Rodrigo, A., Recous, S., Neel, C., Mary, B., 1997. Modelling temperature and moisture effects on C-N transformations in soils: comparison of nine models. *Ecol. Model.* 102 (2–3), 325–339.
- Romanovsky, V.E., Smith, S.L., Shiklomanov, N.I., Streletskiy, D.A., Isaksen, K., Kholodov, A.L., Christiansen, H.H., Drozdov, D.S., Malkova, G.V., Marchenko, S.S., 2017. The Arctic terrestrial permafrost [in “State of the climate in 2016”] State of the climate in 2016. *Bull. Am. Meteorol. Soc.* 98, S1.
- Röbger, N., Sachs, T., Wille, C., Boike, J., Kutzbach, L., 2022. Seasonal increase of methane emissions linked to warming in Siberian tundra. *Nat. Clim. Chang.* 1–6.
- Runkle, B.R., Sachs, T., Wille, C., Pfeiffer, E.M., Kutzbach, L., 2013. Bulk partitioning the growing season net ecosystem exchange of CO₂ in Siberian tundra reveals the seasonality of its carbon sequestration strength. *Biogeosciences* 10 (3), 1337–1349.
- Ruppel, C.D., Kessler, J.D., 2017. The interaction of climate change and methane hydrates. *Rev. Geophys.* 55 (1), 126–168.
- Schadt, C.W., Martin, A.P., Lipson, D.A., Schmidt, S.K., 2003. Seasonal dynamics of previously unknown fungal lineages in tundra soils. *Science* 301, 1359–1361.
- Schimel, J.P., Mikan, C., 2005. Changing microbial substrate use in Arctic tundra soils through a freeze-thaw cycle. *Soil Biol. Biochem.* 37 (8), 1411–1418.
- Schimel, J.P., Weintraub, M.N., 2003. The implications of exoenzyme activity on microbial carbon and nitrogen limitation in soil: a theoretical model. *Soil Biol. Biochem.* 35 (4), 549–563.
- Schindler, D.W., Donahue, W.F., 2006. An impending water crisis in Canada's western prairie provinces. *Proc. Natl. Acad. Sci.* 103 (19), 7210–7216.

- Schmidt, S.K., Wilson, K.L., Meyer, A.F., et al., 2008. Phylogeny and Ecophysiology of Opportunistic “Snow Molds” from a Subalpine Forest Ecosystem. *Microb. Ecol.* 56, 681–687.
- Schmidt, S.K., Wilson, K.L., Monson, R.K., Lipson, D.A., 2009. Exponential growth of “snow molds” at sub-zero temperatures: an explanation for high beneath-snow respiration rates and Q_{10} values. *Biogeochemistry* 95, 13–21.
- Schmidt, M.W., Torn, M.S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I.A., Kleber, M., Kögel-Knabner, I., Lehmann, J., Manning, D.A., Nannipieri, P., 2011. Persistence of soil organic matter as an ecosystem property. *Nature* 478 (7367), 49–56.
- Schnecker, J., Wild, B., Hofhansl, F., et al., 2014. Effects of soil organic matter properties and microbial community composition on enzyme activities in cryoturbated arctic soils. *PLoS One* 9, e94076.
- Schneegurt, M.A., Dore, S.Y., Kulpa, C.F., 2003. Direct extraction of DNA from soils for studies in microbial ecology. *Curr. Issues Mol. Biol.* 5 (1), 1–8.
- Schostag, M., Stibal, M., Jacobsen, S.S., Bælum, T., Taş, N., Elberling, B., Jansson, J.K., Semenchuk, P., Priemé, A., 2015. Distinct summer and winter bacterial communities in the active layer of Svalbard permafrost revealed by DNA-and RNA-based analyses. *Front. Microbiol.* 6, 399.
- Schuur, E.A., McGuire, A.D., Schädel, C., Grosse, G., Harden, J.W., Hayes, D.J., Hugelius, G., Koven, C.D., Kuhry, P., Lawrence, D.M., Natali, S.M., 2015. Climate change and the permafrost carbon feedback. *Nature* 520 (7546), 171–179.
- Segers, R., 1998. Methane production and methane consumption: a review of processes underlying wetland methane fluxes. *Biogeochemistry* 41, 23–51.
- Segers, R., Kengen, S.W.M., 1998. Methane production as a function of anaerobic carbon mineralization: a process model. *Soil Biol. Biochem.* 30 (8), 1107–1117.
- Shao, Y., Taff, G.N., Lunetta, R.S., Weng, Q., 2011. A review of selected moderate-resolution imaging spectroradiometer algorithms, data products, and applications. *Adv. Remote Sensing* 31–55.
- Shastri, A., Sanchez, M., Gai, X., Lee, M.Y., Dewers, T., 2021. Mechanical behavior of frozen soils: Experimental investigation and numerical modeling. *Comput. Geotech.* 138, 104361.
- Shirley, I.A., Mekonnen, Z.A., Grant, R.F., Dafflon, B., Hubbard, S.S., Riley, W.J., 2022. Rapidly changing high-latitude seasonality: implications for the 21st century carbon cycle in Alaska. *Environ. Res. Lett.* 17 (1), 014032.
- Sierra, C.A., Harmon, M.E., Perakis, S.S., 2011. Decomposition of heterogeneous organic matter and its long-term stabilization in soils. *Ecol. Monogr.* 81 (4), 619–634.
- Singh, J.S., Gupta, S.R., 1977. Plant decomposition and soil respiration in terrestrial ecosystems. *Bot. Rev.* 43 (4), 449–528.
- Singh, S.M., Singh, S.K., Yadav, L.S., Singh, P.N., Ravindra, R., 2012. Filamentous soil fungi from Ny-Ålesund, Spitsbergen, and screening for extracellular enzymes. *Arctic* 65 (1), 45–55. <http://www.jstor.org/stable/23187223>.
- Sipes, K., Paul, R., Fine, A., Li, P., Liang, R., Boike, J., Onstott, T.C., Vishnivetskaya, T.A., Schaeffer, S., Lloyd, K.G., 2022. Permafrost active layer microbes from Ny Ålesund, Svalbard (79° N) show Autotrophic and Heterotrophic Metabolisms with Diverse Carbon-Degrading Enzymes. *Front. Microbiol.* 12.
- Sitch, S., Smith, B., Prentice, I.C., Arneth, A., Bondeau, A., Cramer, W., Kaplan, J.O., Levis, S., Lucht, W., Sykes, M.T., Thonicke, K., 2003. Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model. *Glob. Chang. Biol.* 9 (2), 161–185.
- Six, J., Frey, S.D., Thiet, R.K., Batten, K.M., 2006. Bacterial and fungal contributions to carbon sequestration in agroecosystems. *Soil Sci. Soc. Am. J.* 70 (2), 555–569.
- Stieglitz, M., Déry, S.J., Romanovsky, V.E., Osterkamp, T.E., 2003. The role of snow cover in the warming of arctic permafrost. *Geophys. Res. Lett.* 30 (13).
- Stolpovsky, K., Martinez-Lavanchy, P., Heipieper, H.J., Van Cappellen, P., Thullner, M., 2011. Incorporating dormancy in dynamic microbial community models. *Ecol. Model.* 222 (17), 3092–3102.
- Sturm, M., Schimel, J., Michaelson, G., Welker, J.M., Oberbauer, S.F., Liston, G.E., Fahnestock, J., Romanovsky, V.E., 2005. Winter biological processes could help convert arctic tundra to shrubland. *Bioscience* 55 (1), 17–26.
- Sulman, B.N., Phillips, R.P., Oishi, A.C., Shevliakova, E., Pacala, S.W., 2014. Microbe-driven turnover offsets mineral-mediated storage of soil carbon under elevated CO₂. *Nat. Clim. Chang.* 4 (12), 1099–1102.
- Tan, Z., Zhuang, Q., Henze, D.K., Frankenberg, C., Dlugokencky, E., Sweeney, C., Turner, A.J., Sasakawa, M., Machida, T., 2016. Inverse modeling of pan-Arctic methane emissions at high spatial resolution: what can we learn from assimilating satellite retrievals and using different process-based wetland and lake biogeochemical models? *Atmos. Chem. Phys.* 16 (19), 12649–12666.
- Tang, J., Riley, W.J., 2015. Weaker soil carbon-climate feedbacks resulting from microbial and abiotic interactions. *Nat. Clim. Chang.* 5 (1), 56–60.
- Tang, J., Riley, W.J., 2019. A theory of effective microbial substrate affinity parameters in variably saturated soils and an example application to aerobic soil heterotrophic respiration. *J. Geophys. Res. Biogeosci.* 124 (4), 918–940.
- Tarnocai, C., Canadell, J.G., Schuur, E.A., Kuhry, P., Mazhitova, G., Zimov, S., 2009. Soil organic carbon pools in the northern circumpolar permafrost region. *Glob. Biogeochem. Cycles* 23 (2).
- Teepe, R., Ludwig, B., 2004. Variability of CO₂ and N₂O emissions during freeze-thaw cycles: results of model experiments on undisturbed forest-soil cores. *J. Plant Nutr. Soil Sci.* 167 (2), 153–159.
- Thomas, H.R., Cleall, P., Li, Y.C., et al., 2009. Modeling of cryogenic processes in permafrost and seasonally frozen soils. *Geotechnique* 59 (3), 173–184.
- Thormann, M.N., Currah, R.S., Bayley, S.E., 2003. Succession of microfungial assemblages in decomposing peatland plants. *Plant Soil* 250, 323–333.
- Tian, H., Xu, X., Liu, M., Ren, W., Zhang, C., Chen, G., Lu, C., 2010. Spatial and temporal patterns of CH₄ and N₂O fluxes in terrestrial ecosystems of North America during 1979–2008: application of a global biogeochemistry model. *Biogeosciences* 7 (9), 2673.
- Todd-Brown, K.E., Hopkins, F.M., Kivlin, S.N., Talbot, J.M., Allison, S.D., 2012. A framework for representing microbial decomposition in coupled climate models. *Biogeochemistry* 109, 19–33.
- Todd-Brown, K.E., Randerson, J.T., Post, W.M., Hoffman, F.M., Tarnocai, C., Schuur, E.A., Allison, S.D., 2013. Causes of variation in soil carbon simulations from CMIP5 Earth system models and comparison with observations. *Biogeosciences* 10 (3), 1717–1736.
- Todd-Brown, K.E.O., Randerson, J.T., Hopkins, F., Arora, V., Hajima, T., Jones, C., Shevliakova, E., Tjiputra, J., Volodin, E., Wu, T., Zhang, Q., 2014. Changes in soil organic carbon storage predicted by Earth system models during the 21st century. *Biogeosciences* 11 (8), 2341–2356.
- Tojo, M., Newsham, K.K., 2012. Snow moulds in polar environments. *Fungal Ecol.* 5, 395–402.
- Tomczyk, A.M., Bednorz, E., Szyga-Pluta, K., 2021. Changes in air temperature and snow cover in winter in Poland. *Atmosphere* 12 (1), 68.
- Treat, C.C., Bloom, A.A., Marushchak, M.E., 2018. Nongrowing season methane emissions—a significant component of annual emissions across northern ecosystems. *Glob. Chang. Biol.* 24 (8), 3331–3343.
- Treseder, K.K., Balsler, T.C., Bradford, M.A., Brodie, E.L., Dubinsky, E.A., Eviner, V.T., Hofmoeckel, K.S., Lennon, J.T., Levine, U.Y., MacGregor, B.J., Pett-Ridge, J., 2012. Integrating microbial ecology into ecosystem models: challenges and priorities. *Biogeochemistry* 109, 7–18.
- Turetsky, M.R., Abbott, B.W., Jones, M.C., Anthony, K.W., Olefeldt, D., Schuur, E.A., Grosse, G., Kuhry, P., Hugelius, G., Koven, C., Lawrence, D.M., 2020. Carbon release through abrupt permafrost thaw. *Nat. Geosci.* 13 (2), 138–143.
- Uchida, M., Kishimoto, A., Muraoka, H., Nakatsubo, T., Kanda, H., Koizumi, H., 2010. Seasonal shift in factors controlling net ecosystem production in a high Arctic terrestrial ecosystem. *J. Plant Res.* 123 (1), 79–85.
- Vepraskas, M.J., Faulkner, S.P., 2001. In: Richardson, J.L., Vepraskas, M.J. (Eds.), *Redox chemistry of hydric soils*. Wetland Soils CRC Press, Boca Raton, FL, p. 85 to 105.
- Villarreal, S., Hollister, R.D., Johnson, D.R., Lara, M.J., Webber, P.J., Tweedie, C.E., 2012. Tundra vegetation change near Barrow, Alaska (1972–2010). *Environ. Res. Lett.* 7 (1), 015508.
- Waldrop, M.P., McFarland, W., Manies, K.L., Leewis, M.C., Blazewicz, S.J., Jones, M.C., Neumann, R.B., Keller, J.K., Cohen, L., Euskirchen, E.S., Edgar, C., 2021. Carbon fluxes and microbial activities from boreal peatlands experiencing permafrost thaw. *J. Geophys. Res. Biogeosci.* 126 (3). [p.e2020JG005869](https://doi.org/10.1029/2020JG005869).
- Wallenstein, M.D., McMahon, S., Schimel, J., 2007. Bacterial and fungal community structure in Arctic tundra tussock and shrub soils. *FEMS Microbiol. Ecol.* 59 (2), 428–435.
- Walvoord, M.A., Striegl, R.G., 2021. Complex Vulnerabilities of the Water and Aquatic Carbon Cycles to Permafrost Thaw. *Front. Clim.* 3, 730402 <https://doi.org/10.3389/fclim.2021.730402>.
- Wang, Y.P., Law, R.M., Pak, B., 2010. A global model of carbon, nitrogen and phosphorus cycles for the terrestrial biosphere. *Biogeosciences* 7 (7), 2261–2282.
- Wang, G., Mayes, M.A., Gu, L., Schadt, C.W., 2014. Representation of dormant and active microbial dynamics for ecosystem modeling. *PLoS One* 9 (2), e89252.
- Wania, R., 2007. *Modelling Northern Peatland Land Surface Processes, Vegetation Dynamics and Methane Emissions* (Doctoral dissertation. University of Bristol).
- Wania, R., Ross, I., Prentice, I.C., 2010. Implementation and evaluation of a new methane model within a dynamic global vegetation model: LPJ-WHYMe v1. 3.1. *Geosci. Model Dev.* 3 (2), 565–584.
- Waring, B.G., Averill, C., Hawkes, C.V., 2013. Differences in fungal and bacterial physiology alter soil carbon and nitrogen cycling: insights from meta-analysis and theoretical models. *Ecol. Lett.* 16 (7), 887–894.
- Watanabe, K., Kugisaki, Y., 2017. Effect of macropores on soil freezing and thawing with infiltration. *Hydrol. Process.* 31, 270–278.
- Webb, E.E., Schuur, E.A., Natali, S.M., Oken, K.L., Bracho, R., Krapek, J.P., Risk, D., Nickerson, N.R., 2016. Increased wintertime CO₂ loss as a result of sustained tundra warming. *J. Geophys. Res. Biogeosci.* 121 (2), 249–265.
- Wei, D., et al., 2021. Plant uptake of CO₂ outpaces losses from permafrost and plant respiration on the Tibetan Plateau. *Proc. Natl. Acad. Sci.* <https://doi.org/10.1073/pnas.2015283118>.
- Weintraub, M.N., Scott-Denton, L.E., et al., 2007. The effects of tree rhizodeposition on soil exoenzyme activity, dissolved organic carbon, and nutrient availability in a subalpine forest ecosystem. *Oecologia* 154, 327–338.
- Wieder, W.R., Grandy, A.S., Kallenbach, C.M., Taylor, P.G., Bonan, G.B., 2015. Representing life in the Earth system with soil microbial functional traits in the MIMICS model. *Geosci. Model Dev.* 8 (6), 1789–1808.
- Wieder, W.R., Sulman, B.N., Hartman, M.D., Koven, C.D., Bradford, M.A., 2019. Arctic soil governs whether climate change drives global losses or gains in soil carbon. *Geophys. Res. Lett.* 46 (24), 14486–14495.
- Wik, M., Varner, R.K., Anthony, K.W., MacIntyre, S., Bastviken, D., 2016. Climate-sensitive northern lakes and ponds are critical components of methane release. *Nat. Geosci.* 9 (2), 99–105. <https://doi.org/10.1038/ngeo2578>.
- Xu, X., Tian, H., 2012. Methane exchange between marshland and the atmosphere over China during 1949–2008. *Glob. Biogeochem. Cycles* 26 (2).
- Xu, X., Yuan, F., Hanson, P.J., Wullschlegel, S.D., Thornton, P.E., Riley, W.J., Graham, D.E., 2016. Reviews and syntheses: four decades of modeling methane cycling in terrestrial ecosystems. *Biogeosciences* 13 (12), 3735.
- Yang, K., Wang, C., Li, S., 2018. Improved simulation of frozen-thawing process in land surface model (CLM4. 5). *J. Geophys. Res. Atmos.* 123 (23), 13–238.

- Ye, X., Yu, Z., Zhang, Y., Kang, J., Wu, S., Yang, T., Gao, P., 2022. Mineral Composition Impact on the thermal Conductivity of Granites based on Geothermal Field experiments in the Songliao and Gonghe Basins, China. *Minerals* 12, 247.
- Yi, Y., Kimball, J.S., Watts, J.D., Natali, S.M., Zona, D., Liu, J., Ueyama, M., Kobayashi, H., Oechel, W., Miller, C.E., 2020. Investigating the sensitivity of soil heterotrophic respiration to recent snow cover changes in Alaska using a satellite-based permafrost carbon model. *Biogeosciences* 17 (22), 5861–5882.
- Yokohata, T., Saito, K., Takata, K., et al., 2020. Model improvement and future projection of permafrost processes in a global land surface model. *Prog Earth Planet Sci* 7, 69.
- Yoshitake, S., Uchida, M., Koizumi, H., Nakatsubo, T., 2007. Carbon and nitrogen limitation of soil microbial respiration in a High Arctic successional glacier foreland near Ny-Ålesund, Svalbard. *Polar Res.* 26 (1), 22–30.
- Yu, L., Ahrens, B., Wutzler, T., Schrumppf, M., Zaehle, S., 2020. Jena Soil Model (JSM v1.0; revision 1934): a microbial soil organic carbon model integrated with nitrogen and phosphorus processes. *Geosci. Model Dev.* 13 (2), 783–803.
- Zha, J., Zhuang, Q., 2018. Microbial decomposition processes and vulnerable arctic soil organic carbon in the 21st century. *Biogeosciences* 15 (18), 5621–5634.
- Zhang, T., 2005. Influence of the seasonal snow cover on the ground thermal regime: an overview. *Rev. Geophys.* 43 (4).
- Zhang, W., Jansson, P.E., Sigsgaard, C., McConnell, A., Jammet, M.M., Westergaard-Nielsen, A., Lund, M., Friberg, T., Michelsen, A., Elberling, B., 2019. Model-data fusion to assess year-round CO₂ fluxes for an arctic heath ecosystem in West Greenland (69° N). *Agric. For. Meteorol.* 272, 176–186.
- Zhou, J., Wei, C., Lai, Y., Wei, H., Tian, H., 2018. Application of the Generalized Clapeyron Equation to Freezing Point Depression and Unfrozen Water Content. *Water Resour. Res.* 54 (11), 9412–9431.
- Zhu, Q., Zhuang, Q., 2013. Influences of calibration data length and data period on model parameterization and quantification of terrestrial ecosystem carbon dynamics. *Geosci. Model Dev. Discuss.* 6 (4), 6835–6865.
- Zhu, Q., Liu, J., Peng, C., Chen, H., Fang, X., Jiang, H., Yang, G., Zhu, D., Wang, W., Zhou, X., 2014. Modelling methane emissions from natural wetlands by development and application of the TRIPLEX-GHG model. *Geosci. Model Dev.* 7 (3), 981–999.
- Zhu, D., Ciais, P., Krinner, G., et al., 2019. Controls of soil organic matter on soil thermal dynamics in the northern high latitudes. *Nat. Commun.* 10, 3172.
- Zhuang, Q., Romanovsky, V.E., McGuire, A.D., 2001. Incorporation of a permafrost model into a large-scale ecosystem model: Evaluation of temporal and spatial scaling issues in simulating soil thermal dynamics. *J. Geophys. Res. Atmos.* 106 (D24), 33649–33670.
- Zhuang, Q., Melillo, J.M., Kicklighter, D.W., Prinn, R.G., McGuire, A.D., Steudler, P.A., Felzer, B.S., Hu, S., 2004. Methane fluxes between terrestrial ecosystems and the atmosphere at northern high latitudes during the past century: a retrospective analysis with a process-based biogeochemistry model. *Glob. Biogeochem. Cycles* 18 (3).
- Zimov, S.A., Davidov, S.P., Voropaev, Y.V., Prosiannikov, S.F., Semiletov, I.P., Chapin, M.C., Chapin, F.S., 1996. Siberian CO₂ efflux in winter as a CO₂ source and cause of seasonality in atmospheric CO₂. *Clim. Chang.* 33, 111–120.
- Zinger, L., Shahnava, B., Baptist, F., Geremia, R.A., Choler, P., 2009. Microbial diversity in alpine tundra soils correlates with snow cover dynamics. *ISME J.* 3 (7), 850–859.
- Zona, D., Gioli, B., Commane, R., Lindaas, J., Wofsy, S.C., Miller, C.E., Dinardo, S.J., Dengel, S., Sweeney, C., Karion, A., Chang, R.Y.W., 2016. Cold season emissions dominate the Arctic tundra methane budget. *Proc. Natl. Acad. Sci.* 113 (1), 40–45.