



**British
Geological Survey**

NATURAL ENVIRONMENT RESEARCH COUNCIL

DTI Strategic Environmental Assessment Area 7 (SEA7) Geological Metadata

Continental Shelf & Margins Programme

Commissioned Report CR/02/275N



BRITISH GEOLOGICAL SURVEY

COMMISSIONED REPORT CR/02/275 N

DTI Strategic Environmental Assessment Area 7 (SEA7) Geological Metadata

Richard Holmes¹ Jan Fraser¹, Vikki Gunn², Paul Henni¹, Colin
Jacobs², Pat Shannon³, Vikram Unnithan³

Other contributors: Rhys Cooper¹, Alexander Downie⁴, Colin
Graham¹, Tim Lebas², Anne Richardson¹, John Ridgway¹, Steve
Rowlatt⁵, Alan Stevenson¹

Compiler

Richard Holmes

Key words

Metadata, Rockall Trough,
environment, habitat, geology

Front cover

Area of DTI Strategic
Environmental Assessment 7

Bibliographical reference

HOLMES, R, FRASER, J, GUNN,
V, HENNI, P, JACOBS, G,
SHANNON, P. 2002. DTI
Strategic Environmental
Assessment area 7 (SEA7)
Geological Metadata. *British
Geological Survey Commissioned
Report*, CR/02/275N.

1. British Geological Survey, Edinburgh
2. Southampton Oceanography Centre
3. University College Dublin
4. Scottish Natural Heritage
5. Centre for Environment, Fisheries and Aquaculture Science

DTI Contract number SEA678_data_01

BGS Project number ERE73900467

Foreword

This report is the published product of a study by the British Geological Survey (BGS) for research funded by the Department of Trade and Industry Offshore Environment and Decommissioning, Licensing and Consents Unit.

Acknowledgements

In addition to the authors and named contributors many other BGS staff have contributed to the metadata by providing references sourced from unpublished reports. These contributions are gratefully acknowledged.

Contents

1. Introduction.....	1
2. Synopsis.....	1
3. Methods.....	2
4. Sources of metadata.....	3
Non-bibliographic	3
Other websites	21
Endnote © format.....	23
5. References	24
Table 1. Cruises held on the SOC SeaDOG database with seabed and shallow subseabed data.....	19
Table 2. Non-BGS/SOC databases with references to cruises/programmes	22
Table 3. Non BGS/SOC sample databases.	23
Table 4. Marine Telephone Cables.....	23
Figure 1. DTI UK Strategic Environmental Assessment areas	3
Figure 2. Physiography and selected geographical names	4
Figure 3. Oil exploration infrastrucure	5
Figure 4. Unpublished geological reports	5
Figure 5. Unpublished environmental reports	6
Figure 6. Unpublished site investigation reports	7
Figure 7. Unpublished geotechnics/geohazards reports	8
Figure 8. Published offshore regional reports	8

Figure 9. Regional survey: single channel air gun	9
Figure 10. Regional survey: single channel sparker	9
Figure 11. Regional survey: boomer	10
Figure 12. Regional survey: pinger	10
Figure 13. Regional survey: sidescan sonar	11
Figure 14. BGS Rockall Consortium: regional surveys	11
Figure 15. NERC and IFREMER multibeam survey, WFA 3D seabed picks	12
Figure 16. BGS gravity core sites	13
Figure 17. BGS vibrocore sites	13
Figure 18. BGS 1m/5m rockdrill sites	14
Figure 19. BGS and SOC Van Veen, multicore and spadecore (other core OC)	14
Figure 20. BGS Shipek Grab sites	15
Figure 21. BGS maps of solid geology, Quaternary geology, seabed sediment	15
Figure 22. BGS digital regional seabed sediment	16
Figure 23. BGS regional digital bathymetric data	16
Figure 24. BGS inorganic and organic geochemistry	17
Figure 25. Hydrographic Office bathymetry, sidescan sonar, sediment sample data and interpretation.	17
Figure 26. SOC project surveys: mainly Gloria and Tobi sidescan survey	18
Figure 27. Scottish Natural Heritage Special Areas of Conservation	20
Figure 28. Joint Nature Conservancy Council: Project themes and locations	20
Appendix 1. Geological processes	25
Appendix 2. Project liaison	28
Appendix 3. Keywords used in searching the bibliographic database	30
Appendix 4. Bibliography	33
Appendix 5. Occurrence of Carbonate and other mounds in the SEA7 region of the Atlantic Margin.....	87

1. Introduction

The Department of Trade and Industry Offshore Environment and Decommissioning, Licensing and Consents Unit (hereafter DTI) is responsible to the UK government for administering environmental legislation applicable to the oil, gas and renewable energy industry offshore. One such piece of legislation is EC Directive (2001/42/EEC). The EC Directive is generic and wide ranging 'on the assessment of the effects and certain plans and programs on the environment.' The activities regulated include seismic surveys, drilling, other seabed development operations and the decommissioning of offshore development structures. The programme of DTI Strategic Environmental Assessments is aimed at regional assessments of the possible impacts of offshore developments prior to new licensing rounds. The information from the strategic environmental surveys is therefore required to provide the technical basis on whether or not to grant commercial licences and consents for new offshore developments in the energy sector.

The objective of this research is to consider the scope of geological data that may contribute to the technical basis for the strategic environmental assessments:

1. Synopsis
2. Methods
3. Sources of metadata: scope, in terms of geography and type, where the original data is stored, the quality, issues of costs and licensing
4. Metadata

The area of study extends offshore from the coastlines of Scotland and Northern Ireland and across the NE Atlantic Ocean to the western limits of the UK exclusive zone (Figure 1). The seabed habitats are truly oceanic seawards of the shelfbreak and are currently within the category of frontier research areas.

The main part of this BGS report was completed in November 2002 and is supplemented by separately commissioned reports:

1. July 2003, the geological model for seabed mounds in the UK SEA7 and the Irish Sector (Unnithan, V. and Shannon, P. 2003. SEA7 Geology. *Report Marine and Petroleum Geology Research Group*. Dublin: Department of Geology, University College).
2. August 2003, summary of data that crosses the UK side of the SEA 6, 7 and 8 boundaries to approximately 50km within the Irish border (Wheeler, A. 2003. UK-Irish border public domain geological survey metadata. (Cork: University College, Department of Geology and Environmental Research)): inserted as Appendix 3 TAPPIN, D. R., WHEELER, A., ROWLANDS, K., JENKINS, G. AND SLATER, M. 2001. DTI Strategic Environmental Assessment Area 6 (SEA 6) Geological Metadata. *British Geological Survey*
3. *Commissioned Report*, CR/02/287. 17pp.

The reports listed above update some of the references compiled by the BGS in November 2002.

2. Synopsis

An inner physiographical zone extends seawards from a fjord-like coastline to encompass the continental shelf (mainly the Hebrides Shelf), the outer limit of which is defined by a shelfbreak in a range of approximately 140m to 400m or more water depth. An overall deeper-water middle physiographical zone includes the Rockall Trough, Rockall Bank and Rockall Plateau and smaller banks and seamounts such as the George Bligh Bank and Rosemary Bank. An outer physiographical zone includes the Hatton Bank, Endymion Spur and Maury Channel (Figure 2). The wide variety of the environments and the large area of SEA7 have more or less influenced the methodology used to compile the metadata (Section 3).

Prior to the 1980s many of the earliest publications useful to understanding the composition of seabed sediments and the processes forming the seabed sediments were derived from interpretations of single-beam topographic surveys, sidescan sonar images recorded on paper, single and multi-channel reflection seismic and core and grab sample surveys over the basin margins. The most intensive of the seabed surveys were in places where the oil-prospective rock formations crop or subcrop in relatively shallow water on the continental shelf. Subsequently, publications on seabed geology more or less accelerated with time with new development scenarios. These included more efficient ways of acquiring and processing survey data, the expansion of hydrocarbons, communications, fisheries, waste disposal, and defence activities, many extending to new deep-water environments. New agreements on the definition of exclusive national economic zones also contributed to the amount of offshore geological data collected. Increasing publication rates also originated from marine conservation research, some related to the development of tourism.

Some of the earliest UK seabed strategic research was initially funded by the Department of Energy and was completed in 1995 as part of the systematic regional BGS mapping and reporting programme to establish a geological framework for licensing and other development scenarios on the continental shelf and slope. Research for the exploitation of new and established resources continues. For example, there is increasing desire to develop the world-class potential in SEA7 for renewable (wind, wave) energy resources. The burgeoning diversity of development pressures on the environment has meant that measures for coastal and nearshore conservation areas are continuing, but that there are also new provisions for adoption of special areas of conservation in the deep-water environments (e.g. Darwin Mounds, NE Rockall Trough).

3. Methods

Appendix 2 summarises the network of those researchers who participated to review and input the data for this report.

Metadata outputs are in Endnote © version 6 format. For the bibliographic datasets Endnote files were compiled from BGS and SOC research with principal data merged or extracted from the GeoRef and the Gearchive and Aslib online databases, the BGS regional offshore reports series, Western Frontiers Association bibliography compiled in ACCESS format, Stratagem (EU OMARC project cluster) and other publications in MS word. Online geological searches were linked to geographical keywords (Appendix 3). The bibliographic search was completed in November 2002. The breakdown of the geological/geophysical search categories is further explained in Section 4 below.

Maps are considered essential report outputs where the Endnote © format is unsuitable for clearly illustrating the geographical extent, density or interrelationships of the metadata. This is

particularly pertinent to the deep-water and remote regions where much of the seabed habitat is unexplored or patchily reconnoitred. Maps were thus derived from information stored in BGS files in ARCVIEW, ORACLE-GMT and ACCESS formats and from geographical coordinates of the regional geophysical data supplied by the Southampton Oceanography Centre. The BGS offshore science GIS database covers the whole UK zone. The regional metadata from the various sources were clipped to fit the area of SEA7 and output as a series of thematic maps for this report.

To accord with the contract, this report is output in hard-copy and compact disc formats.

4. Sources of metadata

Non-bibliographic

The information in this section has been presented in map form for the reasons explained in Section 3 above.

There are licensing or copyright issues for raw and processed profile and plan data (e.g. Hydrographic Office sonar and bathymetric data) and with interpreted data perhaps with intellectual property rights attached (e.g. BGS digital bathymetry, seabed sediment and geochemical data). Sources of metadata in both the ownership and geographical sense have been merged into the figures. Further explanations of the figures and/or the metadata links are given in the text below the figure titles.

In the following reference has been made to the generic link www.bgs.ac.uk whereas specific offshore data can be accessed at <http://www.bgs.ac.uk/discoverymetadata/home.html>.




Figure 1. DTI UK Strategic Environmental Assessment areas www.offshore-sea.org.uk

The UK zone is organised into 8 SEA areas. Aside from geology the reports for these incorporate aspects of habitat biotope, hydrography and chemistry the details of which are identified within the DTI-funded website for the programme managed by Geotek Ltd www.offshore-sea.org.uk




Figure 2. Physiography and selected geographical names

www.bodc.ac.uk/projects/gebco97.html




Figure 3. Oil exploration infrastructure

Aspects of development infrastructure and the availability of commercial hydrocarbons exploration and development data for the UK zone are incorporated into the DEAL programme managed by UKOAA for BGS. www.ukdeal.co.uk




Figure 4. Unpublished geological reports www.bgs.ac.uk

Figure 4 is derived from the Oracle database: Non-confidential BGS reports or reports likely to be released without charge into the public domain (grey literature). The data include some WFA bibliography and other dedicated databases as appropriate.




Figure 5. Unpublished environmental reports www.bgs.ac.uk

Figure 5 is derived from the Oracle database and shows the potential scope for non-confidential BGS reports describing environmental aspects of the seabed and shallow sub-surface.




Figure 6. Unpublished Site investigation reports www.bgs.ac.uk

The BGS Oracle Database contains reports describing the geology and geotechnics of specific sites related to drilling and civil engineering. These are currently archived in the BGS as area polygons with no indication of content and have been registered with the BGS following UKOOA recommendations. There is currently (November, 2002) no method of retrieving the detail of the quality and content of these reports without retrieving them as the hard-copy report or, more recently, as CDs. Well site investigations: these would typically consist of a 3x3 or 1x1km area surveyed with single or multi channel mini-sleeve/air gun, sparker, pinger / boomer / echosounder / sidescan sonar, with some interpretation calibrated by core. More problematic deep-water site investigations may have employed seabed photography, some with Autonomous Underwater Vehicles. The release of most of this data would require the commercial owners permission and the reports very rarely contain the raw datafiles used by the contractor.




Figure 7. Unpublished geotechnics/geohazards reports www.bgs.ac.uk

Oracle Database: Reports describing the geology and geotechnics of specific sites or larger areas




Figure 8. Published offshore regional reports www.bgs.ac.uk




Figure 9. Regional survey: single channel air gun www.bgs.ac.uk




Figure 10. Regional survey: single channel sparker www.bgs.ac.uk




Figure 11. Regional survey: boomer www.bgs.ac.uk




Figure 12. Regional survey: pinger www.bgs.ac.uk




Figure 13. Regional survey: sidescan sonar www.bgs.ac.uk




Figure 14. BGS Rockall Consortium: regional surveys www.bgs.ac.uk

Mainly single channel airgun with various combinations of high resolution reflection seismic but excluding sidescan sonar.




Figure 15. NERC and IFREMER multibeam survey, WFA 3D seabed picks www.bgs.ac.uk
www.ifremer.fr

The NERC acquired data originate from the Land Ocean Interaction/ Shelfedge study. IFREMER-Regional surveys of swath bathymetry and seabed backscatter are related to deep-water fisheries research. BGS created images of shaded relief by processing seabed picks from 3D commercial seismic exploration surveys on behalf of the Western Frontier Association.




Figure 16. BGS gravity core sites www.bgs.ac.uk




Figure 17. BGS vibrocore sites www.bgs.ac.uk




Figure 18. BGS 1m/5m rockdrill sites www.bgs.ac.uk




Figure 19. BGS and SOC Van Veen, multicore and spadecore (other core OC) www.bgs.ac.uk, www.boscor.org, (boscor temporarily out of order October 2002)




Figure 20. BGS Shipek Grab sites www.bgs.ac.uk




Figure 21. BGS maps solid geology, Quaternary geology, seabed sediment www.bgs.ac.uk
 Hard-copy maps 1:250, 000 and 1:500,000 scale of solid geology Rockall Trough




Figure 22. BGS digital regional seabed sediments. www.bgs.ac.uk
 Folk classes based on proportions of Wentworth grain size scales.




Figure 23. BGS regional digital bathymetric data www.bgs.ac.uk




Figure 24. BGS inorganic and organic chemistry www.bgs.ac.uk




Figure 25. Hydrographic Office bathymetry, sidescan sonar, sediment sample data and interpretation www.bgs.ac.uk

Availability from the BGS varies. BGS holds some data in arrears from the Hydrographic Office which has the up-to-date holdings. Arrangements prior to 1995 meant that much Hydrographic Office data had been incorporated into the BGS seabed sediment maps published at 1:250,000 scale. Arrangements after 1995 are that the raw data are subject to licensing agreements.




Figure 26. SOC project surveys: mainly Gloria and Tobi sidescan survey

<http://www.soc.soton.ac.uk/cgi-bin/seadog/seadog.pl>

As for the BGS data summarised above, the data classes within the SOC project surveys can be subdivided. The SOC survey data has been acquired with more emphasis on seabed project-specific regional deep-water sidescan and swath bathymetric surveys, sometimes accompanied by pinger or chirp profiles. Expanded key to Figure 26, see Table 1.

Cruise Name or Number	Data Type(s)	Date	Data Owner
R/V Colonel Templar 0198	30 kHz TOBI Sidescan Imagery, 3.5 kHz profiles, 7.5 kHz profiles	May 98	AFEN
RRS Discovery 123	6.5 kHz GLORIA Sidescan Imagery, 12 kHz echo-sounder profiles	Sep 81	SOC
R/V Pelagia 0198	30 kHz TOBI Sidescan Imagery, 7.5 kHz profiles	Jul 98	IRISH DATA
Kommander Subsea (BTEL0187)	6.5 kHz GLORIA Sidescan Imagery, 12 kHz echo-sounder profiles(?)	Feb-Mar 87	DTI
RRS Charles Darwin 119L1	30 kHz TOBI Sidescan Imagery, 3.5 kHz profiles, 7.5 kHz profiles	Jun 99	DTI
RRS Charles Darwin 119L2	30 kHz TOBI Sidescan Imagery, 3.5 kHz profiles, 7.5 kHz profiles	Aug-Sep 99	DTI
RRS Charles Darwin 123L2	30 kHz TOBI Sidescan Imagery, 3.5 kHz profiles, 7.5 kHz profiles	Jun-Jul 00	DTI
RRS Charles Darwin 91	30 kHz TOBI Sidescan Imagery, 3.5 kHz profiles, 7.5 kHz profiles	Feb-Mar 95	SOC
R/V Colonel Templar 0298	30 kHz TOBI Sidescan Imagery, 3.5 kHz profiles, 7.5 kHz profiles	May-Jun 98	AFEN
RRS Discovery 248	410 kHz Sidescan imagery, 12 kHz echo-sounder	Jul-Aug 98	SOC
RRS Discovery 78L1	12 kHz echo-sounder	Sep 76	SOC
RRS Charles Darwin 118	SIMRAD EM12 multibeam bathymetry and acoustic backscatter	Apr-May 99	DTI (UNCLOS)
Cruises NOT on the SeaDOG database, but potentially accessible through SOC			
R/V Professor Logachev TTR7 Cruise	Various 3.5 kHz, sidescan sonar and bathymetry	Jul-Aug 97	TTR (contact via SOC)
R/V Professor Logachev TTR8 Cruise	Various 3.5 kHz, sidescan sonar and bathymetry, video images	98	TTR (contact via SOC)
R/V Professor Logachev TTR9 Cruise	Various 3.5 kHz, sidescan sonar and bathymetry, cores	Jun-Jul 99	TTR (contact via SOC)
R/V Professor Logachev TTR10 Cruise	Various 3.5 kHz, sidescan sonar and bathymetry, video images, samples	Jul-Aug 00	TTR (contact via SOC)
R/V Professor Logachev TTR12 Cruise	Various 3.5 kHz, sidescan sonar and bathymetry, video images, samples	Jun-Jul 02	TTR (contact via SOC)

Table 1. List of Cruises held on the SOC SeaDOG database with seabed and shallow subseabed data <http://www.soc.soton.ac.uk/cgi-bin/seadog/seadog.pl>

Seabed habitat surveys have been completed of Special Areas of Conservation (SACs), and maps of these surveys include integration of seabed geology with biotopes. The areas of offshore interest extend from the coast to 12 nautical miles offshore (Figure 27).




Figure 27. Scottish Natural Heritage Special Areas of Conservation. www.snh.org.uk




Figure 28. Joint Nature Conservancy Council: Project themes, locations www.jncc.gov.uk.

Regional research programmes, for example for the Joint Nature Conservancy Council (Figure 28), have been completed as desk studies using existing data. Note that some of the mound fields illustrated on Figure 28 extend across the UK-Irish border. These mounds are more numerous and extensive in the Irish sector and are the subject of a special report completed for SEA7 some 9 months later than this report (Unnithan, V. and Shannon, P. 2003. SEA7. Geology. Marine and Petroleum Geology Research Group . Department of Geology, University College, Dublin).

A pilot scheme (not illustrated) was launched October 2002 with a whole ecosystem approach to regional sea management. This is being funded by DEFRA for the Irish Sea and the northern boundary of this scheme extends from the south end on the Mull of Kintyre to Rathlin in Northern Ireland.

Other websites

PAN EUROPEAN

A huge dataset of European metadata is likely to exist, some of which may be too risky and time consuming (costly) to be of use for environmental assessment. There was not enough money in the project to examine the full potential to see if these other European resources could add significantly to this project.

- GEIXS (Geological Information Exchange System) <http://geixs.brgm.fr/>
- EU-SEASED www.eu-seased.net. The EU-SEASED website consists of metadata from the following EC 4th and 5th Framework projects
 - EUMARSIN (European Marine Sediment Information Network)
 - EUROSEISMIC (European Marine Seismic Metadata and Information Centre)
 - EUROCORE (A searchable Internet database of seabed samples from the Ocean Basins held at European Institutions)
- SEASEARCH (Gateway to Oceanographic and Marine Data & Information in Europe) www.sea-search.net. Includes:
 - EDMED (European Directory of Marine Environmental Datasets)
- PANGAEA <http://www.pangaea.de/>

PANGAEA is a public data library on the Internet aimed at archiving, publishing and distributing geo-coded data with special emphasis on environmental, marine and geological research. It is operated by the Alfred Wegener Institute for Polar and Marine Research and the Centre for Marine Environmental Sciences at the University of Bremen.

- PAN-NATIONAL AGENCY/DEPARTMENT/UNIVERSITY

- NGDF National Geospatial Data Framework (includes 'ask giraffe'), mainly for land based data but extending to the coast www.ngdf.org.uk
- UKMIC UK Marine Information Council www.ukmarine.org
- IACMST . The Inter-Agency Committee on Marine Science and Technology <http://www.marine.gov.uk/>

IACMST is a UK Government Committee reporting to the Office of Science and Technology. IACMST is responsible for the Marine Environmental Data Action Group (MEDAG), which, together with the Marine Environmental Data Co-ordinator, forms the UK Marine Environmental Data (UKMED) Network. The network has set up the [OceanNET](http://www.oceannet.org/) (<http://www.oceannet.org/>) web site as a portal to data and information about the marine environment. OceanNET also contains a new UK Directory of Coastal Data Sets. UKMED is currently funded by the Defence Science and Technology Laboratory (DSTL), Department for Environment, Food and Rural Affairs (DEFRA), the Environment Agency (EA), Fisheries Research Service (FRS), the Met. Office, the Natural Environment Research Council (NERC) and the UK Hydrographic Office (UKHO).

- Marine equivalent of MAGIC www.magic.gov.uk, has been flagged as a possible way forward for collating UK environmental data with www.cefas.co.uk possibly as a front runner.

- INTRA-RESEARCH COUNCIL/UNIVERSITY

- www.NERC.ac.uk/data/

- INTRA-SURVEY/INSTITUTION

- www.bgs.ac.uk BGS Intranet/Geoscience/Metadata, referred to above for the offshore areas but also covers the coastal boundary
- SOC <http://www.soc.soton.ac.uk/cgi-bin/seadog/seadog.pl>).

Table 2: Non-BGS/SOC databases with references to cruises/programmes

Database Name	URL
British Oceanographic Data Centre	www.bodc.ac.uk
US National Geophysical Data Centre	www.ngdc.noaa.gov

European Directory of Marine Environmental Data (EDMED)	http://www.bodc.ac.uk/frames/index4.html?../services/edmed/index.html&2
---	---

Table 3. List of non BGS/SOC sampling databases.

There are too many samples of many different types and age outside of the BGS and SOC sites referred to above to make a sensible list for this report.

Database Name	URL
US National Geophysical Data Centre	www.ngdc.noaa.gov
1.1 Lamont-Doherty Deep-sea Sample Repository	www.ldeo.columbia.edu/CORE_REPOSITORY/RPH1.html

Table 4. Marine Telephone Cables.

Almost all of the data collected during survey and installation remains as proprietary data and requests for use must be addressed to owners/operators.

Cable Name	European Terminus	Other Terminus	Owner(s)/Operator(s)	Note(s)
TAT-10 Transatlantic 10	Norden, Germany	Rhode Island, U.S.A	AT&T, Deutsche Telekom, Netherlands PTT	In-Service: 1992
AC1	Beverwyjk(Neth)	Brookhaven U.S.A.	Global Crossing	In-Service: May 1998
TAT-14 Transatlantic 14		New Jersey, U.S.A.	AT&T, BT, Sprint	Under Construction: October 2000

Endnote © format

There are more than 690 references saved in digital Endnote © format.

Bibliographic references have all been assumed to be quality rank 3-5 but have not been individually ranked. All published maps have been assigned a ranking of 5.

The notes field contains items a to d relating to the following legend:

- a. location of data (where is it archived/stored)
- b. cost (free or cost)

- c. quality ranks: 1. only to be used in extremis 2. quality uncertain or very patchy or of doubtful cost effectiveness to retrieve 3. fair to mediocre 4. data useable and free of charge or at cost 5. data should be used even if it incurs cost (eg. licensing or processing)
- d. latitude and longitude limits for maps

For the output in Endnote © without ranking see Appendix 4.

5. References

See Endnote compilation.

Appendix 1 Geological processes

The SEA7 project area comprises a major part of the coast, the continental shelf, the slopes and the deep-water basins occurring to the west of UK (Figure 2).

The coast and parts of the adjacent continental shelf were connected to the land by regional ice sheets during at least 4 regional glaciations at intervals between 10,000 years and more than approximately 800,000 years ago. The significance of the former regional glaciations to the geomorphology of SEA7 and to the composition of the seabed sediments is that they were associated with thick continental ice sheets that flowed and caused severe sub-ice sheet erosion of the underlying ground. The erosion formed the complex fjord-like shape of the coastline and transferred mainly non-biogenic gravelly and muddy sediments from the Scottish mainland and inner shelf to the seaward margins of the continental ice sheets. They also contributed to modern micro-seismicity and crustal warp originating from the post-glacial adjustments to removal of large volumes of sediment, changes of ice thickness, and sea-level rise. Although the modern SEA7 has very little non-biogenic sediment input it is in a boreal temperate climate zone set in a fertile sea with strong near-bottom currents on the continental shelf and slope. In these environments the modern sediments mainly consist of mixtures of reworked glacial sediments and post-glacial shell carbonates. Thus the remobilised sediments are muddy in locations that are sheltered from near-bottom currents, they are sandy in the more exposed locations and all the sediment size fractions typically contain >10% biogenic carbonate.

The seabed environments are strongly influenced by the geomorphology, sediment texture and sediment composition. In turn these are influenced by the origins and relative abundances of the seabed sediment sources and the amount of sediment reworking by biological processes and by geostrophic, tide, wind and storm-surge generated near-bottom currents.

Coastal geological processes are strongly influenced by the tidal range, which varies from 1- 4m with position on the coast. Coastal and nearshore processes are also strongly influenced by exposure to strong waves. For example, extreme-wave-orbital near-bottom current-strengths of more than 600cm/sec have been estimated for the 50-year storm in water depths of less than 100m at sites west of the Hebrides (eg. Pantin, 2001). Under these conditions seabed lithic gravel (>2mm diameter) is readily mobilised.

Further offshore, mean tidal, wave and geostrophically-generated near-bottom currents range from less than 50cm/sec to more than 200cm/sec, so that the exposed nearshore, continental shelf and upper slope seabed environments in SEA7 are characterised by mobile sands and silts (eg. Pantin, 1991). During the major part of modern times the near bottom currents on the middle to outer Hebrides Shelf at the shelfbreak are directed towards the northeast (e.g. Huthnance, 1986). Here, the current speeds vary on a regional scale with the shelf and slope geometry, that is, with the geological structure that underpins the changes to the seabed geomorphology.

At the mesoscale, the near-bottom currents and shelf and shelfedge seabed properties also vary with changes of seabed topography associated with rock platforms, basins, moraines and submarine fans, most of which have also been more or less shaped by glaciogene processes (e.g. Holmes and Stoker, 1990; Stoker *et al*, 1993; Stoker, 1995). At smaller scales, gravel ridges and intervening sand infill are associated with iceberg scour in a zone extending from the mid-shelf and over the shelfedge to approximately 450m water depth on the upper slope. Mobile sediment waves are found in environments ranging from the coast and continental shelf to the deepest-water basins and vary in size from the almost-ubiquitous sand ripples (<60cm wavelength) to the less common very large bedforms (>100m wavelength).

Outside of the areas affected by canyons and submarine landslides, the modern deep-basin sedimentary processes are dominated by erosion and deposition from contour-following currents (contouritic processes) and by deposition from the water column by slow settling and slow lateral advection (hemipelagic processes).

Regional slope angles at the margins of the Rockall Trough typically range from 1-4° and occasionally 7-14° on the upper slope. Relatively abrupt increases of slope angles and slope angles in the order of 2° or more commonly map to sediment-drift (contouritic) bedforms, rock crop at or near seabed in areas with very strong currents and to features formed by canyon and submarine landslide processes, these last occurring more frequently south of approximately 57°N (Armishaw *et al.*, 1998). Abrupt local changes in seabed gradients in the NE Rockall Basin are also related to isolated submarine landslides and are linked to evidence for sub-seabed fluid migration and seabed fluid expulsion (Baltzer *et al.*, 1998) in an area also noted for seabed mounds and pockmarks (Masson *et al.*, 2001). Modern submarine landslides occur south of approximately 56.5°N on the Barra and Donegal Fans. These map to relatively steep slope angles, prograding deposits associated with fan build-out towards deep-water and to proximity to the epicentres of modern seismicity (Holmes, 2002).

The deep-water areas of the western Rockall Trough and further west have been relatively sediment starved compared to those of the eastern Rockall Trough and there are relatively few submarine landslides. The seabed morphology of the unconsolidated sediments is predominantly shaped by along-slope sediment transport and hemipelagic sediment deposition and is characterised by sediment plastering or sediment drape over underlying structures.

Selected References

- Armishaw, J.E., Holmes, R. and Stow, D.A.V., 1998. Morphology and sedimentation on the Hebrides Slope and Barra Fan, NW UK continental margin. In: M.S. Stoker, D. Evans and A. Cramp (Editors), *Geological Processes on Continental Margins: Sedimentation, Mass-Wasting and Stability*. Geological Society, London, Special Publications, vol 129. Geological Society, London, pp. 81-104.
- Baltzer, A., Holmes, R. and Evans, D., 1998. Debris flows on the Sula Sgeir Fan, NW of Scotland. In: M.S. Stoker, D. Evans and A. Cramp (Editors), *Geological Processes on Continental Margins: Sedimentation, Mass-Wasting and Stability*. Geological Society, London, Special Publications, Vol. 29. Geological Society of London, London, pp. 105-115.
- Holmes, R. and Stoker, M.S., 1990. Pleistocene glaciomarine facies on the continental shelf and slope north-west of Scotland, Sediments 1990. 13th International Sedimentological Congress. Nottingham, England. 26th-31st August 1990. Abstract of Papers. International Association of Sedimentologists, Comparative Sedimentology Division. Utrecht., pp. 230.
- Holmes, R., 2002. Holocene shelf-margin submarine landslides, Donegal Fan, eastern Rockall Trough. In: J. Meinert and P.P.E. Weaver (Editors), *European Margin Sediment Dynamics: Sidescan Sonar and Seismic Images*. Springer-Verlag, New York.
- Huthnance, J.M., 1986. The Rockall slope current and shelf-edge processes. In: J. Mauchline (Editor), *The oceanography of the Rockall Channel*. Royal Society of Edinburgh, pp. 83-101.
- Masson, D., Bett, B., Billett, D. and Wheeler, A., 2001. The Darwin mounds - possible fluid escape features in the Northern Rockall Trough. *Journal of Conference Abstracts*, Southampton, 6: 751.
- Pantin, H.M., 1991. The Sea-bed sediments around the United Kingdom: Their bathymetric and Physical environment, grain size, mineral composition and associated bedforms. British Geological Survey Research Report SB/90/1.
- Stoker, M.S., Hitchen, K. and Graham, C.C. 1993. United Kingdom offshore regional report: the geology of the Hebrides and West Shetland shelves, and adjacent deep water areas. (London, HMSO for the British Geological Survey).

Stoker, M.S., 1995. The influence of Glacigenic Sedimentation on Slope-Apron development on the Continental Margin off NW Britain. In: R.S. Scrutton, M.S. Stoker, G.B. Shimmiel and A.W. Tudhope (Editors), Geological Society Special Publication. The Tectonics, Sedimentation and Palaeoceanography of the North Atlantic Region. Geological Society of London, London, pp. 159-177.

Appendix 2. Project liaison

Quentin Huggett,
Contracting Authority
Geotek Ltd
3 Faraday Close
Drayton Fields
Daventry
Northants. NN11 5RD
Quentin@Geotek.co.uk

Richard Holmes ¹ (BGS Edinburgh Compiler)
Jan Fraser ¹ (BGS Edinburgh Endnote © 6)
Colin Graham ¹ (BGS Edinburgh Arcview reports SEA7, geology)
Paul Henni, ¹ (BGS Edinburgh GMT from Oracle, EndNote SEA7, BGS Arcview)
Anne Richardson ¹ (BGS Edinburgh bibliography SEA7 in ACCESS)
Alan Stevenson ¹ (BGS Edinburgh European project network metadata data including rig decommissioning information)
John Ridgway ² (BGS Keyworth and liaison CEFAS Subcontract, SEA 6&8 bibliographies contamination seabed sediments and seawater overlap with seabed geology)

¹ British Geological Survey
West Mains Road
Edinburgh EH9 3LA
rho@bgs.ac.uk
Tel 0131 667 1000
Fax 0131 668 4140

² British Geological Survey
Keyworth,
Nottingham NG12 5GG

Steve Rowlett CEFAS (SEA 6,7,8 contamination seabed sediments and seawater, bibliographies overlap with seabed geology)
The Centre for Environment, Fisheries and Aquaculture Science (CEFAS)
Burnham Laboratory
Remembrance Avenue
Burnham-on-Crouch

Essex

CN0 8HA

S.M.Rowlatt@CEFAS.co.uk

Telephone 01621 78 72 00

Direct no 01621 78 72 44

Mr Colin Jacobs

Vikki Gunn (SOC assisting)

Challenger Division for Seafloor Processes

Southampton Oceanography Centre

European Way

Southampton

SO14 3ZH

Tel 023 8059 6550 (Gunn)

023 8059 6576 (Jacobs)

Fax 023 8059 6554

VKG@soc.soton.ac.uk

C.Jacobs@soc.soton.ac.uk

Professor Patrick M Shannon (assisting)

Vikram Unnithan (assisting)

University College Dublin

Dept of Geology

Bellfield

Dublin 4

Tel 00353 1706 77 77

Fax 00353 1283 77 33

P.shannon@ucd.ie

Vikram.Unnithan@ucd.ie

Appendix 3 Geographical Keywords

N.E. Atlantic Ocean	South Uist	Clyde
Rockall Trough	Barra	Ailsa Craig
Northern Rockall Trough	Barra Head	Bute
N E Rockall Basin	Malin Sea	Great Cumbrae
Maury Channel	North Minch	Little Cumbrae
Endymion Spur	South Minch	Inner Clyde, Clyde
Hatton Bank	The Little Minch	Enard Bay
Hatton-Rockall Basin	Sound of Raasay	Rubha Coiseach
Rockall Plateau	Inner Sound	Edrochillin Bay
Rockall	Raasay	Cape Wrath
George Bligh Bank	Rona	Solan Bank
Bill Bailey Bank	Skye	Glasgow
North Feni Ridge	Monach Island	Stanton Bank(s)
Feni Ridge	Rhum	Geikie Bulge
Rosemary Bank	Eigg	Darwin Mounds
Wyville Thomson Ridge	Muck	Flannan Trough
Anton Dohrn Seamount	Tiree	Geikie Escarpment
St Kilda	Coll	Larne
Hebrides Shelf	Mull	Kishorn
Hebridean Shelf	Staffa	Ronan Basin
Barra Fan	Treshnish Islands	Hatton Drift
Donegal Fan	Skerrymore	Iceland Basin
Sula Sgeir Fan	Blackstones Bank	Peach Slide
Sula Sgeir	Mull of Kintyre	Summer Islands
Malin Shelf	Rathlin Island	Priest Island
North Rona	North Channel	Finnan Islands
Butt of Lewis	Ailsa Craig	Western Island
Outer Hebrides	Firth of Clyde	Shiant Islands
Inner Hebrides	Beauforts Dyke	Loch Roag
Scottish Mainland	Luce Bay	W. Loch Tarbet
Lewis	Isle of Man	Pabbay
South Harris	Dundalk Bay	Monach Islands
Benbecula	Belfast	Sound of Barra
North Uist	Belfast Loch	Vatersay Sound

Loch Resort	Loch Carron	Ardrossan
The Narrows	Loch Torridon	Greenock
Loch Maddy	Loch Garloch	Gourock
Loch Uskavagh	Loch Etive	Gareloch
Loch Carman	Gruinard Bay	Rhu Marrows
Loch Skipport	Loch Broom	Faslane
Loch Eynort	Loch Kanaird	Loch Goil
Loch Boisdale	Loch Inver	Loch Striven
Castle Bay	Point of Stoer	Dunoon
Loch Watersay Sound	Eddrachillis Bay	Inchmarnock Water
Red Point	Loch Lauford	Loch Gilp
Berneray Sound	Loch Inchard	Loch Tarbet
Shillay Sound	Kilbrannan Sound	Gigha
Stornoway	Ailsa Craig	Sound of Gigha
Loch Grimashadur	Loch Ryan	Corryvreckan
Loch Luirbost	Stranraer	Loch Crinan
Sound of Harris	Mull of Galloway	Loch Etive
Village Bay	Burrow Head	Loch Crenan
Loch Bracadale	Peel	Loch Leven
Ardnamurchan Point	Isle of Man	Loch Eil
Tiree Passage	Port Erin	Ballachulish Bay
Loch Soridain	Calk Sound	Lismore Island
Jura Sound	Port St Mary	Linn of Morven
Firth of Lorne	Castletown Bay	Lynn of Lorne
Oban	Loch Carlingford	Loch Ailine
Loch Fyne	Strangford Loch	L. A'Choire
Loch Long	Donaghadee Sound	Loch Nevis
Loch Linnhe	Ardglass Harbour	Loch Tudth
Loch Faslane	Killough Harbour	Sound of Iona
Greenock	Larne Harbour	Treshnish
Dumbarton	The Maidens	Gott Bay
Sound of Mull	Portrush	Gunna Sound
Loch Sunart	Kilbrannon Sound	Loch Eathama
Sound of Sleat	Campbelton Loch	Mallaig
Loch Hourne	Sanda Island	Kyle Rhea
Loch Alsh	Holy Loch	Loch Euich

Loch Alsh
Kyleakin
Strome Narrows
Broadford Bay
Loch Kishorn
Caol Moire
Loch Sheldaig
Ayr Bay
Brodict Bay
Irvine Harbour
Troon Harbour
Ayr Harbour
Ullapool
Firth of Lorne

**End of geographical
keywords**

Appendix 4 Bibliography

Annotated references in Endnote© format are on compact disc.

The following references are output from the Endnote © format files:

Abrantes, F., Baas, J., Hafliðason, H., Rasmussen, T.L., Klitgaard, D., Loncaric, N. & Gaspar, L. 1998. Sediment fluxes along the north-eastern European Margin: inferring hydrological changes between 20 and 8 Ka. *Marine Geology*, **152**, 7-23.

(AFEN), A.F.E.N. 1998. UK Continental Shelf 17th Round Atlantic Margins Environmental Survey [CD-ROM]. *CD-ROM*, (Southampton).

(AFEN), A.F.E.N. 2000. Environmental Surveys of the Seafloor of the UK Atlantic Margin. *CD-ROM*, (Daventry).

Akhmetzanov, A.M., van Weering, T.C.E., Kenyon, N.H. & Ivanov, M.K. 1998. Carbonate mounds and reefs at the Rockall Trough and Porcupine margins. 69 in *Carbonate mud mounds and cold water reefs: deep-biosphere-geosphere coupling. TTR-7 post cruise conference*. Mol, B.D. (editor). (Gent, Belgium: University of Gent.)

Alcock, G.A. 1982. Offshore bottom pressure records. North Channel of Irish Sea 1979, Malin Shelf 1979, Rockall Trough 1979.

Allen, J.A. 1960. Manganese deposition on the shells of living molluscs. *Nature*, **185**, 336-337.

Allen, N.H. 1983. Recent temperate carbonate deposits on the continental shelf north and west of Scotland: distribution, sedimentology and reserves. Unpublished thesis, University of Strathclyde.

Andersen, M.S. 1988. Late Cretaceous and early Tertiary extension and volcanism around the Faeroe Islands. In: Morton, A.C. & Parson, L.M. (eds.). *Early Tertiary volcanism and the opening of the NE Atlantic*. **39**. Geological Society of London, London, 115-122.

Andersen, M.S. & Boldreel, L.O. 1995. Effect of Eocene-Miocene Compression Structures on the bottom-water currents in the Faeroe-Rockall area. In: Scrutton, R.S., Stoker, M.S., Shimmield, G.B. & Tudhope, A.W. (eds.). *Geological Society Engineering Geology Special Publication. Quaternary Engineering Geology*. **90**. 141-143.

Andersen, M.S., Nielsen, T., Sorensen, A.B., Boldreel, L.O. & Kuijpers, A. 2000. Cenozoic sediment distribution and tectonic movements in the Faroe region. In: Chalmers, J.A. & Cloetingh, S. (eds.). *Global and Planetary Change. Neogene uplift and tectonics around the North Atlantic*. **24**; **3-4**. Elsevier, Amsterdam, Netherlands, 239-259.

Anderson, S.J. & Bracey, J.S. 1999. Environmental management of deep-water exploration, Rockall Trough. 17-19 in AAPG, Tulsa, Oklahoma, USA.)

Anderton, R., Bridges, P.H., Leeder, M.R. & Sellwood, B.N. 1979. *A dynamic stratigraphy of the British Isles: a study in crustal evolution*. George Allen & Unwin, London.

Andrews, J.T., Austin, W.E.N., Bergsten, H. & Jennings, A.E. 1996. The Late Quaternary palaeoceanography of North Atlantic margins: an introduction. *In: Andrews, J.T., Austin, W.E.N., Bergsten, H. & Jennings, A.E. (eds.). Geological Society Special Publication, 111. Late Quaternary Palaeoceanography of the North Atlantic Margins*. Geological Society, London, 1-6.

Anon. 1982. Current measurements and hydrography in the Rockall Trough and Faroe-Shetland channel: RRS Frederick Russell Cruise 82/2, March 16-March 29 1982. *IOS Cruise Rep [Godalming]*, **127**.

Anon. 1992. New survey of the UK Rockall continental shelf. *Geographical Journal*, **158**, Pt 2, 250.

Ardus, D.A. & Chesher, J.A. 1977. RRS Challenger Cruise - Leg 2, 30 September - 14 October 77/CH/02.

Ardus, D.A. & Harrison, D.J. 1990. The assessment of aggregate resources from the UK continental shelf. *In: Ardus, D.A. & Champ, M.A. (eds.). Ocean Resources, Vol. 1, Assessment and Utilisation. 1*. Kluwer, Dordrecht, 113-128.

Armishaw, J.E., Holmes, R. & Stow, D.A.V. 1996. The Barra Fan: a mixed contourite/turbidite sediment body on the West Hebridean margin. 66 in *Applied Geoscience conference Warwick University 15-18 April 1996*. Geological Society, London.)

Armishaw, J.E., Holmes, R. & Stow, D.A.V. 1998. Morphology and sedimentation on the Hebrides Slope and Barra Fan, NW UK continental margin. *In: Stoker, M.S., Evans, D. & Cramp, A. (eds.). Geological Processes on Continental Margins: Sedimentation, Mass-Wasting and Stability. Geological Society, London, Special Publications, vol 129*. Geological Society, London, 81-104.

Armishaw, J.E. 1999. Bottom current accumulation and sediment fluxes on the Hebridean Slope. Unpublished thesis, University of Southampton.

Armour, A.R. & Bott, M.H.P. 1977. The Hebridean Margin Seismic Project of 1975. *The Geophysical Journal of the Royal Astronomical Society*, **49**, Pt 1, 284.

Austin, W.E.N. 1991. Late Quaternary Benthonic Foraminiferal Stratigraphy of the Western UK Continental Shelf. Unpublished PhD. thesis, University of Cardiff.

Austin, W.E.N. 1992. Late Quaternary Benthonic Foraminiferal Stratigraphy of the Western UK Continental Shelf. *Quaternary Newsletter*, **68**, 18-19.

Austin, W.E.N., Kroon, D., Shimmiel, G.B. & Derrick, S. 1994. High resolution marine records of the last glacial/interglacial transition. (Abstract). *Mineralogical Magazine*, **58A**, 31.

Austin, W.E.N., Kroon, D. & Stoker, M.S. 1996. Piston coring the Hebridean margin - an initiative within the Images programme. NERC Small Research Grant: GR/01595'A'.

Austin, W.E.N., Derrick, D., Kroon, D., Shimmield, G.B. & Thompson, R. 1996. Palaeoceanographic records from the continental slope N.W. Scotland. 64 in *Applied Geoscience conference Warwick University 15-18 April 1996*. Geological Society, London.)

Austin, W.E.N. 1996. Quantifying down-slope transportation: stable isotope records from benthic foraminifera. 68 in *Applied Geoscience conference Warwick University 15-18 April 1996*. Geological Society, London.)

Austin, W.E.N. & Kroon, D. 1996. The Lateglacial palaeoceanographic evolution of the Hebridean Continental Shelf, N.W. Scotland. In: Andrews, J.T., Austin, W.E.N. & Bergsten, H.E. (eds.). *The Lateglacial Palaeoceanography of the North Atlantic Margins. Geological Society Special Publication 111*. Geological Society of London, London, 187-214.

Austin, W.E.N. 1996. North East Atlantic benthic foraminifera: Modern distribution patterns and palaeoecological significance. In: Andrews, J.T., Austin, W.E.N., Bergsten, H. & Jennings, A.E. (eds.). *Geological Society Special Publication, 111. Late Quaternary Palaeoceanography of the North Atlantic Margins*. Geological Society, London, 679-691.

Austin, W.E.N. & Kroon, D. 1998. North Atlantic thermohaline circulation during the last 15000 years: testing the 'Superconveyor' Theory. 109 in *GeoScience 98, (Keele University, 14-18 April 1998) Abstracts Volume*. Geological Society of London.

Austin, W.E.N. & Evans, J.R. 2000. Benthic foraminifera and sediment grain size variability at intermediate water depths in the Northeast Atlantic during the late Pliocene - early Pleistocene. *Marine Geology*, **170**, 423-441.

Austin, B.J. 2000. Rockall Trough Shallow Stratigraphy and Geohazards Study, RSG Project No: 98/23.

Austin, W.E.N. & Kroon, D. 2001. Deep sea ventilation of the northeastern Atlantic during the last 15,000 years. *Global and Planetary Change*, **30**, 13-31.

Backstrom, D.L. & Kuijpers, A. 1997. A late Quaternary paleoceanographic record (core ENAM-30) from near the Faeroe Bank Channel outlet. *1st ENAM II Workshop*. Geomar.

Backstrom, D.L. 2000. *Late Quaternary paleoceanography and paleoclimate of the North Atlantic Ocean*. (Late Quaternary paleoceanography and paleoclimate of the North Atlantic Ocean). Goteborg University, Earth Sciences Centre. Goteborg, Sweden.

Bacon, M. 1972. Cruise report for surveys in Moray Firth and Sea of Hebrides. MGU project no 72/05. *Marine Geophysics Report No. 35*.

Bailey, R.J., Grzywacz, J.M. & Buckley, J.S. 1974. Seismic Reflection Profiles of the Continental Margin bordering the Rockall Trough. *Journal of the Geological Society of London*, **130**, 55-69.

Bailey, R.J. & Haynes, J.R. 1974. New dredge samples from the continental margin bordering Rockall Trough. *Marine Geology*, **16**, Pt 3, M57-M62.

Ballantyne, C.K., McCarroll, D., Nesje, A., Dahl, S.O. & Stone, J.O. 1998. The last ice sheet in North-West Scotland: reconstruction and implications. *Quaternary Science Reviews*, **17**, 1149-1184.

Ballantyne, C.K., McCarroll, D., Nesje, A., Dahl, S.O., Stone, J.O. & Fifield, L.K. 1998. High-resolution reconstruction of the last ice sheet in NW Scotland. *Terra Nova*, **10**, 63-67.

Ballantyne, C.K. & Hallam, G.E. 2001. Maximum altitude of late Devensian glaciation on South Uist, Outer Hebrides, Scotland. *Proceedings of the Geologists' Association*, **112**, 155-167.

Baltzer, A., Evans, D. & Long, D. 1996. Differentiation of the origin of debris flows on the West Shetland slope. *Applied Geoscience conference Warwick University 15-18 April 1996*. Geological Society, London.

Baltzer, A., Howe, J.A. & Stoker, M.S. 1997. Paleo current formed sediment waves on the Geikie Edge, NW of UK margin. *Terra Nova, Volume 9, Abstract Supplement 1*, **9**, 262.

Baltzer, A. 1997. Sediment waves on the Geikie edge, NW of UK Margin. *1st ENAM II Workshop*. Geomar.

Baltzer, A., Holmes, R. & Evans, D. 1998. Debris flows on the Sula Sgeir Fan, NW of Scotland. *In: Stoker, M.S., Evans, D. & Cramp, A. (eds.). Geological Processes on Continental Margins: Sedimentation, Mass-Wasting and Stability. Geological Society, London, Special Publications, Vol. 29. Geological Society of London, London, 105-115.*

Baltzer, A., Walter-Simmonet, A. V., Evans, D. and Long, D. 1999. Clay mineral variations in glacial versus classic debris flows on the continental slope west and north of Scotland. 16 in *North East Atlantic Slope Processes: Multi-Disciplinary Approaches, Abstract Book. Conference 24-27th January 1999*. Friend, P.a.K., N. (editor). Southampton Oceanography Centre, Southampton, UK.

Banaszek, A.D. & MacDonald, D.C.C. 1985. UK Continental slope Experiment (CONSLEX) 1982/83 Offshore bottom pressure records.

Banner, F.T. 1979. *Seabed resources, potential and actual (excluding hydrocarbons)*. The North-West European shelf seas: the sea bed and the sea in motion - II. Physical and Chemical Oceanography, and Physical Resources. Elsevier, Amsterdam.

Banner, F.T. & Culver, S.J. 1979. Sediments of the North-Western European Shelf. *In: Banner, F.T., Collins, M.B. & Massie, K.S. (eds.). The North-West European shelf seas: the sea bed and the sea in motion - I. Geology and sedimentology*. Elsevier, Amsterdam, 271-300.

Barber, P.L., Dobson, M.R. & Whittington, R.J. 1979. The geology of the Firth of Lorne as determined by seismic and dive sampling methods. *Scottish Journal of Geology*, **15**, 217-230.

Barton, A.J. & White, R.S. 1997. Volcanism on the Rockall continental margin. *Journal of the Geological Society of London*, **154**, Pt 3, 531-536.

Belderson, R.H., Kenyon, N.H. & Stride, A.H. 1970. Holocene sediments on the continental shelf west of the British Isles. pages 157-170.

Benn, D.I. 1997. Glacier fluctuations in western Scotland. *Quaternary International*, **38**, 137-147.

Bentley, P.A.D. 1986. Geophysical studies in southern and central Rockall Trough, northeast Atlantic. Unpublished thesis, University of Edinburgh.

Bentley, P.A.D. & Scutton, R.A. 1987. Seismic investigations into the basement structure of southern Rockall Trough. *In: Brooks, J. & Glennie, K.W. (eds.). Petroleum Geology of North West Europe*. Graham & Trotman, London, 667-675.

Berggren, W.A. & Schnitker, D. 1983. Cenozoic marine environments in the North Atlantic and Norwegian-Greenland Sea. *In: Bott, M.H.P., Saxov, S., Talwani, M. & Thiede, J. (eds.). Structure and development of the Greenland-Scotland Ridge; new methods and concepts*. Plenum, New York and London, 495-548.

Berthois, L. & Roa, P. 1967. Interprétation des analyses de minéraux lourds par pourcentages pondérés, dans la région des Feroër-Écosse-Hébrides. *Compte Rendu Academie des Sciences Serie D*, **265**, 195-197.

Berthois, L. & Buit, M.H.D. 1971. Contribution à l'étude de la sédimentation en l'Atlantique nord: zone comprise entre les Archipels Faerøe - Shetland - Orcades - Hébrides. *Bulletin de l'Institut du Geologie du Bassin d'Aquitaine*, **11**, 365-374.

Berthois, L., Latouche, C. & Parra, M. 1973. Etude minéralogique et géochimique de quelques sédiments de la zone comprise entre les Archipels: Faerøe, Shetland, Orcades, et Hébrides. Contribution à la connaissance de l'origine des sédiments dans le domaine profond de l'Atlantique Nord-oriental. *Bulletin de l'Institut du Geologie du Bassin d'Aquitaine*, **14**, 3-17.

Bertram, C.J., Elderfield, H., Shackleton, N.J. & Macdonald, J.A. 1995. Cadmium/calcium and carbon isotope reconstructions of the glacial northeast Atlantic Ocean. *Palaeoceanography*, **10**, 563-578.

Bett, B.J. 1997. RRS Charles Darwin Cruise 101C Leg 2, 14 Jul-20 Aug 1996, Atlantic Margin Environmental Survey: seabed survey of the shelf edge and slope west of Shetland.

Bett, B.J. 1998. The Biological Environment – the Atlantic Margin Environmental Surveys. *Atlantic Frontier Environmental Conference 6th & 7th October 1998 (Abstracts of Presentations)*. Atlantic Frontier Environmental Forum, A. (editor).

Bett, B. 1998. Observations on the seabed mound field centred at approximately 59.50N 007.30W. Report to CONOCO UK and Atlantic Frontiers Environmental Network, 15 pages. *Unpublished manuscript*.

Bett, B.J. 1999. RRS Charles Darwin Cruise 112C, 19 May-24 Jun 1998. Atlantic margin environmental survey: seabed survey of deep-water areas (17th round tranches) to the north and west of Scotland. *Cruise Report of the Southampton Oceanography Centre*, **Issue 25**, 1-171.

Bett, B.J., Pearson, T.H. & Sørensen, J. 2000. Expert review of seabed fauna and chemistry. pages 58.

Bett, B.J. & Axelsson, M. 2000. Section 4.2: Survey Photography. *In*: Network, A.F.E. (ed.). *Environmental Surveys of the Seafloor of the UK Atlantic Margin [CD-ROM]*. Geotek Ltd, Daventry.

Bett, B.J. 2001. UK Atlantic Margin Environmental Survey: Introduction and overview of bathyal benthic ecology. *Continental Shelf Research*, **21**, 917-956.

Bett, B.J. 2001. RRS Charles Darwin cruise 119C. Initial assessment of seabed observations made during the White Zone (WhiZ) Environmental Survey: Seabed survey of the deep-waters to the north and west of Shetland.

BGS. 1985. Isle of Man 54N 06W. Seabed sediment and Quaternary geology 1:250 000. Ordnance Survey.

BGS. 1986. Rona 59N 06W. Seabed sediments 1: 250 000. Ordnance Survey.

BGS. 1986. Malin 55N 08W. Seabed sediments 1: 250 000. Ordnance Survey.

BGS. 1986. Clyde 55N 06W. Seabed sediment and Quaternary geology 1: 250 000. Ordnance Survey.

BGS. 1988. Little Minch (Including part 57N 06W Great Glen) 57N 08W. Seabed sediment and Quaternary geology 1: 250 000. Ordnance Survey.

BGS. 1988. Tiree (including part Argyll) 56N 08W. Seabed sediments and Quaternary geology 1: 250 000. Ordnance Survey.

BGS. 1988. Sula Sgeir, 59N 08W. Seabed sediments 1: 250 000. Ordnance Survey.

BGS. 1989. Sub-Pleistocene geology of the British Isles and adjacent continental shelf 1:1 000 000.

BGS. 1989. Sutherland 58N 06W. Seabed sediments 1: 250 000. Ordnance Survey.

BGS. 1989. Geikie, 58N 10W. Seabed sediments 1: 250 000. Ordnance Survey.

BGS. 1989. Lewis, 58N 8W. Seabed sediments 1: 250 000. Ordnance Survey.

BGS. 1990. Peach 56N 10W. Seabed sediments 1: 250 000. Ordnance Survey.

BGS. 1991. Geology of the United Kingdom, Ireland and adjacent continental shelf: North sheet. British Geological Survey 1:1 000 000.

BGS. 1991. St Kilda 57N 10W. Seabed sediments 1: 250 000. Ordnance Survey.

BGS. 1994. East Irish Sea (including Isle of Man and Lake District) 54N 06W solid geology. Solid geology 1:250 000. Ordnance Survey.

Binns, P.E. 1972. IGS Continental Shelf Unit II, The Sea of The Hebrides, Cruise Report m.v. Whitehorn 7 Apr/8 May 1972 Cruise CSU II 72/W2.

Binns, P.E. 1972. Helen's Reef, Rockall: Field report on joint IGS/NIO diving survey from RFA Engadine, 16-30 June 1972.

Binns, P.E. 1972. The Sea of the Hebrides, MV Whitehorn, 7th April - 8th May 1972: Cruise CSU II 72/W2. *Marine Geology Internal Report*.

Binns, P.E., McQuillan, R. & Kenolty, N. 1974. The geology of the Sea of the Hebrides.

Binns, P.E. & Fannin, N.G.T. 1975. Cruise Report m.v. Whitehorn 20 June to 3 July 1973 Cruise CSU II 73/WH/13.

Biscaye, P.E. 1965. Mineralogy and Sedimentation of Recent Deep-Sea Clay in the Atlantic Ocean and Adjacent Seas and Oceans. *Geological Society of America Bulletin*, **76**, 803-832.

Bishop, P. 1977. Glacial and post-glacial sedimentation in the Minches, North West Scotland. Unpublished thesis, University of London.

Bishop, P. & Jones, E.J.W. 1979. Patterns of glacial and post-glacial sedimentation in the Minches, North-West Scotland. *In*: Banner, F.T., Collins, M.B. & Massie, K.S. (eds.). *The North-West European shelf seas: the sea bed and the sea in motion - 1. Geology and sedimentology*. Elsevier, Amsterdam, 89-94.

Björck, S., Kromer, B., Johnsen, S., Bennike, O., Hammarlund, D., Lemdahl, G., Possnert, G., Rasmussen, T.L., Wohlfarth, B., Hammer, C.U. & Spurk, M. 1996. Synchronized terrestrial-atmospheric deglacial records around the North Atlantic. *Science*, **274**, 1155-1160.

Bjorklund, K.R. & Ciesielski, P.F. 1994. Ecology, morphology, stratigraphy, and the paleoceanographic significance of *Cycladophora davisiana davisiana*; Part I, Ecology and morphology. *Marine Micropaleontology*, **24**, 71-88.

Blakiston Houston, L.M.L. 1993. Distribution and dynamics of nutrients on the Scottish NW Shelf Region. Unpublished thesis, University of Belfast.

Boldreel, L.O. & Andersen, M.S. 1993. Late Paleocene to Miocene compression in the Faeroe-Rockall area. 1025-1034 in *Proceedings of the 4th conference on Petroleum Geology in Northwest Europe, 1992*. Geological Society of London.)

Boldreel, L.O. & Andersen, M.S. 1995. The relationship between the distribution of Tertiary sediments and Tectonic processes and deep water circulation around the Faeroe Islands. *In*: Scrutton, R.S., Stoker, M.S., Shimmield, G.B. & Tudhope, A.W. (eds.). *The Tectonics, Sedimentation and Palaeoceanography of the North Atlantic Region*. Geological Society of London Special Publication. 90.145-158.

- Boldreel, L.O. & Anderson, M.S. 1995. The relationship between the distribution of Tertiary sediments, tectonic processes and deep-water circulation around the Faroe Islands. *In: Scrutton, R., Stoker, M.S., Shimmield, D.B. & Tudhope, A.W. (eds.). The tectonics, sedimentation and palaeoceanography of the North Atlantic region.* (Geological Society of London Special Publication). **90**. Geological Society of London, London.
- Boldreel, L.O. & Andersen, M.S. 1998. Paleogene and Neogene vertical tectonic movement on the Faeroe-Rockall Plateau. 81-82 in *Neogene Uplift and Tectonics around the North Atlantic - International workshop, Copenhagen, May 18-19 1998*. Boldreel, L.a.J. P. (editor). Geological Survey of Denmark and Greenland (GEUS).
- Bond, G., Showers, W., Cheseby, M., Lotti, R., Almasi, P., de Menocal, P., Priore, P., Cullen, H., Hajdas, I. & Bonani, I. 1997. A pervasive millennial-scale cycle in North Atlantic Holocene and Glacial climates. *Science*, **278**, 1257-1266.
- Booth, D.A. & Ellet, D.J. 1975. The Scottish continental slope current. *Continental Shelf Research*, **2**, 127-146.
- Borenas, K.M. & Lundberg, P.A. 1988. On The Deep-Water Flow Through The Faroe Bank Channel. *Journal Of Geophysical Research*, **93**, 1281-1292.
- Borthwick, A. 1995. Glacial Geology and Sedimentation of the North Lewis Basin. *QRA Annual Discussion Meeting, 1995 Abstract Volume The Lateglacial Palaeoceanography of the North Atlantic Margins*. Andrews, J.T., Austin, W.E.N. & Bergsten, H.E. (editors).
- Bott, M.H.P. & Watts, A.B. 1970. Deep sedimentary basins proved in the Shetland-Hebridean continental shelf and margin. *Nature*, **227**, 265-268.
- Bott, M.H.P. 1978. The origin and development of the continental margins between the British Isles and southeastern Greenland. *In: Bowes, D.R. & Leake, B.E. (eds.). Crustal evolution in northwestern Britain and adjacent regions.* (Geological Society of London Special Publications). **10**. Geological Society of London, London, 377-392.
- Bott, M.H.P., Saxov, S., Talwani, M. & Thiede, J. 1983. *Structure and development of the Greenland-Scotland Ridge: New Methods and Concepts*. Plenum Press, New York and London.
- Boulton, G.S., Chroston, N.P. & Jarvis, J. 1981. A marine seismic study of late Quaternary sedimentation and inferred glacier fluctuations along western Inverness-shire, Scotland. *Boreas*, **10**, 39-51.
- Boulton, G.S., Peacock, J.D. & Sunderland, D.G. 1991. Quaternary. *In: Craig, G.T. (ed.). Geology of Scotland*. Geological Society of London Publishing House, Bath, 503-543.
- Bout Roumazeilles, V., Vanderaveroet, P., Recourt, P. & Davies, G. 2001. Mineralogical and geochemical records of glacial-interglacial cycles in the North Atlantic Ocean over the last million years. *Journal of Conference Abstracts*, **6**, 126.
- Bowden, R. & Orford, J.D. 1984. *Residual sediment cells on the morphologically irregular coastline of the Ards Peninsula, Northern Ireland*. (Proceedings of the Royal

Irish Academy, Section B: Biological, Geological and Chemical Science). Royal Irish Academy, Dublin, Ireland.

Bowen, D.Q. 1991. Time and space in glacial sediment systems of the British Isles. *In*: Elders, J., Gibbard, P.L. & Rose, J. (eds.). *Glacial Deposits of Great Britain and Ireland*. A A Balkema, Rotterdam, 3-11.

Bowen, D.Q., Phillips, F.M., McCabe, A.M., Knutz, P.C. & Sykes, G.A. 2002. New data for the Last Glacial Maximum in Great Britain and Ireland. *Quaternary Science Reviews*, **21**, 89-101.

Britsurvey. 1995. Shell UK Exploration and Production: North Atlantic surveys - North Feni Ridge, Rockall Trough and Maury Channel. Volume 3: Oceanographic, water and sediment sample analyses.

Britsurvey. 1995. Shell UK Exploration and Production: North Atlantic surveys - North Feni Ridge. Volume 2: Detailed report.

Brodie, J. & White, N. 1994. Sedimentary basin inversion caused by igneous underplating: Northwest European Continental Shelf. *Geology*, **22**, 147-150.

Brooks, J. & Glennie, K.W. 1987. *Petroleum geology of North West Europe*. Graham and Trotman, London.

Browne, M.A.E. & McMillan, A.A. 1984. Shoreline inheritance and coastal history in the Firth of Clyde. *Scottish Journal of Geology*, **20**, 119-120.

Buchanan, J.Y. 1981. On the composition of oceanic and littoral manganese nodules. *Transactions of the Royal Society of Edinburgh*, **36**, 459-483.

Buckley, J.S. & Bailey, R.J. 1975. Geophysical evidence on the nature of the Hebrides Terrace Seamount. *Scottish Journal of Geology*, **11**, 37-45.

Bulat, J. 1991. The Rockall continental margin: A review of existing seismic data. *BGS marine Geology Technical Report*.

Bulat, J. & Long, D. 1998. Creation of seabed feature maps from 3D seismic horizon data sets.

Bulat, J. & Long, D. 2000. Use of 3D exploration surveys for regional images of the seabed. *Geoscience 2000, Manchester*.

Bulat, J. & Long, D. 2000. Use of commercial 3D seismic in imaging sedimentary processes west of Britain. *Seismic expression of contourites and related deposits: a seismic workshop, Trieste*, 36-37.

Burrows, M., Thorpe, S.A. & Meldrum, D.T. 1999. Dispersion over the Hebridean and Shetland shelves and slopes. *Continental Shelf Research*, **19**, 49-55.

Burrows, M. & Thorpe, S.A. 1999. Drifter observations of the Hebrides slope current and nearby circulation patterns. *Annales Geophysicae-Atmospheres Hydrospheres And Space Sciences*, **17**, 280-302.

- Burton, J.D. & Young, M.L. 1979. Trace metals in the shelf seas of the British Isles. *In: Banner, F.T., Collins, M.B. & Massie, K.S. (eds.). The North-West European shelf seas: the sea bed and the sea in motion - II. Physical and chemical oceanography, and physical resources.* Elsevier, Amsterdam, 494-516.
- Butler, C.A. 1995. Basement fault reactivation: the kinematic evolution of the Outer Hebrides Fault Zone, Scotland. Unpublished thesis, University of Durham.
- Calvert, S.E. & Price, N.B. 1970. Composition of manganese nodules and manganese carbonates from Loch Fyne, Scotland. *Mineralogy and Petrology*, **29**, 215-233.
- Cameron, T.D.J. & Holmes, R. 1999. The continental shelf. *In: Bowen, D.Q. (ed.). A revised correlation of Quaternary deposits in the British Isles.* (Geological Society of London Special Publications). **23**. Geological Society of London, London, 125-139.
- Carrie, R.H., Mitchell, L. & Black, K.D. 1998. Fatty acids in surface sediment at the Hebridean shelf edge, west of Scotland. 1583-1593 in *Advances in organic geochemistry 1997; proceedings of the 18th international meeting on Organic geochemistry; Part II, Biogeochemistry.* Horsfield, B., Radke, M., Schaefer, R. & Wilkes, H. (editors). Pergamon. Oxford-New York, International.)
- Carter, R.W.G., Devoy, R.J.N. & Shaw, J. 1989. Late Holocene sea levels in Ireland. *In: Late Quaternary sea-level changes and crustal movements in the British Isles. Journal of Quaternary Science.* **4**. John Wiley and Sons for the Quaternary Research Association, Chichester, 7-24.
- Cartwright, D.E. 1969. Extraordinary tidal currents near St. Kilda. *Nature*, **223**, 928-932.
- Castillo, P.F. 1999. Sedimentation of organic matter on the Hebridean Shelf. Unpublished PhD thesis, University of Wales.
- Caston, G.F. 1975. Igneous dykes and associated scour hollows of the North Channel, Irish Sea. *Marine Geology*, **18**, M77-M85.
- Caston, G.F. 1976. The floor of the North Channel, Irish sea: a side-scan sonar survey.
- Cater, D.J.T., Loynes, J. & Challenor, P.G. 1987. Estimates of extreme current speeds over the continental slope off Scotland.
- Chapman, T.J., Broks, T.M., Corcoran, C.V., Duncan, L.A. & Dancer, P.N. 1999. The structural evolution of the Erris Trough, offshore northwest Ireland, with implications for hydrocarbon generation. *In: Fleet, A.J. & Boldy, S.A.R. (eds.). Petroleum Geology of Northwest Europe: Proceedings of the 5th Conference.* Geological Society, London, 455-469.
- Cherriman, J.W. & Gould, W.J. 1982. RRS 'Frederick Russell' Cruise 82/2, March 16 - March 29, 1982. Current Measurements and Hydrography in the Rockall Trough and Faeroe-Shetland Channel.
- Chesher, J.A. & Evans, D. 1976. Offshore Geology of the Western Isles and its Economic Significance.

Chesher, J.A., Smythe, D.K. & Bishop, P. 1983. The geology of The Minches, Inner Sound and Sound of Raasay. *IGS Report*.

Chesher, J.A. 1985. British Magnus Cruise 85/09, Leg 1, 5th - 28th August 1985: St Kilda sheet, NW Atlantic Margin. Cruise Report. *Marine Geology Internal Report*.

Chesher, J.A. & Smythe, D.K. 1986. Little Minch, Sheet 57N - 08W, 1:250 000 Map Series, Solid Geology. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.

Chesher, J.A. 1987. Cruise Report of Magnus Heinason, 17 September - 4th October 1987. NE Atlantic.

Chesher, J.A. 1992. The Barra Fan Cruise report. *BGS Technical Report*.

Chi, J. & Mienert, J. 1996. Linking physical property records of Quaternary sediments to Heinrich events. *Marine Geology*, **131**, 57-73.

Chuter, A. 1979. Study to see if 'north-west frontier' oil is commercial [Rockall Trough]. *Engineer [London]*, **249/6432**, 12.

Clift, P.D. & Turner, J. 1998. Paleogene igneous underplating and subsidence anomalies in the Rockall-Faeroe-Shetland area. *Marine and Petroleum Geology*, **15**, 223-243.

Clift, P.D., Charter, A. & Hurford, A.J. 1998. The erosional and uplift history of NE Atlantic passive margins: constraints on a passing plume. *Journal of the Geological Society of London*, **155**, 787-800.

CLIMAP. 1976. The Surface of the Ice-Age Earth. Science. *Science, New York*, **191**, 1131-1137.

Cloetingh, S., Gradstein, F.M., Kooi, H., Grant, A.C. & Kaminski, M. 1990. Plate reorganization: a cause of rapid late Neogene subsidence and sedimentation around the North Atlantic? *Journal of the Geological Society of London*, **147**, 495-506.

Cockburn, A.M. 1935. The geology of St Kilda. *Transactions of the Royal Society of Edinburgh*, **58**, 511-547.

Cole, G.A.J. & Crook, T. 1910. On the rock specimens dredged from the floor of the Atlantic off the coast of Ireland.

Condensed Report. 1995. Survey of Coastal Erosion in the Western Isles.

Corfield, S., Murphy, N. & Parker, S. 1999. The structural and stratigraphic framework of the Irish Rockall Trough. In: Fleet, A.J. & Boldy, S.A.R. (eds.). *Petroleum Geology of Northwest Europe: Proceedings of the 5th Conference*. 407-420.

Cremer, M., Faugeres, J.M. & Poutiers, J. 1982. Les glacis continentaux de marge stable; morphologie et dynamique sedimentaire, evolution au Quaternaire recent
Translated Title: The continental slope of a stable margin; morphology and sedimentary

dynamics; evolution during the upper Quaternary. *In: Bulletin de la Societe Geologique de France. Oceans-paleoceans; Seance specialisee de la Societe geologique de France. Translated Title: Oceans--paleo-oceans; special session of the Geological Society of France.* **24**. Societe Geologique de France. Paris, France, 433-446.

Cremer, M., Faugeres, J.C., Grousset, F. & Gonthier, E. 1993. Late Quaternary sediment flux on sedimentary drifts in the Northeast Atlantic. *Sedimentary Geology*, **82**, 89-101.

Crofts, R. & Ritchie, W. 1973. Beaches of Mainland Argyll.

Cronan, D.S. 1969. Recent sedimentation in the central NE Irish Sea. *IGS Report*.

Cronan, D.A. 1970. Geochemistry of Recent sediments from the central NE Irish Sea. *IGS Report*.

Cunningham, G.A. & Shannon, P.M. 1997. The Erris Ridge; a major geological feature in the NW Irish offshore basins. *Journal of the Geological Society of London*, **154**, Pt 3, 503-508.

Curry, D. 1978. A Correlation of Tertiary rocks in the British Isles.

Curry, G.B. 1983. Ecology of the Recent deep-water rhynchonellid brachiopod *Cryptopora* from the Rockall Trough. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **44**, 93-102.

Davies, A.J. 1974. Geophysical surveys in the Celtic Sea area, Project 74/6, MV Briarthorn, April - June 1974. *Marine Geophysics Unit Report*.

Davies, H.C., Dobson, M.R. & Whittington, R.J. 1984. A revised seismic stratigraphy for Quaternary deposits on the inner continental shelf west of Scotland between 55°30'N and 57°16'30'N. *Boreas*, **13**, 49-66.

Davies, G.J. 1987. Aspects of the biology and ecology of deep-sea Scaphopoda (Mollusca). 270.

Davies, T.A., Bell, T., Cooper, A.K., Josenhans, H., Polyak, L., Solheim, A., Stoker, M.S. & Stravers, J.A. 1997. *Glaciated Continental Margins - An atlas of acoustic images*. Chapman & Hall, London.

Davies, R., Cartwright, J. & Rana, J. 1999. Giant hummocks in deep-water marine sediments; evidence for large-scale differential compaction and density inversion during early burial. *Geology, Boulder*, **27**, 907-910.

Davies, R., Cartwright, J., Pike, J. and Line, C. 2001. Early Oligocene initiation of North Atlantic deep water formation. *Nature.*, **410**, 917-920.

Dawson, A.G. 1980. Late Quaternary sea levels and crustal deformation, Scotland. *In: International Union for Quaternary Research. Subcommission on shorelines for NW Europe, Symposium*. Polytechnic, Coventry.

- Dawson, D.A.G. 1980. Shore erosion by frost; an example from the Scottish Lateglacial. *In: Lowe, J., Gray, J.M. & Robinson, J.E. (eds.). Studies in the Lateglacial of North-West Europe.* Pergamon Press, Oxford, 45-53.
- Dawson, A.G., Dawson, S., Mighall, T.M., Waldmann, G. & Mactaggart, F. 2001. Intertidal peat deposits and early Holocene relative sea-level changes, Traigh Eileraig, Isle of Coll, Scottish Hebrides. *Scottish Journal of Geology*, **37 Issue 1**, 11-18.
- Day, G.A. & Sunderland, J. 1970. Cruise report for MV Surveyor, 29th August - 2nd September, and 14th - 17th October, 1970: North Channel survey, MGU project 70/04. *Marine Geophysics Report No. 13.*
- de Haas, H., T C E van Weering & Stoker, M.S. in press. Development of a sediment drift: Feni Drift, northeast Atlantic margin. *In: Meinert, J. & Weaver, P.P.E. (eds.). European Margin Sediment Dynamics: Sidescan Sonar and Seismic Images.* Springer-Verlag, New York.
- Deegan, C.E. 1972. Firth of Clyde, MV Whitehorn 27 February - 5th April 1972: Cruise CSU II 72/W1. *Marine Geology Internal Reports.*
- Deegan, C.E. 1973. Cruise report MV Whitehorn Cruise 73/WH/01, 27th February - 4th April 1973: The Firth of Clyde and Malin Sea.
- Deegan, C.E., Kirby, R., Rae, I. & Floyd, R. 1973. The superficial deposits of the Firth of Clyde and its sea lochs. *IGS Report*, pages 42.
- Deegan, C.E. 1978. Boreholes and outcrop sampling in sheet 55°N/6°W. pages 65-76.
- Department of Energy. 1985. Environmental parameters on the United Kingdom Continental Shelf.
- Dickson, R.R., Gould, W.J., Müller, T.J. & Maillard, C.D. 1985. Estimates of the mean circulation in the deep layer (> 2000m) of the eastern North Atlantic. *Oceanology*, **14**, 103-127.
- Dietrich, V.J. & Jones, E.J.W. 1980. Volcanic rocks from Rosemary Bank (Rockall Trough, NE Atlantic). *Marine Geology*, **35**, 287-297.
- Dobinson, A. 1978. Geophysical studies south and west of Kintyre. *In: McLean, A.C. & Deegan, C.E. (eds.). The solid geology of the Clyde sheet (55°N/6°W).* Report of the Institute of Geological Sciences, 77-91.
- Dobson, M.R. & Haynes, R. 1973. Association of foraminifera with hydroids on the deep shelf. *Micropalaeontology*, **19**, 78-90.
- Dobson, M.R. & Evans, D. 1974. Geological structure of the Malin Sea. *Journal of the Geological Society of London*, **130**, 475-478.
- Dobson, M.R., Garrett, P., Haynes, J.R., Jenkins, D.G. & Medani, A.H. 1976. Upper Cretaceous and Cenozoic carbonates from the margins of Rockall Trough, N. Atlantic. *Journal of the Geological Society of London*, **132**, Pt 6, 611-621.

- Dobson, M.R. & Whittington, R.J. 1992. Aspects of the geology of the Malin Sea area. *In: Parnell, J. (ed.). Post-Devonian development in the north-west seaboard of the British Isles.* **62**. Geological Society of London, London.
- Dodd, L.R. 1995. An investigation into the conditions and controls of submarine slope and stability in the Var Canyon, Nice, France and the Barra Fan, Hebrides, Scotland. Unpublished thesis, University of North Wales.
- Dodge, J.D. 1994. Biogeography of marine armoured dinoflagellates and dinocysts in the NE Atlantic and North Sea. *Review of Palaeobotany and Palynology*, **84**, 169-180.
- Dokken, T.M. & Jansen, E. 1999. Rapid changes in the mechanism of ocean convection during the last glacial period. *Nature*, **401**, 458-461.
- Dooley, H.D., Martin, J.H.A. & Payne, R. 1976. Flow across the Continental slope off northern Scotland. *Deep Sea Research*, **23**, 875-880.
- Dore, A.G., Lundin, E.R., Jensen, L.N., Birkeland, O., Eliassen, P.E. & Fichler, C. 1999. Principal tectonic events in the evolution of the Northwest European Atlantic margin. 41-61 in *Petroleum Geology of Northwest Europe; proceedings of the 5th conference*. Fleet, A.J. & Boldy, S.A.R. (editors). Geological Society of London.)
- Doré, A.G. & Lundin, E.R. 1996. Cenozoic compressional structures on the NE Atlantic margin: nature, origin and potential significance for hydrocarbon exploration. *Petroleum Geoscience*, **2**, 299-311.
- Doré, A.G., Lundin, E.R., Fichler, C. & Olesen, O. 1997. Patterns of basement structure and reactivation along the NE Atlantic margin. *Journal of the Geological Society of London*, **154**, 85-92.
- Dowling, L.M., Duplessey, J.C., McCave, I.N. & Shackleton, N.J. 1989. Deglacial events and chronology determined from sediments in the Rockall Trough. *Terra Abstracts*, **1**, 188.
- Dowling, L.M. 1990. Deglacial Climatic History recorded in sediments of the Rockall Trough. Unpublished PhD thesis, University of Cambridge.
- Dowling, L.M. & McCave, I.N. 1993. Sedimentation on the Feni Drift and late Glacial bottom water production in the northern Rockall Trough. *Sedimentary Geology*, **82**, 79-87.
- Eden, R.A., Ardu, D.A., Binns, P.E., McQuillin, R. & Wilson, J.B. 1971. Geological investigations with a manned submersible of the west coast of Scotland 1969-1971.
- Eden, R.A., Deegan, C.E., Rhys, G.H., Wright, J.E. & Dobson, M.R. 1973. Geological investigations with a manned submersible off the west coast of Scotland 1969-1970.
- Edwards, P.G. 1982. Ecology and distribution of selected foraminiferal species in the North Minch Channel, northwestern Scotland. *In: Banner, F.T. (ed.). Aspects of micropalaeontology; papers presented to Professor Tom Barnard*. George Allen and Unwin, London, 111-139.

Edwards, L.E. 1992. New semiquantitative (paleo)temperature estimates using dinoflagellate cysts, an example from the North Atlantic Ocean. *In: Head, M.J. & Wrenn, J.H. (eds.). Neogene and Quaternary dinoflagellate cysts and acritarchs.* Publishers Press. Salt Lake City, UT, United States., 69-87.

Eldholm, O. & Thiede, J. 1980. Cenozoic Continental separation between Europe and Greenland. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **30**, 243-259.

Ellet, D.J. 1988. Bottom topography to the west of the Wyville-Thomson Ridge. *Deutsche Hydrographische Zeitschrift*, **41**, Pt 1, 23-33.

Ellet, D.J. 1991. Cruise report: RRS "Challenger" Cruise 81/1991, 1-8 July 1991. Scottish continental shelf and Rockall Channel. *Cruise Report*, 5.

Ellet, D.J. & Edwards, A. 1996. The hydrography of the Rockall Channel - an overview. *Proceedings of the Royal Society of Edinburgh*, **88B**, 61-81.

Ellett, D.J. & Martin, J.H.A. 1973. The Physical and Chemical Oceanography of the Rockall Channel. *Deep-Sea Research*, **20**, 585-625.

Ellett, D.J. & Roberts, D.G. 1973. The overflow of Norwegian Sea deep water across the Wyville-Thomson Ridge. *Deep-Sea Research*, **20**, 819-835.

Ellett, D.J., Kruseman, P., Prangma, G.J., Pollard, R.T., Aken, H.M., A van Edward, Dooley, H.D. & Gould, W.J. 1983. Water masses and mesoscale circulation of North Rockall Trough waters during JASIN 1978. *Philosophical Transactions of the Royal Society of London. Series A*, **308**, 231-256.

Ellett, D.J. 1998. Norwegian Sea Deep Water overflow across the Wyville Thomson Ridge during 1987-1988. *Cooperative Research Report- International Council For The Exploration Of The Sea*, **225**, 195-205.

Emery, K.O. & Uchupi, E. 1984. *The Geology of the Atlantic Ocean.*

Ericson, D.B., Ewing, M., Woolin, G. & Heeaen, B.C. 1961. Atlantic Deep-Sea Sediment Cores. *Geological Society of America Bulletin*, **72**, 193-286.

Etheridge, M.A., Symonds, P.A. & Lister, G.S. 1989. Application of the detachment model to reconstruction of conjugate passive margins. *In: Tankard, A.J. & Balkwill, H.R. (eds.). Extensional Tectonics and Stratigraphy of the North Atlantic Margin.* Memoir of the American Association of Petroleum Geologists, No. 46, 23-40.

Evans, D. 1974. Geophysical studies in the Malin Sea. Unpublished thesis, University of Wales.

Evans, D. 1975. Whitehorn Cruise 75/WH/05, 4th-21st September 1975: Malin Sea. *Marine Geology Internal Report.*

Evans, D. & Ardu, D.A. 1976. Challenger Cruise 76/CH/06, 1-16 May, 1976.

Evans, D., Wilkinson, C.G. & Craig, D.L. 1979. The tertiary sediments of the Canna Basin, Sea of the Hebrides. *Scottish Journal of Geology*, **15**, 329-332.

Evans, D., Kenolty, N., Dobson, M.R. & Whittington, R.J. 1980. The geology of the Malin Sea.

Evans, D. & Ruckley, N.A. 1980. The geology and origin of narrow closed channels in the Sound of Jura. *Scottish Journal of Geology*, **16**, 65-72.

Evans, D., Stoker, M S. and McNeill, E. 1988. Plio-Pleistocene depocentres on the UK and Norwegian margins. 55-56 in *Neogene Uplift and Tectonics around the North Atlantic - International workshop, Copenhagen, May 18-19 1998*. Boldreel, L.a.J., P. (editor). Geological Survey of Denmark and Greenland (GEUS.)

Evans, D., Abraham, D.A. & Hitchen, K. 1989. The Geikie Igneous Centre, West of Lewis: its Structure and Influence on Tertiary Geology. *Scottish Journal of Geology*, **25**, 339-352.

Evans, D. & McElvanney, E.P. 1989. Geikie, Sheet 58°N - 10°W, 1:250 000 Map Series, Quaternary Geology. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.

Evans, D. & McElvanney, E.P. 1989. Lewis, Sheet 58°N - 08°W, 1:250 000 Map Series, Quaternary Geology. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.

Evans, D. & Strachan, P. 1989. Geikie, Sheet 58°N - 10°W, 1:250 000 Map Series, Sea Bed Sediments. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.

Evans, D., Hitchen, K. & Abraham, D.A. 1989. Geikie, Sheet 58°N - 10°W, 1:250 000 Map Series, Solid Geology. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.

Evans, D. & Hitchen, K. 1989. Sutherland, Sheet 58°N - 06°W, 1:250 000 Map Series, Solid Geology. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.

Evans, D. & McElvanney, E.P. 1989. Sutherland, Sheet 58°N - 06°W, 1:250 000 Map Series, Sea-bed Sediments and Quaternary Geology. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.

Evans, D. & Strachan, P. 1989. Lewis, Sheet 58°N - 08°W, 1:250 000 Map Series, Sea-bed Sediments. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.

Evans, D., Hitchen, K. & Robertson, S. 1990. Lewis, Sheet 58°N - 08°W, 1:250 000 Map Series, Solid Geology. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.

Evans, C.D.R., Selby, I. & Hopson, P.M. 1991. St. Kilda, Sheet 57N - 10 W, 1:250 000 Map Series, Sea-Bed Sediments. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.

Evans, C.D.R. 1992. St. Kilda, Sheet 57N - 10W, 1:250 000 Map Series, Solid Geology. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.

Evans, C.D.R. 1992. St. Kilda, Sheet 57N - 10W, 1:250 000 Map Series, Quaternary Geology. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.

Evans, D., Graham, C.C., Brett, C.P., Gillespie, E.J. & Wild, J.B.L. 1996. Slope Stability on the West Shetland Slope: a study of seismic profiles.

Evans, D., Morton, A.C., Wilson, S., Jolley, D. & Barreiro, B.A. 1997. Palaeoenvironmental significance of marine and terrestrial Tertiary sediments on the NW Scottish shelf in BGS borehole 77/7. *Scottish Journal of Geology*, **33**, 31-42.

Evans, D. 1997. Physiography, pathways and processes on the European Continental Margin. British Geological Survey 1: 5 000 000, Edinburgh.

Evans, D., Holmes, R., King, E., Meinert, J. & Nielson, T. 1997. Downslope processes. Part B in European North Atlantic Margin (ENAM): sediment pathways, processes and fluxes. Final 36 month report, MAST-CT93-0064.

Fairbanks, R.G. 1989. A 17,000 year glacio-eustatic sea level record: influence of glacial melting rates on the Younger Dryas event and deep-ocean circulation. *Nature*, **342**, 637-642.

Farrow, G.E., Cucci, M. & Scoffin, T.P. 1978. Calcareous sediments on the nearshore continental shelf of western Scotland. *Proceedings of the Royal Society of Edinburgh*, **76**, 55-76.

Farrow, G.E. 1978. Recent sediments and sedimentation in the Inner Hebrides. *Proceedings of the Royal Society of Edinburgh*, **83**, 91-105.

Farrow, G., Scoffin, T., Brown, B. & Cucci, M. 1979. An underwater television survey of facies variations on the inner Scottish shelf between Colonsay, Islay and Jura. *Scottish Journal of Geology*, **15**, 13-29.

Farrow, F.G. & Fyfe, J.A. 1988. Bioerosion and carbonate mud production on high-latitude shelves. In: Nelson, C.S. (ed.). *Non-tropical shelf carbonates - modern and ancient*. **60**. Sedimentary Geology.

Faugeres, J.C., Gonthier, E., Grousset, F. & Poutiers, J. 1981. The Feni Drift: The importance and meaning of slump deposits on the eastern slope of the Rockall Bank. *Marine Geology*, **40**, 49-57.

Faugeres, J.C., Stow, D A V., Imbert, P. and Viana, A. 1999. Seismic features diagnostic of contourite drifts. *Marine Geology*, **162 (1)**, 1-38.

Ferentinos, G.K. 1976. Sedimentary distribution and transport processes on the Outer Continental Shelf of the Hebridean Sea. *Marine Geology*, **20**, 41-56.

- Ferguson, M. 1996. *Environmental concerns in deepwater areas - the Atlantic Margin. Extended abstract.* (Towards 2000 - metres or millennium? Deepwater site investigation). Society for Underwater Technology,
- Ferragne, A., Latouche, C. & Parra, M. 1984. Lower Eocene red-clay sedimentation on the George Bligh/ Rockall channel. *Geo-Marine Letters*, **4**, 77-82.
- Ferry, S., Pastouret, L., J L de-Beaulieu & Mandier, P. 1985. Synchronisme remarquable entre les cycles calcaire-marne des vases quaternaires atlantiques et les alternances tourbe-limon d'anciens lacs periglaciaires europeens Translated Title: Remarkable synchronism between the Quaternary Atlantic lime-marl cycles. *Comptes-Rendus des Seances de l'Academie des Sciences, Serie 2: Mecanique-Physique, Chimie, Sciences*, **300**, 573-578.
- Flinn, D. 1969. On the development of coastal profiles in the North of Scotland, Orkney and Shetland. *Scottish Journal of Geology*, **5**, 393-399.
- Flood, R.D. & Hollister, C.D. 1975. Studies and significance of deep-sea bedforms in the North Atlantic. *Abstracts with Programs - Geological Society of America*, **7**, Pt 7, 1076.
- Flood, R., Lonsdale, P. & Hollister, C.D. 1976. Microtopography studies in the Rockall Trough, Northeast Atlantic. *Abstracts with Programs - Geological Society of America*, **8**, 868.
- Flood, R.D., Hollister, C.D. & Lonsdale, P. 1979. Disruption of the Feni sediment drift by debris flows from Rockall Bank. *Marine Geology*, **32**, 311-334.
- Folk, R.L. 1954. The distinction between grain size and mineral composition in sedimentary-rock nomenclature. *Journal of Geology*, **62**, 344-359.
- Ford, G.D., Simpson, B.A. & Walker, A.B. 1997. NW Scotland Offshore Seismicity: annual report to 31 December 1996.
- Francis, T.J.G. 1981. RRS 'Discovery' cruise 118, 11 February - 30 March 1981. Geophysics and sediment sampling in the North East Atlantic. pages 33.
- Frederiksen, R., Jensen, A. & Westerberg, H. 1992. The distribution of the scleractinian coral *Lophelia pertusa* around the Faroe Islands and the relation to internal tidal mixing. *Sarsia.*, **77**, 157-171.
- Freiwald, A., Wilson, J.B. & Ruediger, H. 1999. Grounding Pleistocene icebergs shape Recent deep water coral reefs. *Sedimentary Geology*, **125**, 1-8.
- Fyfe, J.A., Long, D., Evans, D. & Abraham, D.A. 1993. The geology of the Malin-Hebrides sea area. *Offshore Regional Report No. 4*, pages 91.
- Ganssen, G.M. & Kroon, D. 2000. The isotopic signature of planktonic foraminifera from NE Atlantic surface sediments; implications for the reconstruction of past oceanic conditions. In: Rothwell, R.A. (ed.). *Journal of the Geological Society of London. NE Atlantic palaeoceanography and climate change*. **157**. Geological Society of London., 693-699.

Gaston, G.F. 1976. The floor of the North Channel, Irish Sea: a side scan sonar survey.

GEBCO. 1977. Digital Atlas. *Intergovernmental Oceanographic Commission & International Hydrographic Organization, BODC, Merseyside, UK*, No. IOC/IHO.

Geikie, J. 1878. On the glacial phenomena of the Long Island, or Outer Hebrides. Second Paper. *Quarterly Journal of the Geological Society of London*, **34**, 819-870.

Gilbertson, D.D., Schwenninger, J.L., Kemp, R.A. & Rhodes, E.J. 1999. Sand-drift and soil formation along an exposed North Atlantic coastline: 14,000 years of diverse geomorphological, climatic and human impacts. *Journal Of Archaeological Science*, **26**, 439-469.

Gould, W.J. 1983. Current Measurements in the North Rockall Trough. *MIAS News Bulletin.*, **6**, 6.

Gradstein, F.M. 1998. Stratigraphic Resolution on Accelerated Neogene Subsidence, circum North Atlantic. 49-50 in *Neogene Uplift and Tectonics around the North Atlantic - International workshop, Copenhagen, May 18-19 1998*. Boldreel, L. & Jaspén, P. (editors). Geological Survey of Denmark and Greenland (GEUS).

Graham, D.K., Harland, R., Gregory, D.M., Long, D. & Morton, A.C. 1990. The Palaeontology and Chronostratigraphy of BGS borehole 78/4, North Minch, Scotland. *Scottish Journal of Geology*, **26**, 65-76.

Graham, C., Holmes, R., Wild, J.B. & Tulloch, G. 1996. Charles Darwin Cruise 101C - Geological Observations.

Graham, C., Holmes, R. & Long, D. 1997. Selected evidence for shallow gas from seabed to 20m below seabed. 1:1M. HSE: Offshore Safety Division.

Graham, C., Stewart, H.A., Poulton, C.V.L. & James, J.W.C. 2001. A Description Of Offshore Gravel Areas Around The UK. *Commissioned Report*, pages 26.

Graham, C., Campbell, E., Cavill, J., Gillespie, E. & Williams, R. 2001. JNCC Marine Habitats GIS Version 3: its structure and content. *Commissioned Report*, pages 45.

Gray, J.M. & Lowe, J.J. 1977. The Scottish Lateglacial environment; a synthesis. *In: Studies in the Scottish lateglacial environment*. Pergamon Press, Oxford, 163-181.

Gregersen, S. & Basham, P.W. 1989. *Earthquakes at North-Atlantic Passive Margins: Neotectonics and Postglacial Rebound*.

Gribbin, J. 1985. Rockall: rock, island or microcontinent. *New Scientist*, **107**, 18.

Hackett, B. & Roed, L.P. A numerical study of the slope current northwest of the British Isles. *Continental Shelf Research*, **18**, 1-30.

Hailwood, E.A., Kidd, R.B. & Dowling, L. 1987. The magnetic fabric of Neogene and Quaternary sediments on the Feni and Gardar Drifts, northeastern Atlantic, Deep Sea

Drilling Project Sites 620 and 611. *Initial Report of the Deep Sea Drilling Project*, **94**, 1083-1088.

Hall, J. 1970-71. Seismic studies in the region of the Firth of Clyde. Unpublished thesis, University of Glasgow.

Hall, B.D. & White, N. 1994. Origin of anomalous Tertiary subsidence adjacent to North Atlantic continental margins. *Marine and Petroleum Geology*, **11**, 702-714.

Hanisch, J. 1984. The Cretaceous opening of the Northeast Atlantic. *Tectonophysics*, **101**, 1-23.

Haq, B.U., Hardenbol, J. & Vail, P.R. 1987. The Chronology of fluctuating sea levels since the Triassic. *Science*, **235**, 1156-1167.

Harland, R. 1978. Quaternary and Neogene dinoflagellate cysts. In: Thusu, B. (ed.). *Distribution of biostratigraphically diagnostic dinoflagellate cysts and miospores from the Northwest European continental shelf and adjacent areas*. **100**. Institutt for Kontinentalsokkelundersokelser. Trondheim, Norway, 7-17.

Harland, R. 1983. Distribution maps of recent dinoflagellate cysts in bottom sediments from the North Atlantic Ocean. *Palaeontology*, **26**, 321-387.

Harland, R. 1994. Dinoflagellate cysts from the glacial/postglacial transition in the north-east Atlantic Ocean. *Palaeontology*, **37**, 263-283.

Harland, R. & Howe, J.A. 1995. Dinoflagellate cysts and Holocene oceanography of the Northeastern Atlantic Ocean. *Holocene*, **5 (2)**, 220-228.

Harrison, R.K. 1975. Expeditions to Rockall 1971-72.

Haughton, P., Praeg, D., Tyrrell, S. & Shannon, P.M. 2001. Maintenance and Collapse of a Cretaceous High on the SE Rockall Slope: Evidence from Shallow Seismic-Borehole Correlations in the Bróna Basins. In: Murphy, N.J. & Davies, M. (eds.). *Ireland's Deepwater Frontier: Results from the Petroleum Infrastructure Programme (PIP)*. *Extended Abstracts, Dublin*. 82-84.

Hawkes, J.R., Herriman, R.J., Harding, R.R. & Darbyshire, D.P.F. 1975. Rockall Island: new geological, petrological, chemical and Rb-Sr age data. In: Harrison, R.K. (ed.). *Expeditions to Rockall 1971-72*. Report of the Institute of Geological Sciences, No. 75/1.

Heaps, N.S. 1985. Storm-current events on the Scottish Continental Shelf. *MIAS News Bulletin*, **8**, 8-9.

Henriet, J.P., de Mol, B., Pillen, S., Vanneste, M., van Rooij, D., Versteeg, W., Croker, P.F., Shannon, P.M., Unnithan, V., Bouriak, S., Chachkine, P. & Belgica '97 Shipboard Party. 1998. Gas hydrate crystals may help build reefs. *Nature*, **391**, 648-649.

Hern, D. 1990. Quaternary topography in the Little Minch Sheet, 57-58N/6-8W. *BGS Marine Geology Technical Report*.

Hill, P.R. 1987. Characteristics of sediments from Feni and Gardar Drifts, Sites 610 and 611, Deep Sea Drilling Project Leg 94. *In: Initial Reports of the Deep Sea Drilling Project. 94.* US Government Printing Office, Washington DC, 1075-1082.

Hill, A.W. 1996. Site Investigation Studies into Seabed Stability of Slope Areas. 61 in *Applied Geoscience conference Warwick University 15-18 April 1996.* Geological Society, London.)

Himsworth, E.M. 1973. The Wyville-Thomson Ridge. *Quarterly Journal of the Geological Society of London, 129*, 322-323.

Himsworth, E.M. 1973. Marine geophysical studies between north-west Scotland and the Faeroe Plateau. Unpublished thesis, University of Durham.

Hine, N.H. 1990. Late Cenozoic Calcareous Nannoplankton from the Northeast Atlantic. Unpublished thesis, University of East Anglia.

Hiscock, K. 1992. The Ecology and Conservation of Sublittoral Hard Substratum Ecosystems in Scotland. *Proceedings of the Royal Society Edinburgh, 100B*, 95-112.

Hitchen, K. & Stoker, M.S. 1991. Judd, Sheet 60°N - 06°W, 1:250 000 Map Series, Solid Geology. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.

Hitchen, K. 1992. The crustal characteristics, volcanic and sedimentary history of the Rockall Continental Margin. *BGS Technical Report.*

Hitchen, K. & Stoker, M.S. 1993. Mesozoic rocks from the Hebrides Shelf and implications for hydrocarbon prospectivity in the northern Rockall Trough. *Marine and Petroleum Geology, 10*, 246-254.

Hitchen, K., Morton, A.D., Mearns, E.W., Whitehouse, M. & Stoker, M.S. 1997. Geological implications from geochemical and isotopic studies of Upper Cretaceous and lower Tertiary igneous rocks around the northern Rockall Trough. *Journal of Geological Society of London, 154*, Pt 3, 517-521.

Hobbs, P. 1988. Geotechnical testing of glaciomarine clays from St. Kilda.

Hoffmann, N. 1976. The importance of calcareous nannoplankton in marine geology and early biostratigraphical results of investigating sedimentary cores from the Rockall Plateau (north-east Atlantic). *Z Geol Wiss [Berlin], 4/4*, 647-659.

Holland, G.S. & Gardiner, R.A. 1975. The first map of Rockall. *Geographical Journal, 141*, 94-98.

Holliday, N.P., Pollard, R.T. & Read, J.F. 1999. Water mass properties and fluxes in the Rockall Trough; 1975 to 1998. *Deep-Sea Research.*

Holliday, N.P., Pollard, R.T. & Leach, H.a.R., J F. 1999. Water mass transports in the Rockall Trough, 1975 to 1998. *North-East Atlantic Slope Processes: Multi-Disciplinary Approaches, TTR-8 Post Cruise Conference/4th ENAM II Workshop/IGCP Workshop 432, SOC, Southampton, 22-27 January 1999.* Southampton Oceanography Centre.)

Holmes, N.A. 1987. Distribution, Ecology and Palaeoenvironmental analysis of Recent Planktonic Foraminifera from the Rockall Trough, NE Atlantic. Unpublished PhD thesis, University of Wales.

Holmes, R. & Stoker, M.S. 1990. Pleistocene glaciomarine facies on the continental shelf and slope north-west of Scotland. 230 in *Sediments 1990. 13th International Sedimentological Congress. Nottingham, England. 26th-31st August 1990. Abstract of Papers*. International Association of Sedimentologists, Comparative Sedimentology Division. Utrecht.)

Holmes, R., Ruckley, N., Jeffery, D. & Wingfield, R. 1992. Compilation of the 1:1M scale map of Quaternary sediments around the UK. *BGS Marine Geology Technical Report*.

Holmes, R. 1994. Seabed topography and other geotechnical information for the shelf edge 55-60N NW of Britain. *BGS Technical Report*.

Holmes, R. 1995. Proposals for piston core sites west of the Hebrides and Shetland Isles.

Holmes, R., Long, D. & Dodd, L.R. 1996. Debris flows and sediment slides on the Barra Fan, northwest of Britain. 67 in *Applied Geoscience conference Warwick University 15-18 April 1996*. Geological Society, London.)

Holmes, R., Finlayson, K., Griffiths, M.A. & Andresen, P.C. 1996. Regional Shallow Gas Distribution Study.

Holmes, R., Alexander, S., Ball, K., Bulat, J., Evans, R., Long, D., MacBeth, C., McCormack, M. & Sankey, M. 1996. The issues surrounding a shallow gas database in relation to offshore hazards. *Health & Safety Executive Offshore Safety Division Report OTH 96 504*.

Holmes, R. 1997. Shelf Edge and Oceanward: Britains Western Frontier Wealth and Health. *Scottish Association for Marine Science*, **15**.

Holmes, R. 1997. Quaternary stratigraphy: the offshore record: Chapter 7. In: Gordon, J.E. (ed.). *Reflections on the ice age in Scotland: an update on Quaternary studies*. Scottish Natural Heritage.

Holmes, R., Long, D. & Dodd, L.R. 1998. Large-scale debrites and submarine landslides on the Barra Fan, west of Britain. In: Stoker, M.S., Evans, D. & Cramp, A. (eds.). *Geological Processes on Continental Margins: Sedimentation, Mass-Wasting and Stability*. Geological Society, London, *Special Publications, vol 129*. Geological Society, London, 66-79.

Holmes, R., Long, D. & Jones, S. 1998. RRS Charles Darwin Cruise 112C - geological observations from seabed and shallow sediments 17th Round Tranches.

Holmes, R. 1999. Regional Shallow Gas Study: UK 17th Round License Tranches, NW of Scotland.

- Holmes, R. 2002. Holocene shelf-margin submarine landslides, Donegal Fan, eastern Rockall Trough. *In: Meinert, J. & Weaver, P.P.E. (eds.). European Margin Sediment Dynamics: Sidescan Sonar and Seismic Images.* Springer-Verlag, New York.
- Holmes, R., Bulat, J., Hamilton, I. & Long, D. In press. Morphology of an ice-sheet limit and contractional glacially-fed slope front, Faroe-Shetland Channel. *In: Meinert, J. & Weaver, P.P.E. (eds.). European Margin Sediment Dynamics: Sidescan Sonar and Seismic Images.* Springer-Verlag, New York.
- Hovland, M. & Judd, A.G. 1988. *Seabed pockmarks and seepages: impact on geology, biology and the marine environment.* Graham & Trotman, London.
- Howarth, M.J. & Harrison, A.J. 1980. RRS "Shackleton" Cruise 7/79, 3-15 August 1979, RRS "Challenger" Cruise 14/79, 21 September - 2 October 1979. North Channel/Malin Shelf Sea. *Institute of Oceanographic Sciences Cruise Report*, pages 45.
- Howarth, M.J. & Heaps, N.S. 1981. RRS "John Murray" Cruise 9/80, 27 August - 9 September 1980, RRS "Challenger" Cruise 16/80, 17-30 October 1980. Celtic Sea. *Institute of Oceanographic Sciences Cruise Report*, pages 45.
- Howe, J.A., Stoker, M.S. & Stow, D.A.V. 1994. A late Cenozoic sediment drift complex, Northern Rockall Trough, North Atlantic. *Palaeoceanography*, **9**, 989-999.
- Howe, J.A. 1994. Turbidite and contourite sediment waves from the northern Rockall Trough, North Atlantic. S2.8-S2.9 in *14th International Sedimentological Congress. abstracts.* International Association of Sedimentologists, Comparative Sedimentology Division. Utrecht, Netherlands.)
- Howe, J.A. 1994. Bottom currents, contourites and related sedimentation in the northern Rockall Trough, North Atlantic Ocean. Unpublished thesis, University of Southampton.
- Howe, J.A. & Harland, R. 1995. Dinoflagellate cysts and the Holocene Oceanography of the north-eastern Atlantic Ocean. *Holocene*, **5**, 220-228.
- Howe, J.A. 1995. Sedimentary processes and variations in slope current activity during the last glacial-interglacial episode on the Hebrides Slope, Northern Rockall Trough, North Atlantic Ocean. *Sedimentary Geology.*, **96**, 201-230.
- Howe, J.A. & Humphery, J.D. 1995. Photographic evidence for slope current activity, Hebrides Slope, Northern Rockall Trough, North Atlantic Ocean. *Scottish Journal of Geology*, **31**, 107-115.
- Howe, J.A. 1996. Sedimentary Processes on the Hebrides Slope, northern Rockall Trough, North Atlantic Ocean. 66 in *Applied Geoscience conference Warwick University 15-18 April 1996.* Geological Society, London.)
- Howe, J.A. 1996. Sandy Contourite occurrence, deposition and the implications for the facies model. 70 in *Applied Geoscience conference Warwick University 15-18 April 1996.* Geological Society, London.)
- Howe, J.A. 1996. Turbidite and contourite sediment waves in the northern Rockall Trough, North Atlantic Ocean. *Sedimentology*, **43**, 219-234.

- Howe, J.A., Harland, R., Hine, N.H. & Austin, W.E.N. 1998. Late Quaternary stratigraphy and palaeoceanographic change in the northern Rockall Trough, North Atlantic Ocean. *In: Stoker, M.S., Evans, D. & Cramp, A. (eds.). Geological Processes on Continental Margins: Sedimentation, Mass-Wasting and Stability.* (Geological Society of London Special Publications). **129**. Geological Society of London, London, 269-286.
- Howe, J.A., Stoker, M.S. & Woolfe, K.J. 2001. Deep-marine seabed erosion and gravel lags in the northwestern Rockall Trough, North Atlantic Ocean. *Journal of the Geological Society*, **158**, 427-438.
- Howe, J.A., Overnell, J., Inall, M.E. & Wilby, A. In press. A side scan image of a glacially-overdeepened sea loch, upper Loch Etive, Argyll. *Scottish Journal of Geology*.
- Hoy, T., Boldreel, L.O., Andersen, M.S. & Kiorboe, L. 1992. A high resolution seismic survey in the Faroe-Rockall area. *Journal of Applied Geophysics*, **29**, 85.
- Huggett, Q. 1985. The distribution of Glacial Erratics in the Northeast Atlantic: Final Report.
- Hughes, D.J. 2001. Quantitative analysis of a deep-water bryozoan collection from the Hebridean continental slope. *Journal of the Marine Biological Association of the UK*, **81**, 987-993.
- Hunt, J.B., Fannin, N.G., Hill, P.G. & Peacock, J.D. 1995. The tephrochronology and radiocarbon dating of North Atlantic, Late Quaternary sediments: an example from the St. Kilda Basin. *In: Scrutton, R.S., Stoker, M.S., Shimmield, G.B. & Tudhope, A.W. (eds.). Geological Society Engineering Geology Special Publication. Quaternary Engineering Geology.* **90**. 227-248.
- Huthnance, J.M. 1986. The Rockall slope current and shelf-edge processes. *In: Mauchline, J. (ed.). The oceanography of the Rockall Channel.* **88B**. Royal Society of Edinburgh, 83-101.
- Hutton, D.H.W. 1987. Strike-slip terranes and a model for the evolution of the British and Irish Caledonides. *Geological Magazine*, **124**, 405-425.
- Hydrographic Office. 1990. Admiralty tide tables 1991. **1**.
- IGS. 1968. RRS Discovery Cruise 24, September 1968: Geological investigations west of the British Isles and north of Scotland. *NIO Cruise Report*.
- Inall, M.E., Shapiro, G.I. & Sherwin, T.J. 2001. Mass transport by non-linear internal waves on the Malin Shelf. *Continental Shelf Research*, **21**, 1449-1472.
- Ivanov, M., Kenyon, N.H., Henriot, J.P., Swennen, R. & Limonov, A. 1998. Carbonate mud mounds and cold water corals in the Porcupine Seabight and Rockall Bank: are they methane related? 37 in *Carbonate mud mounds and cold water reefs: deep-biosphere-geosphere coupling. TTR-7 post cruise conference.* De Mol, B. (editor). (Gent, Belgium: University of Gent.)

- Jackson, D.I., Mulholland, P., Jones, S.M. & Warrington, G. 1987. The geological framework of the East Irish Sea Basin. *In: Brooks, J. & Glennie, K.W. (eds.). Petroleum Geology of North West Europe*. London: Graham and Trotman, 191-203.
- Jackson, P.D., Gunn, D.A., Stoker, M.S., Holmes, R., Hobbs, P.R.N., Long, D., Walker, A. & Musson, R. 1999. Impact of earthquake ground motion on the stability of slope sediments using existing ground models.
- Jacob, A.W.B., Neilson, G. & Ward, V. 1983. A Seismic event near the Hebrides Terrace Seamount. *Scottish Journal of Geology*, **19**, 287-296.
- Jacobs, C.L., Masson, D.G. & Bett, B.J. 1999. Distribution and biology of recently discovered carbonate mounds in the northern Rockall Trough. *Reprint*, 2.
- James, J.W.C. & Wright, S.A. 1990. Peach, Sheet 56N - 10W, 1:250 000 Map Series, Sea-bed Sediments. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.
- James, J.W.C. & Hitchen, K. 1992. Peach, Sheet 56N - 10W, 1:250 000 Map Series, Solid Geology. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.
- James, J.W.C. 1992. Peach, Sheet 56N - 10W, 1:250 000 Map Series, Quaternary Geology. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.
- Jansen, E. & Veum, T. 1990. Evidence for two-step deglaciation and its impact on North Atlantic deep-water circulation. *Nature*, **343**, 612-616.
- Jansen, E., Raymo, M.E. & Blum, P. 1996. 3. Sites 980/981. *In: Jansen, E., Raymo, M.E. & Blum, P. (eds.). Proceedings ODP, Initial reports, 162*. Ocean Drilling Program, College Station, TX, 49-90.
- Japsen, P. 1997. Regional Neogene exhumation of Britain and the western North Sea. *Journal of the Geological Society of London*, **154**, 239-247.
- Japsen, P. & Chalmers, J.A. 2000. Neogene uplift and tectonics around the North Atlantic: overview. *Global and Planetary Change*, **24**, 165-173.
- Jardine, W.G. 1977. The Quaternary marine record in south west Scotland and the Scottish Hebrides. *In: Kidson, C. & Tooley, M.J. (eds.). The Quaternary History of the Irish Sea*. Seel House Press, Liverpool, 99-118.
- Jehu, T.J. & Craig, R.M. 1934. Geology of the Outer Hebrides. Part 4 North Harris and Lewis. *Transaction of the Royal Society of Edinburgh*, **57**, 839-874.
- Johnson, G.L. & Schneider, E.D. 1969. Depositional Ridges in the North Atlantic. *Earth and Planetary Science Letters*, **6**, 416-422.
- Johnson, G.L., Vogt, P.R. & Schneider, E.D. 1971. Morphology of the Northeastern Atlantic and Labrador Sea. *Deutsche Hydrographische Zeitschrift*, **B24**, 49-73.

- Johnson, M.A., Kenyon, N.H., Belderson, R.H. & Stride, A.H. 1982. Sand transport. *In*: Stride, A.H. (ed.). *Offshore tidal sands - processes and deposits*. Chapman and Hall, London, 58-94.
- Jones, E.J.W., Ewing, M., Ewing, J.I. & Eittreim, S.L. 1970. Influences of Norwegian Sea overflow water on sedimentation in the Northern North Atlantic and Labrador Sea. *Journal of Geophysical Research*, **75**, 1655-1680.
- Jones, E.J.W., Mitchell, J.G., Shido, F. & Phillips, J.D. 1972. Igneous rocks dredged from the Rockall Plateau. *Nature Physical Science*, **237**, 118-120.
- Jones, E.J.W., Ramsay, A.T.S., Preston, N.J. & Smith, A.C.S. 1974. A Cretaceous guyot in the Rockall Trough. *Nature*, **251**, 129-131.
- Jones, E.J.W. & Ramsay, A.T.S. 1982. Volcanic ash deposits of early Eocene age from the Rockall Trough. *Nature*, **299**, 342-344.
- Jones, E.J.W., Perry, R G. and Wild, J L. 1986. Geology of the Hebridean Margin of the Rockall Trough. *Proceedings of the Royal Society of Edinburgh.*, **88B**, 27-51.
- Jones, E.J.W., Siddall, R., Thirlwall, M.F., Chroston, P.N. & Lloyd, A.J. 1994. Anton Dohrn Seamount and the evolution of the Rockall Trough. *Oceanologica Acta*, **17**, Pt 3, 237-247.
- Joppen, M., White, R.S., Spence, G.D. & Westbrook, G.K. 1987. The seismic structure of Rockall Trough. *Eos, Transactions, American Geophysical Union*, **68**, Pt 44, 1372.
- Joy, A.M. 1992. Right place, wrong time: anomalous post-rift subsidence in sedimentary basins around the North Atlantic Ocean. *In*: Storey, B.C., Alabaster, T. & Pankhurst, R.J. (eds.). *Magmatism and the Causes of Continental Break-up*. **68**. Geological Society Special Publication, London, 387-393.
- Judd, A., Davies, G., Wilson, J., Holmes, R., Baron, G. & Bryden, I. 1997. Contributions to atmospheric methane by natural seepages on the UK continental shelf. *In*: T C E van Weering, Klaver, G.T. & Prins, R.A. (eds.). *Marine Geology. Gas in marine sediments, geology, geochemistry, microbiology*. **137**. Elsevier, Amsterdam, Netherlands, 165-189.
- Jung, S.J.A. & Sarnthein, M. 1995. North Atlantic minima in deep water ventilation linked to Heinrich events during the last 250,000 years. *In*: Anon (ed.). *Global change and marine geology; 85. Jahrestagung der Geologischen Vereinigung e.V.; Zusammenfassung der Tagungsbeiträge*. **1-95**. Alfred-Wegener-Stiftung. Bonn, Federal Republic of Germany, 23-24.
- Kenyon, N.H. & Stride, A.H. 1970. The tide-swept continental shelf sediments between the Shetland Isles and France. *Sedimentology*, **14**, 159-173.
- Kenyon, N.H. & Pelton, C.D. 1980. Seabed conditions west of the Outer Hebrides. *Report*, pages 8.
- Kenyon, N.H. 1986. Evidence from bedforms for a strong poleward current along the upper Continental Slope of Northwest Europe. *Marine Geology*, **72**, 187-198.

Kenyon, N.H. 1987. Mass wasting features on the continental slope of northwest Europe. *Marine Geology*, **74**, 57-77.

Kenyon, N.H., Ivanov, M.K. & Akhmetzanov, A.M. 1998. Cold water carbonate mounds and sediment transport on the Northeast Atlantic margin: preliminary results of geological and geophysical investigations during the TTR-7 cruise of R/V "Professor Logachev". *Technical Series*, pages 178.

Kenyon, N.H. 1998. II. Rockall Trough (Leg II). II.1. Geological setting and objectives. *In: Cold water carbonate mounds and sediment transport on the Northeast Atlantic margin: preliminary results of geological and geophysical investigations during the TTR-7 cruise of R/V "Professor Logachev"*. 108-109.

Kenyon, N.H. & Shipboard Scientific Party of RV Professor Logachev TTR7 Cruise. 1999. Efforts of bottom trawling in deep water, west of Ireland and Scotland. 45 in *North East Atlantic Slope Processes: Multi-Disciplinary Approaches, Abstract Book. Conference 24-27th January 1999*. Friend, P. & Kenyon, N. (editors). Southampton Oceanography Centre, Southampton, UK.)

Kenyon, N.H., Ivanov, M.K. & Akhmetzhanov, A.M. 1999. *Geological processes in the Northeast Atlantic margin; June-August 1998*. (Serie Technique - UNESCO, Commission Oceanographique Intergouvernementale. Geological processes on the Northeast Atlantic margin, June-August 1998). UNESCO, Commission Oceanographique Intergouvernementale, Paris, International,

Kidd, R.B. & Hill, P.R. 1986. Sedimentation on mid-ocean sediment drifts. *In: Summerhayes, C.P. & Shackleton, N.J. (eds.). North Atlantic Palaeoceanography*. (Geological Society of London Special Publications). **21**. Blackwells, Oxford, 87-102.

Kidd, R.B. & Hill, P.R. 1987. Sedimentation on Feni and Gardar sediment drifts. *Initial Report of the Deep Sea Drilling Project*, **94**, Pt 2, 1217-1244.

Kidd, Robert, B. 1982. Long-range sidescan sonar studies of sediment slides and the effects of slope mass sediment movement on abyssal plain sedimentation. *In: Saxov, S. & Nieuwenhuis, J.K. (eds.). NATO Conference Series. IV. Marine Sciences. Marine slides and other mass movements*. **6**. Plenum, New York, 289-303.

Kilenyi, T. & Standley, R. 1985. Petroleum prospects in the northwest seaboard of Scotland. *Oil and Gas Journal*, **83**, 100-101, 104, 106, 108.

Kiriakoulakis, K., Bett, B.J. & Wolff, G.A. 2001. Biogeochemistry of the Darwin Mounds - Preliminary Results. *Journal of Conference Abstracts*, **6**, 751.

Kirton, S.R. & Hitchen, K. 1987. Timing and style of crustal extension north of the Scottish mainland. *In: Continental extensional tectonics*. **28**. Geological Society of London, London, 501-510.

Knaap, R.J. 1972-73. The form and structure of the Islay, Jura and Arran Tertiary basic dyke swarms. Unpublished thesis, University of City of London Polytechnic.

Knott, S.D., Burchell, M.T., Jolley, E.W. & Fraser, A.J. 1993. Mesozoic to Cenozoic plate reconstructions of the North Atlantic and hydrocarbon plays of the Atlantic

margins. In: Parker, J.R. (ed.). *Petroleum Geology of Northwest Europe: Proceedings of the 4th conference*. Geological Society of London, London, 953-974.

Knox, R.W.O.B. & Morton, A.C. 1988. The record of early Tertiary North Atlantic volcanism in sediments of the North Sea Basin. In: Morton, A.C. & Parson, L.M. (eds.). *Early Tertiary Volcanism and the Opening of the NE Atlantic*. Geological Society Special Publication, No. 39, 407-419.

Knutz, P.C. 1998. Glaciomarine sedimentary cycles in the northern Rockall Trough, NE Atlantic margin. 170 in *GeoScience 98, (Keele University, 14-18 April 1998) Abstracts Volume*. Geological Society of London.)

Knutz, P.C. 1999. Contourite deposition, mass flows and ice rafting events on the lower Barra Fan, Rockall Trough. 46 in *North East Atlantic Slope Processes: Multi-Disciplinary Approaches, Abstract Book. Conference 24-27th January 1999*. Friend, P. & Kenyon, N. (editors). Southampton Oceanography Centre, Southampton, UK.)

Knutz, P.C. 1999. Late Pleistocene mass flow events on the northern Barra Fan: relationship between seismic facies and lithology. 46 in *North East Atlantic Slope Processes: Multi-Disciplinary Approaches, Abstract Book. Conference 24-27th January 1999*. Friend, P. & Kenyon, N. (editors). Southampton Oceanography Centre, Southampton, UK.)

Knutz, P.C. 2000. Late Pleistocene glacial fluctuations and palaeoceanography on the continental margin of north-west Britain. Unpublished thesis, University of Wales.

Knutz, P.C., Austin, W.E.N. & Jones, E.J.W. 2001. Millennial-scale depositional cycles related to British ice sheet variability and North Atlantic paleocirculation since 45 kyr B.P., Barra Fan, U.K. margin. *Palaeoceanography*, **16**, 53-64.

Knutz, P.C., Jones, E.J.W., Austin, W.E.N. & T C E van Weering. 2002. Glacimarine slope sedimentation, contourite drifts and bottom current pathways on the Barra Fan, UK North Atlantic margin. *Marine Geology*, **188**, 129-146.

Koster, K., Boldreel, L.O. & Kuijpers, A. 1999. Evidence of bottom current activity around Lousy Bank, NE Atlantic, illustrated by high-resolution reflection seismic data. 47 in *North East Atlantic Slope Processes: Multi-Disciplinary Approaches, Abstract Book. Conference 24-27th January 1999*. Friend, P. & Kenyon, N. (editors). Southampton Oceanography Centre, Southampton, UK.)

Kroon, D. & Austin, W.E.N. 1995. High resolution records of the last glacial/interglacial transition. *Proceedings of the Royal Dutch Academy of Science*, **44**, 125-130.

Kroon, D., Austin, W.E.N. & Chapman, M.R. 1995. High resolution palaeoceanographic studies of the Hebridean continental margin, N W Scotland. *QRA Annual Discussion Meeting, 1995 Abstract Volume The Lateglacial Palaeoceanography of the North Atlantic Margins*. Andrews, J.T., Austin, W.E.N. & Bergsten, H.E. (editors).

Kroon, D., Austin, W.E.N., Chapman, M.R. & Ganssen, G.M. 1997. Deglacial surface circulation changes in the northeastern Atlantic: Temperature and salinity records off NW Scotland on a century scale. *Palaeoceanography*, **2**, 755-763.

- Kroon, D., Shimmiel, G., Austin, W.E.N., Derrick, S., Knutz, P. & Shimmiel, T. 2000. Century - to millennial-scale sedimentological-geochemical records of glacial-Holocene sediment variations from the Barra Fan (NE Atlantic). *In: Rothwell, R.A. (ed.). Journal of the Geological Society of London. NE Atlantic palaeoceanography and climate change.* **157**. Geological Society of London, 643-653.
- Kuijpers, A., Troelstra, S.R., Wisse, M., Nielsen, S.H. & T C E van Weering. 1998. Norwegian Sea overflow variability and NE Atlantic surface hydrography during the past 150,000 years. *Marine Geology*, **152**, 75-99.
- Kuijpers, A., Andersen, M.A., Kenyon, N.H., Kunzendorf, H. & T C E van Weering. 1998. Quaternary sedimentation and Norwegian Sea overflow pathways around Bill Bailey Bank, north-eastern Atlantic. *Marine Geology.*, **152**, 101-127.
- Kuijpers, A.H., Andersen, M.S., Kenyon, N.H., Kunzendorf, H. & van Weering, T.C.E. 1998. Quaternary sedimentation and Norwegian Sea overflow pathways around Bill Bailey Bank, northeastern Atlantic. *Marine Geology*, **152**, Pt 1-3, 101-127.
- Labeyrie, L. & Duplessy, J.C. 1985. Changes in the oceanic $^{13}\text{C}/^{12}\text{C}$ ratio during the last 140 000 years: high latitude surface water records. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **50**, 217-240.
- Labeyrie, L., Elliot, M., Raffalli, G., Cortijo, E., Balbon, E., Tj C E Van Weering & Rasmussen, T. 1996. The Northern Atlantic records of the Dansgaard Oeschger cycling, causes and consequences. *AGU 1996 Fall Meeting, Dec. 15-19, San Francisco, supplm. EOS, Transactions, AGU*, **77**.
- Labracherie, M. & Moyes, J. 1978. Importance of the radiolarians as indicators of deep oceanic circulation. As for example, Northeast Atlantic Ocean in the Rockall and Faeroe Regions. *Series D: Sciences Naturelles*, **286**, 1425-1428.
- Lachelt, A. 1975. Geochemical - Lithological conditions of the Rockall plateau. *Z Geol Wiss [Berlin]*, **3/11**, 1404-1413.
- Lambeck, K. 1991. Glacial rebound and sea-level change in the British Isles. *Terra Nova*, **3**, 379-389.
- Lambeck, K. 1995. Glacial isostasy and water depths in the Late Devensian and Holocene on the Scottish Shelf west of the Outer Hebrides. *Journal of Quaternary Science*, **10**, 83-86.
- Lassen, S., Jansen, E., Knudsen, K.L., Kuijpers, A., Kristensen, M. & Christensen, K. 1999. Northeast Atlantic sea surface circulation during the past 30-10 ^{14}C kyr BP. *Palaeoceanography*, **14**, 616-625.
- Laughton, A.S. & Roberts, D.G. 1978. Morphology of the Continental Margin. *Philosophical Transactions of the Royal Society of London*, **290A**, 75-85.
- Laughton, A.S., Roberts, D.G. & Hunter, P.M. 1982. Bathymetry of the northeast Atlantic. Sheet 1 : Reykjanes Ridge and Rockall Plateau. Scale 1 : 2,400,000 at 41N. Bathymetry 1:2500 000.

Le Coeur, C. 1988. Late Tertiary warping and erosion in Western Scotland. *Geografiska Annaler*, **70**, 361-368.

Lee, S.H. 1971-72. Sample disturbance in soft alluvial Clyde Estuary clay. Unpublished thesis, University of Strathclyde.

Lee, A.J. & Ramster, J.W. 1981. Atlas of the Sea around the British Isles.

Leslie, A.B. 1990. Shallow Plio-Pleistocene Contourites on the Northwest UK Continental Slope in the Northern Rockall Trough. 305 in *Sediments 1990. 13th International Sedimentological Congress. Nottingham, England. 26th-31st August 1990. Abstract of Papers*. International Association of Sedimentologists, Comparative Sedimentology Division. Utrecht.)

Leslie, A.B., Scott, W.D., Briden, J. & Stoker, M.S. 1991. Sedimentology and Magnetostratigraphy of Plio-Pleistocene ice-distal fine-grained clastics, north-west UK Continental Margin. 351 in *Terra Abstracts. Sixth Meeting of European Union of Geosciences*. Blackwell Scientific Publications. Oxford.)

Leslie, A.B. 1992. A sedimentological study of Tertiary and Quaternary sediments in borehole 88/7, 7A, Hebrides Slope, Northern Rockall Trough.

Leslie, A.B. 1993. Shallow Plio-Pleistocene contourites on the Hebrides Slope, north-west UK continental margin. *Sedimentary Geology*, **82**, 61-78.

Leslie, A. 1993. Shallow Plio-Pleistocene contourites on the Hebrides Slope, Northwest U.K. continental margin. *Sedimentary Geology*, **82**, Pt 1-4, 61-78.

Liu, A.K., Kasischke, E.S., Borchardt, S.R. & Shuchman, R.A. 1983. Detection of bottom features on SEASAT SAR imagery. *Eos, Transactions, American Geophysical Union*, **64 part 52**, 1052.

Lohse, L., Rikus, T., Kloosterhuis, H.C., de Stigter, W.H., van Raaphorst, W. & van Weering, T.C.E. 2000. Carbonate removal by acidification causes loss of nitrogenous compounds in continental margin sediments. *Marine Chemistry*, **69 Issues 3-4**, 193-201.

Long, D. & Fannin, N.G.T. 1979. Potential pipeline approaches to Western Scotland.

Long, D. & Bone, B. 1990. Sediment instability on the continental slope of north-western Europe. *Proceedings of Oceanology International 1990.*, **2**.

Long, D. 1992. A review of sediment instability on the continental slope of north-western Europe (Abstract). *Annales Geophysicae*, **10**, C109.

Long, D., Walker, P M., Hill, A W. and Ardu, D A. 1998. The Western Frontiers Association - an example of joint industry funded regional studies of the shallow geology in a frontier area. 123-138 in *Offshore Site Investigation and Foundation Behaviour - new frontiers. Conference proceedings 22-24 September 1998*. Ardu, D.A., Hobbs, R., Horsnell, M., Jardine, R. & Long, D. (editors). Society for Underwater Technology, London.)

- Long, D. 1998. Geological Characteristics and Geohazards on the Hebridean and West Continental Shelf and Slope. pages 54.
- Long, D., Holmes, R. & Wild, J.B.L. 1998. RRS Charles Darwin Cruise 112C - sample descriptions.
- Long, D., Roberts, J.M. & Gillespie, E.J. 1999. Occurrences of *Lophelia Pertusa* on the Atlantic Margin.
- Long, D. 2001. The Western Frontiers Association: Evaluating seabed conditions west of the UK. *Continental Shelf Research.*, **21 Issues 8-10**, 811-824.
- Lonsdale, P.F. & Hollister, C.D. 1979. A near bottom traverse of Rockall Trough: Hydrographic and geologic inferences. *Oceanologica Acta*, **2**, 91-105.
- Lonsdale, P.F. 1982. Sediment drifts of the north-east Atlantic and their relationship to the observed abyssal currents. *Bulletin de l'institut du Geologie du Bassin d'Aquitaine*, **31**, 141-149.
- Lovell, J.P.B. 1979. composition of Holocene sands of Mull and adjacent offshore areas: a study of provenance. *IGS Report*, pages 9.
- Lundberg, P. 1993. Comments on Cold outflow from the Faeroe Bank Channel. *Journal of Physical Oceanography*. No 6, **23**, 1285-1291.
- MacKensen, A. 1987. Benthische Foraminiferer auf Island-Schottland Rucken: Umwelt-Anzeiger an der Grenze zweier ozeanischer Raume. (Benthic Foraminifers on the Iceland-Scotland Ridge, environmental indicator at the boundary between two oceanic regions. *Palaeontologische Zeitschrift*, **61**, 149-179.
- Makris, J., Ginzburg, A., Shannon, P.M., Jacob, A.W.B., Bean, C.J. & Vogt, U. 1991. A new look at the Rockall region, offshore Ireland. *Marine and Petroleum Geology*, **8**, 410-416.
- Manighetti, B. 1993. The Glacial to Holocene Sedimentary Regime in the NE Atlantic Ocean. Unpublished thesis, University of Cambridge.
- Manighetti, B. & McCave, I.N. 1995. Late glacial and Holocene palaeocurrents around Rockall Bank, NE Atlantic Ocean. *Palaeoceanography*, **10**, 611-626.
- Manighetti, B. & McCave, I.N. 1995. Depositional fluxes, paleoproductivity and ice rafting in the NE Atlantic over the past 30ka. *Palaeoceanography*, **10**, 579-592.
- Marca, S. & Cambray, H. 1996. Late Pleistocene Changes Recorded in Physical Properties in NE Atlantic Sediments. *AGU 1996 Fall Meeting, Dec.15-19, San Francisco, supplem. EOS, Transactions, AGU*, **77**, **abstract:F22**.
- Maslin, M.A. 1993. A study of the Palaeoceanography of the NE Atlantic in the Late Pleistocene. Unpublished thesis, University of Cambridge.

- Maslin, M.A. & Shackleton, N.J. 1995. Surface water temperature, salinity, and density changes in the north-east Atlantic during the last 45,000 years: Heinrich events, deep water formation, and climatic rebounds. *Palaeoceanography*, **10**, 527-544.
- Maslin, M.A., Thomas, E., Shackleton, N.J., Hall, M.A. & Seidov, D. 1997. Glacial northeast Atlantic surface water pCO₂: Productivity and deep-water formation. *Marine Geology*, **144**, 177-190.
- Masson, D.G. & Kidd, R.B. 1986. Tertiary seismic stratigraphy of the Southern Rockall Trough. *In: Ruddiman, W.F. & Kidd, R.B. (eds.). Initial Reports of the Deep Sea Drilling Project, Leg 94*. Washington DC: US Government Printing Office, 1117-1126.
- Masson, D.G. 1996. Atlantic Margin Environmental Survey - Interpretation of Sidescan Sonar Data: Final Report. *In: Network, A.F.E. (ed.). UKCS Atlantic Margins Environmental Survey of the Seafloor [CD-ROM]*. Southampton Oceanography Centre, Southampton.
- Masson, D.G. 1997. RRS Charles Darwin Cruise 101C Leg 1, 05 June-13 Jul 1996, TOBI surveys of the continental slope west of Shetland.
- Masson, D.G. 1997. RRS Charles Darwin Cruise 101C Leg 1.
- Masson, D.G., Bett, B.J. & Birch, K.G. 1997. Atlantic margin environmental survey. *Sea Technology*, **38**, 52-59.
- Masson, D.G. 1998. The Atlantic frontier - the physical environment. *Atlantic Frontier Environmental Conference 6th & 7th October 1998 (Abstracts of Presentations)*. Atlantic Frontier Environmental Forum, A. (editor).
- Masson, D.G. & Jacobs, C.L. 1998. R.V. Colonel Templar Cruises 01 and 02/98: TOBI surveys of the continental slope north and west of Scotland. *Cruise Report*.
- Masson, D. & Bett, B. 1999. Atlantic margin environmental surveys. *AAPG Bulletin. AAPG international conference and exhibition; abstracts*.
- Masson, D.G. & Bett, B. 1999. Atlantic margin environmental surveys. *AAPG Bulletin*, **83**, Pt 8, 1328.
- Masson, D.G., Jacobs, C.L., Bas, T.P.L. & Huhnerbach, V. 2000. Section 4.1: Surficial Geology. *In: Network, A.F.E. (ed.). Environmental Surveys of the Seafloor of the UK Atlantic Margin [CD-ROM]*. Geotek Ltd, Daventry.
- Masson, D., Bett, B., Billett, D. & Wheeler, A. 2001. The Darwin mounds - possible fluid escape features in the Northern Rockall Trough. *Journal of Conference Abstracts*, **6**, 751.
- Masson, D.G., Wynn, R.B., Bett, B.J., Stoker, M.S. & Howe, J.A. 2001. Bottom current bedforms on the Northwest UK continental margin. 55 in *BSRG 2001, Plymouth, UK, 16-19 Dec 2001*. University of Plymouth.)
- Mather, A.S. 1979. *Beaches of southwest Scotland*. University of Aberdeen, Aberdeen.

Mauchline, J. 1979. Artificial radioisotopes in the marginal seas of northwestern Europe. *In: Banner, F.T., Collins, M.B. & Massie, K.S. (eds.). The North-West European shelf seas: the sea bed and the sea in motion - II. Physical and chemical oceanography, and physical resources.* Elsevier, Amsterdam, 517-542.

Mauchline, J., Ellett, D.J., Gage, J.D., Gordon, J.D.M. & Jones, E.J.W. 1986. A Bibliography of the Rockall Trough. *Proceedings of the Royal Society of Edinburgh*, **88B**, 319-354.

McCandliss, R.R. 2001. Distribution and dynamics of particulate matter at the Hebridean shelf edge. Unpublished thesis, University of Wales.

McCave, I.N. & Tucholke, B.E. 1986. Deep current-controlled sedimentation in the western North Atlantic. *In: Vogt, P.R. & Tucholke, B.E. (eds.). The western North Atlantic region.* Boulder, Colorado: Geological Society of America, 451-468.

McCave, I.N. & NEAPACC Investigators. 1998. North East Atlantic palaeoceanography: a climate change-driven system in the Late Quaternary. 108 in *GeoScience 98, (Keele University, 14-18 April 1998) Abstracts Volume.* Geological Society of London.)

McDonnell, A. & Shannon, P.M. 2001. Comparative Tertiary stratigraphic evolution of the Porcupine and Rockall basins. *In: Shannon, P.M., Haughton, P.D.W. & Corcoran, D.V. (eds.). The Petroleum Exploration of Ireland's Offshore Basins.* **188.** Geological Society of London, Special Publications, London, 323-344.

McGrane, K., Readman, P.W., O'Reilly, B.M., Jacob, A.W.B. & Keary, R. 1998. Gravity and side-scan sonar studies in the Rockall and Porcupine troughs, offshore Ireland. *Annales Geophysicae*, **16**, 284.

McGrane, K., Unnithan, V., Readman, P.W., Jacob, A.W.B., Keary, R. & Kenyon, N.H. 1999. Depositional and erosional processes in the Rockall Trough imaged by GLORIA. 56 in *North-East Atlantic Slope Processes: Multi-Disciplinary Approaches. Incorporating: TTR-8 Post Cruise Conference; 4th ENAM II Workshop; IGCP Workshop 432 - Contourites and Bottom Currents.* (SOC, Southampton, UK: SOC.)

McIntyre, A., Kipp, N.G., Be, A.W.H., Crowley, T., Gardner, J.V., Prell, W. & Ruddiman, W. 1976. Glacial North Atlantic 18,000 years ago: a CLIMAP reconstruction. *Memoirs of the Geological Society of America*, **145**, 43.

McKinley, I.G., Baxter, M.S., Ellett, D.J. & Jack, W. 1981. Tracer applications of radiocaesium in the sea of the Hebrides. *Estuarine, Coastal and Shelf Science*, **13**, 69-82.

McLean, A.C., Wren, A.E. & Walker, J.H.D. 1970. Gravity and magnetic and sparker surveys in the Firth of Clyde. *Proceedings of the Geologists' Association*, **166**, 75-76.

McLean, A.C. & Deegan, C.E. 1978. The solid geology of the Clyde Sheet (55°N/6°W).

McMahon, N.A. 1995. The role of uplifts in the sedimentation and rifting history of the North Atlantic. Unpublished thesis, University of Edinburgh.

- McManus, J. 1992. A Hydrographic Framework For Marine Conservation In Scotland. *Proceedings of the Royal Society Edinburgh*, **100B**, 3-26.
- McQuillin, R., Eden, R.A. & Tully, M.C. 1968. Geophysical and geological investigations in the Sea of Hebrides and Minches, 20th September - 3rd November, 1968. MGU Project No 68/05. *Marine Geophysics report*.
- McQuillin, R. & Binns, P.E. 1973. Geological Structure in the Sea of the Hebrides. *Nature; Physical Science (London)*, **241**, 2-4.
- McQuillin, R. & Watson, J. 1973. Large-scale Basement Structures of the Outer Hebrides in the Light of Geophysical Evidence. *Nature; Physical Science (London)*, **245**, 1-3.
- Megson, J.B. 1987. The evolution of the Rockall Trough and implications for the Faeroe Shetland Trough. In: Brooks, J. & Glennie, K.W. (eds.). *Petroleum Geology of North West Europe*. 653-665.
- Mienert, J. & Chi, J. 1995. Astronomical time-scale for physical property records from Quaternary sediments of the northern North Atlantic. In: Anon (ed.). *Geologische Rundschau. Global environmental change; the northern North Atlantic*. **84**. Springer International. Berlin, Federal Republic of Germany., 67-88.
- Mienert, J., Chi, J. & Baas, J. 1995. Linking physical property records of Quaternary sediments from the European Atlantic margin to major iceberg surges in the North Atlantic. In: Anon (ed.). *Global change and marine geology; 85. Jahrestagung der Geologischen Vereinigung e.V.; Zusammenfassung der Tagungsbeitraege*. **1-95**. Alfred-Wegener-Stiftung. Bonn, Federal Republic of Germany., 32-33.
- Mienert, J., Posewang, J. & Baumann, M. 1998. Gas hydrates along the northeastern Atlantic Margin; possible hydrate-bound margin instabilities and possible release of methane. In: Henriot, J.P. & Mienert, J. (eds.). *Geological Society Special Publications. Gas hydrates; relevance to world margin stability and climate change*. **137**. Geological Society of London, London, United Kingdom., 275-291.
- Mienert, J., Abrantes, F., Auffret, G.A., Evans, D., Kenyon, N.H., Kuijpers, A.H., Sejrup, H.P. & van Weering, T.C.E. 1998. European North Atlantic Margin (ENAM I): Sediment pathways, processes and fluxes - an introduction. *Marine Geology*, **152**, 3-6.
- Miller, K.G. & Tucholke, B.E. 1983. Development of Cenozoic abyssal circulation south of the Greenland-Scotland Ridge. In: Bott, M.H.P., Saxov, S., Talwani, M. & Thiede, J. (eds.). *Structure and development of the Greenland-Scotland Ridge; new methods and concepts*. **IV. Marine Sciences**. **8**. Plenum, New York and London, 549-589.
- Miller, D. 1983. A geotechnical study of Clyde alluvia. Unpublished thesis, University of Strathclyde.
- Miller, J.A. & Pye, K. 1987. Dating movement on geological faults. *Terra Cognita*, **7**, 335.
- Milliman, J.D. 1980. Coccolithophorid production and sedimentation, Rockall Bank. *Deep-Sea Research. Part A: Oceanographic Research Papers*, **27**, 959-963.

Mitchell, C.J. 1971. Analysis of tephra from borehole 88/7, 8/7A, Rockall - Faeroes trough.

Morgan, G.E. 1984. Palaeomagnetism. 38-39 in St Kilda: an illustrated account of the geology. *In*: Harding, R.R., Merriman, R.J. & Nancarrow, P.H.A. (eds.). **16, No. 7**. Report of the British Geological Survey.

Morgan, J.V. 1988. Seismic studies over continental margins. Unpublished thesis, University of Cambridge.

Morgan, J.V. & Barton, P.J. 1990. A geophysical study of the Hatton Bank volcanic margin: a summary of the results from a combined seismic, gravity and magnetic experiment. *Tectonophysics*, **173**, 517-526.

Morton, A.C. 1987. Distribution and significance of volcanic glass shards in vibrocore 59-07/276, Wyville-Thomson Ridge. pages 20.

Morton, A.C. 1988. Petrology and Geochemistry of igneous rock samples from Hebrides Terrace Seamount and Epsom Shoal, UK Continental Shelf.

Morton, A.C. & Parson, L.M. 1988. Early Tertiary Volcanism and the Opening of the NE Atlantic. *Geological Society of London Special Publication*, No. 39.

Morton, A.C. & Taylor, P.N. 1991. Geochemical and isotopic constraints on the nature and age of basement rocks from Rockall Bank, NE Atlantic. *Journal of the Geological Society of London*, **148**, 631-634.

Morton, A.C., Hitchen, K., Ritchie, J.D., Hine, N.H., Whitehouse, M. & Carter, S.G. 1995. Late Cretaceous basalts from Rosemary Bank, northern Rockall Trough. *Journal of the Geological Society of London*, **152**, 947-952.

Muir, R.J. 1991. The Precambrian basement and related rocks of the southern Inner Hebrides, Scotland. Unpublished Ph.D., E3d thesis, University of Aberystwyth, Wales.

Muir Wood, R. 1989. Fifty million years of 'passive margin' deformation in north west Europe. *In*: Gregerson, S. & Basham, P.W. (eds.). *Earthquakes at North-Atlantic Passive Margins: Neotectonics and Postglacial Rebound*. Kluwer Academic Publishers, Dordrecht, 7-36.

Mukhin, A.I. 1975. Decrease in water temperature in the Northeastern Atlantic. *Oceanology*, **14**, 821-823.

Murray, J., Hjort, J., Aooelloff, H.H. & Heland-Hansen, B. 1912. *The depths of the ocean*. Macmillan, London.

Murray, J.W. & Taplin, C.M. 1984. Larger agglutinated foraminifera from the Faeroe Channel and Rockall Trough collected by W.B. Carpenter. *Journal of Micropalaeontology*, **3**, 59-62.

Musgrove, F.W. & Mitchener, B. 1996. Analysis of the pre-Tertiary rifting history of the Rockall Trough. *Petroleum Geoscience*, **2**, 353-360.

Musson, R.M.W., Long, D., Pappin, J.W., Lubkowski, Z.A. & Booth, E. 1997. UK continental shelf seismic hazard.

Mykura, W. & Phemister, J. 1976. *Geology of offshore Ireland and West Britain*. Graham and Trotman, London.

Nadin, P.A., Houchen, M.A. & Kuszniir, N.J. 1999. Evidence for pre-Cretaceous rifting in the Rockall Trough: an analysis using quantitative 2D structural/stratigraphic modelling. In: Fleet, A.J. & Boldy, S.A.R. (eds.). *Petroleum Geology of Northwest Europe: Proceedings of the 5th Conference*. Geological Society of London, London, 371-378.

Naylor, D. & Mounteney, S.N. 1975. *Geology of the North-West European continental shelf*. Graham Trotman Dudley Publ. Ltd., London.

Naylor, D. & Shannon, P.M. 1982. *Geology of Offshore Ireland and West Britain*. Graham & Trotman, London.

Naylor, D., Shannon, P. & Murphy, N. 1999. Irish Rockall Basin region - a standard structural nomenclature system.

Neish, J.C.K. 1990. Rockall Trough: Crustal Structure of a Complex Environment from Composite Seismic Surveys. *Society of Economic Geologists, Abstracts*, **60**, 301-303.

Neish, J.C.K. 1992. Seismic structure of the Hatton-Rockall area: an integrated seismic/modelling study from composite data sets. *Petroleum Geology of NW Europe, Abstracts*, **131**.

New, A.L. & Smythe Wright, D. 2001. Aspects of circulation in the Rockall Trough. *Continental Shelf Research*, **21 Issues 8-10**, 777-810.

Nielson, T. 1998. Slope stability and sediment accumulations in the North Atlantic: Evidence of Cenozoic tectonic episodes. 83-84 in *Neogene Uplift and Tectonics around the North Atlantic - International workshop, Copenhagen, May 18-19 1998*. Boldreel, L. & Jaspén, P. (editors). Geological Survey of Denmark and Greenland (GEUS).

Nilsen, T.H. 1978. Sedimentation in the Northeast Atlantic Ocean and Norwegian Sea. In: Bowes, D.R. & Leake, B.E. (eds.). *Crustal evolution in northwestern Britain and adjacent regions*. (Geological Society of London Special Publications). **10**. Geological Society of London, London, 433-454.

NIO. 1970. RRS Discovery Cruise 29, August - October 1969: GLORIA in the Azores and geophysics on and around the Rockall Plateau. *NIO Cruise Report*.

NIO. 1971. M.V. Surveyor Cruise 71/1, February - April, 1971: Geology and geophysics on the Hebridean Shelf and on the Rockall Plateau. *NIO Cruise Report*.

NIO. 1971. RRS Discovery cruise 33, April - May 1970: DSDP site surveys and geology and geophysics around King's Trough. *NIO Cruise Report*.

NIO. 1971. RRS Discovery Cruise 39, April - June 1971: Plankton investigations at 60N, 20W and 53N, 20W. *NIO Cruise Report*.

NIO. 1972. RRS Discovery Cruise 42, 21st - 30th September 1971: Long range detection of herring by GLORIA. *NIO Cruise Report*.

Nolan, C. & Wilson, I. 1999. Northeast Irish Rockall; assessing geological risk in a deepwater, frontier province. *AAPG Bulletin. AAPG international conference and exhibition; abstracts*.

Noor Bin Salleh, M. 1975. Sediment balance of the Upper Clyde estuary. Unpublished thesis, University of Strathclyde.

Ollif, N.J. 1973. Bathymetric and Magnetic Traverses Over the Rockall Plateau: RRS Discovery, Cruise 47, June-July, 1972.

Omran, M.A. & Whittington, R.J. 1986. Geophysical Studies of the Hebrides Terrace Seamount. *Geological Society Newsletter*, **15**, 43.

Omran, M.A. 1990. Geophysical studies in the Hebrides Terrace Seamount area. Unpublished thesis, University of Wales.

O'Reilly, B.M., Readman, P.W., Shannon, P.M. & Jacob, A.W.B. 2002. A model for the development of a carbonate mound population in the Rockall trough based on deep-towed sidescan sonar data. *Marine Geology (in press)*.

Pain, S. 1985. Rockall; a controversy in deep water. *New Scientist*, **107**, 1464.

Palmer, T.J., McKerrow, W.S. & Cowie, J.W. 1980. Sedimentological evidence for a stratigraphical break in the Durness Group. *Nature*, **287**, 721-722.

Pantin, H.M. 1991. The seabed sediments around the United Kingdom: their bathymetric and physical environment, grain size, mineral composition and associated bedforms. *Offshore Geology Series, Research Report*, pages 47 (+ 10 charts).

Parker, J.G. 1982. Grain-size characteristics of Recent sediments in Belfast Lough. *Marine Geology*, **50**, Pt 1-2, 143-154.

Parra, M. 1982. North Atlantic sedimentation and paleohydrology during the late Quaternary mineralogical and geochemical data. *Oceanologica Acta*, **5**, 241-248.

Parra, M., Delmont, P., Ferrange, A., Latouche, C., Pons, J.C. & Puechmaille, C. 1985. Origin and Evolution of Smectites in Recent Marine Sediments of the NE Atlantic. *Clay Minerals*, **20**, 335-346.

Parson, L.M., Masson, D.G., Miles, P.R. & Pelton, C.D. 1986. Structure and evolution of the Rockall and East Greenland Continental Margins. Report of work undertaken by IOS in the period up to April 1985. *IOS Report*, pages 71.

Paul, M.A. & Jobson, L.M. 1987. On the Geotechnical and Acoustic Properties of Sediments from the British Continental Margin W of the Hebrides. 350.

Paul, M.A. & Talbot, L.A. 1991. Thematic geotechnical analysis of sediments from the Continental Slope Northwest of the British Isles.

Paul, M.A., Talbot, L.A. & Stoker, M.S. 1993. Geotechnical properties of sediments from the Continental Slope NW of the British Isles. *In: Ardu, D.A., Clare, D., Hill, A., Hobbs, R., Jardine, R.J. & Squire, J.M. (eds.). Advances in Underwater Technology. Offshore Site Investigation and Foundation Behaviour.* **28**. Kluwer Academic Publishers, 77-106.

Paul, M.A. & Talbot, L.A. 1995. Geotechnical and Seismic analysis of a deep sedimentary sequence on the UK Continental Margin.

Paul, M.A., Talbot, L.A. & Stoker, M.S. 1998. Shallow geotechnical profiles, acoustic character and depositional history of glacially influenced sediments from the Hebrides and west Shetland slopes. *In: Stoker, M.S., Evands, D. & Cramp, A. (eds.). Geological Processes on Continental Margins: Sedimentation, Mass-Wasting and Stability.* (Geological Society of London Special Publications). **129**. Geological Society of London, London, 117-131.

Payne, R. 1976. Flow across the Continental Slope off Northern Scotland. *Deep-Sea Research and Oceanographic Abstracts*, **23**, 875-880.

Peacock, J.D., Graham, D.K., Robinson, J.E. & Wilkinson, I. 1977. Evolution and chronology of Lateglacial marine environments at Lochgilphead, Scotland. *In: Gray, J.M. & Lowe, J.J. (eds.). Studies in the Scottish lateglacial environment.* Pergamon Press, Oxford, 89-100.

Peacock, J.B., Graham, D.K. & Wilkinson, I.P. 1978. Late glacial and post-glacial marine environments at Ardyne, Scotland, and their significance in the interpretation of the history of the Clyde Sea area. *IGS report*, pages 25.

Peacock, J.D. 1981. Scottish late-glacial marine deposits and their environmental significance. *In: Neale, J.W. & Henley, J.R. (eds.). The Quaternary in Britain.* Pergamon Press, Oxford, 222-236.

Peacock, J.D. 1981. Glacial deposits of the Hebridean region. *In: Ehlers, J., Gibbard, P.L. & Rose, J. (eds.). Glacial Deposits in Great Britain and Ireland.* A A Balkema, Rotterdam.

Peacock, J.D. 1984. Quaternary Geology of the Outer Hebrides.

Peacock, J.D. & Harkness, D.D. 1990. Radiocarbon ages and the full-glacial to Holocene transition in Seas adjacent to Scotland and Southern Scandinavia: a review. *Transactions of the Royal Society of Edinburgh: Earth Sciences*, **81**, 385-396.

Peacock, J.D. 1991. Glacial deposits of the Hebridean region. *In: Ehlers, J., Gibbard, P.L. & Rose, J. (eds.). Glacial deposits in Great Britain and Ireland.* 109-119.

Peacock, J.D., Austin, W.E.N., Selby, I., Graham, D.K., Harland, R. & Wilkinson, I.P. 1992. Late Devensian and Flandrian Palaeoenvironmental changes on the Scottish Continental Shelf West of the Outer Hebrides. *Journal of Quaternary Science*, **7**, 145-161.

Peacock, J.D. 1993. Late Quaternary marine mollusca as palaeoenvironmental proxies: a compilation and assessment of basic numerical data for NE Atlantic species found in shallow water. *Quaternary Science Reviews.*, **12**, 263-275.

Pearce, G.J. 1980. Benthonic Foraminifera from the Continental Slope and their Fossil Counterparts. Unpublished thesis, University of Wales.

Pendlebury, D.C. 1974-75. Recent sediments, shelly fauna and foraminifera of the Malin Sea. Unpublished thesis, University of Aberystwyth, Wales,.

Pendlebury, D.C. & Dobson, M.R. 1976. Sediments and microfaunal distributions in the eastern Malin Sea, as determined by sidescan sonar and sampling. *Scottish Journal of Geology*, **11**, 315-332.

Perry, R.G. 1987. A geological and geophysical investigation of the Hebridean continental margin. Unpublished thesis, University of London.

Peypouquet, J.P. 1975. Le renouvellement de la faune d'Ostracodes du bassin de Rockall entre le Miocene et le Pleistocene superieur; interet paleohydrologique
Translated Title: The revival of the ostracod fauna in the Rockall Basin between the Miocene and upper Pleistocene; pa. *Bulletin de la Societe Geologique de France*, **17**, 886-895.

Pichon, X.L., Cressard, A., Mascle, J., Pautot, G. & Sichler, B. 1970. Structures sous-marines des bassins sedimentaires de Porcupine et de Rockall; Translated Title: Submarine structures of the Porcupine and Rockall sedimentary basins. *Comptes Rendus Hebdomadaires des Seances de l'Academie des Sciences, Serie D: Sciences Naturelles*, **270**, 2903-2906.

Picken, M.J. 1979. Report on the environmental factors relating to sites for the bottom standing structures west of the Outer Hebrides. N. E. L. Oscillating Water Column.

Picken, M.J. 1980. Interim Report on marine fouling assessments around the Outer Hebrides and preliminary topography survey of the wave energy area.

Pillai, M.H. 1998. Flow and water column structure at the Hebridean shelf-edge. Unpublished thesis, University of Wales.

Pitcher, W.S. 1969. Northeast trending faults of Scotland and Ireland, and chronology of displacements. In: Kay, M. (ed.). *North Atlantic; geology and continental drift*. **12**. American Association of Petroleum Geologists, Tulsa, OK, United States, 724-733.

Pnencz-Castillo, F., Ezzi, I., Jones, K., Lampitt, R. & Tett, P. 1996. Particulate organic matter in waters on the Hebrides continental slope: distribution and sedimentation patterns. 179 in *UK Oceanography*. Anon (editor). (University of Wales, Bangor:

Powell, A.D.J. 1988. Palaeobathymetric analysis of Tertiary sediments in the Northern North Sea and north-east Atlantic Ocean. Unpublished thesis, University of Exeter.

Praeg, D., Haughton, P.D.W., O'Reilly, B.M., Øvrebø, L.K. & Shannon, P.M. 2001. History of a Slippery Slope: Neogene to Quaternary Sedimentary Processes on the Eastern Rockall Margin. In: Murphy, N.J. & Davies, M. (eds.). *Ireland's Deepwater*

Frontier: Results from the Petroleum Infrastructure Programme (PIP). Extended Abstracts. PIP, Dublin, 42-44.

Prescott, C.N. 1988. Marine geophysical investigation of the Hatton Bank volcanic passive continental margin. Unpublished thesis, University of Durham.

Proctor, R. & Davies, A.M. 1996. A three dimensional hydrodynamical model of tides off the north-west coast of Scotland [the Malin-Hebrides shelf]. *J Mar Sys [Amsterdam]*, **7/1**, 43-66.

Rae, J.B., Elford, W.E. & Cartwright, D.E. 1977. RRS "Discovery" Cruise 78 (3 legs), 8 September - 18 October 1976. Oceanic tidal recording SE and SW of Iceland and tests and use of side-scan sonar west of Hebrides. *IOS Cruise Report*, pages 22.

Rashid, B.M. 1978. Interpretation of geophysical data in the Firth of Lorne. Unpublished thesis, University of Glasgow.

Rasmussen, T.L., van Weering, T.C.E. & Labeyrie, L. 1997. Climatic instability, ice sheets and ocean dynamics at high northern latitudes during the last glacial period (58-10 ka bp). *Quaternary Science Review*, **16**, 71-80.

Rasmussen, T.L., Backstrom, D.L., Heinemeier, J., Klitgaard Kristensen, D., Knutz, P.C., Kuijpers, A., Lassen, S., Thomsen, E., Troelstra, S.R. & van Weering, T.C.E. 2002. The Faroe-Shetland Gateway: Late Quaternary water mass exchange between the Nordic seas and the northeastern Atlantic. *Marine Geology*, **188**, 165-192.

Raymo, M.E. & Ruddiman, W.F. 1992. Tectonic forcing of late Cenozoic climate. *Nature*, **359**, 117-122.

Readman, P.W., O'Reilly, B.M., Shannon, P.M., Jacob, A.W.B., Kenyon, N. & Blake, T. 2001. Images of slope failure along the glaciated eastern margin of the Rockall Trough from TOBI sidescan sonar. In: Murphy, N.J. & Davies, M. (eds.). *Ireland's Deepwater Frontier: Results from the Petroleum Infrastructure Programme (PIP). Extended Abstracts.* PIP, Dublin, 90-93.

Rees, A.I., Brown, C.M., Hailwood, E.A. & Riddy, P.J. 1982. Magnetic Fabric of Bioturbated Sediment from the Northern Rockall Trough; Comparison with modern currents. *Marine Geology*, **46**, 161-173.

Richards, P.C., Ritchie, J.D. & Thomson, A.R. 1987. Evolution of deep-water climbing dunes in the Rockall Trough; implications for overflow currents across the Wyville-Thomson Ridge in the (?)late Miocene. *Marine Geology*, **76**, Pt 3-4, 177-183.

Richardson, A.E. 1999. Western Frontiers Association Bibliography User Guide Version 4.0.

Richter, T.O., Lassen, S., van Weering, T.C.E. & de Haas, H. 2001. Magnetic susceptibility patterns and provenance of ice-rafted material at Feni Drift, Rockall Trough; implications for the history of the British-Irish ice sheet. *Marine Geology*, **173**, Pt 1-4, 37-54.

Ridley, G. 1980. The British Sub-Aqua Club St. Kilda survey expedition. *Underwater World*, **3**, 12-15.

Ritchie, W. 1966. The post-glacial rise in sea level and coastal changes in the Uists. *Transactions of the Institute of British Geographers*, **39**, 79-86.

Ritchie, W. & Mather, A.S. 1970. The beaches of Lewis and Harris: A survey of the beach, dune and machir areas of Lewis and Harris.

Ritchie, W. & Whittington, G. 1994. Nonsynchronous Aeolian Sand Movements In The Uists - The Evidence Of The Intertidal Organic And Sand Deposits At Cladach-Mor, North-Uist. *Scottish Geographical Magazine*, **110**, 40-46.

Ritchie, J.D., Gatliff, R.W. & Richards, P.C. 1999. Early Tertiary magmatism in the offshore NW UK margin and surrounds. In: Fleet, A.J. & Boldy, S.A.R. (eds.). *Petroleum Geology of Northwest Europe: Proceedings of the 5th Conference*. 573-584.

Ritchie, W., Whittington, G. & Edwards, K.J. 2001. Holocene changes in the physiography and vegetation of the Atlantic littoral of the Uists, Outer Hebrides, Scotland. *Transactions Of The Royal Society Of Edinburgh-Earth Sciences*, **92**, 121.

Roberts, D.G. 1969. New Tertiary volcanic centre on the Rockall bank, eastern north Atlantic Ocean. *Nature*, **223**, 819-820.

Roberts, D.G., Bishop, D.G., Laughton, A.S., Ziolkowski, A.M. & Scrutton, R.A. 1970. New sedimentary basin on Rockall Plateau. *Nature*, **225**, 170-172.

Roberts, D.G. 1971. New geophysical evidence on the origin of the Rockall Plateau and Trough. *Deep-Sea Research*, **18**, 350-360.

Roberts, D.G. 1972. Site survey in Hatton-Rockall Basin (sites 116 and 117). In: Laughton, A.S. (ed.). *Initial Reports of the Deep Sea Drilling Project*. **12**. US Government Printing Office, Washington D.C, USA, 1201-1208.

Roberts, D.G. 1973. The Overflow of Norwegian Sea Deep Water across the Wyville-Thomson Ridge. *Deep-Sea Research and Oceanographic Abstracts*, **20**, 819-835.

Roberts, D.G., Hogg, N.G., Bishop, D.G. & Flewellen, C.G. 1974. Sediment distribution around moated seamounts in the Rockall Trough. *Deep-Sea Research and Oceanographic Abstracts*, **21**, 175-184.

Roberts, D.G. 1974. Structural development of the British Isles, the continental margin, and the Rockall Plateau. In: Burk, C.A. & Drake, C.L. (eds.). *The geology of continental margins*. Springer-Verlag, New York, 343-359.

Roberts, D.G., Fleming, N.C., Harrison, R.K. & Binns, P.E. 1974. Helen's Reef: a Cretaceous microgabbroic intrusion in the Rockall intrusive centre. *Marine Geology*, **16**, M21-M30.

Roberts, D.G. & Eden, R.A. 1974. DE "Vickers Voyager" and "Pisces III" June - July 1973. Submersible investigations of the geology and benthos of the Rockall Bank. *Cruise Report*, pages 22.

Roberts, D.G. 1974. RRS "Discovery" Cruise 60, 26 February - 22 March 1974. Seismic reflection profiling on the Rockall Plateau and in the South Western approaches. *Cruise Report*, pages 17.

Roberts, D.G. 1975. Sediment distribution on the Rockall Bank, Rockall Plateau. *Marine Geology*, **19**, 239-257.

Roberts, D.G. 1975. Marine Geology of the Rockall Plateau. *Philosophical Transactions of the Royal Society of London*, **278**, 447-509.

Roberts, D.G. 1975. The solid geology of the Rockall Plateau. *In: Harrison, R.K. (ed.). Expeditions to Rockall 1971-72*. Report of the Institute of Geological Sciences, No. 75/1.

Roberts, D.G. 1975. Tectonic and stratigraphic evolution of the Rockall Plateau and trough. *In: Woodland, A.W. (ed.). Petroleum and the continental shelf of North-west Europe; geology*. John Wiley & Sons, New York, 77-91.

Roberts, D.G. 1975. Geology and tectonics of the area beyond the shelf west of the British Isles. *In: Offshore Europe '75. Paper OE-75 202; 8 pages*. Spearhead publications, Kingston-upon-Thames, UK.

Roberts, D.G., Hunter, P.M. & Laughton, A.S. 1977. Continental margin around the British Isles 1:2 400 000. Hydrographer of the Navy for Institute of Oceanographic Sciences, Taunton.

Roberts, D.G., Hunter, P.M. & Laughton, A.S. 1979. Bathymetry of the northeast Atlantic: continental margin around the British Isles. *Deep-Sea Research*, **26A**, 417-428.

Roberts, D.G. & Kidd, R. 1979. Abyssal sediment wave fields on the Feni Ridge, Rockall Trough: Long range sonar studies. *Marine Geology*, **33**, 175-191.

Roberts, D.G., Montadert, L. & Searle, R.C. 1979. The western Rockall Plateau : stratigraphy and structural evolution. *Initial Reports of the Deep Sea Drilling Project*, **48**, 1061-1088.

Roberts, D.G. 1980. Geological setting and principal results of drilling on the margins of the Bay of Biscay and Rockall Plateau during Leg 48. *Philosophical Transactions of the Royal Society of London*, **294A**, 65-75.

Roberts, D.G., Masson, D.G. & Miles, P.R. 1981. Age and structure of the southern Rockall Trough; new evidence. *Earth and Planetary Science Letters*, **52**, 115-128.

Roberts, D.G., Bott, M.H.P. & Uruski, C. 1983. Structure and origin of the Wyville-Thomson Ridge. *In: Bott, M.H.P., Saxov, S., Talwani, M. & Thiede, J. (eds.). Structure and development of the Greenland-Scotland Ridge; new methods and concepts*. (NATO Conference Series IV: Marine Sciences). **8**. Plenum, New York, 133-158.

Roberts, D.G. 1989. Basin inversion in and around the British Isles. *In: Cooper, M.A. & Williams, D.G. (eds.). Inversion Tectonics*. Geological Society of London Special Publication, No. 44, London, 131-150.

- Roberts, J.M., Chenery, S.R., Long, D., Gage, J.D., Wilson, J.B. & Mortensen, P.B. 1999. An interdisciplinary study of *Lophelia pertusa*: west of Shetland distribution and initial skeletal isotope and trace element analysis. 68 in *North East Atlantic Slope Processes: Multi-Disciplinary Approaches, Abstract Book. Conference 24-27th January 1999*. Friend, P. & Kenyon, N. (editors). Southampton Oceanography Centre, Southampton, UK.)
- Roberts, D.G., Thompson, M., Mitchener, B., Hossack, J., Carmichael, S. & Bjørnseth, H.M. 1999. Palaeozoic to Tertiary rift and basin dynamics: mid-Norway to the Bay of Biscay - a new context for hydrocarbon prospectivity in the deep water frontier. In: Fleet, A.J. & Boldy, S.A.R. (eds.). *Petroleum Geology of Northwest Europe: Proceedings of the 5th Conference*. Geological Society of London, London, 7-40.
- Roberts, J.M., Harvey, S.M., Lamont, P.A., Gage, J.D. & Humphery, J.D. 2000. Seabed photography, environmental assessment and evidence of deep-water trawling on the continental margin west of the Hebrides. *Hydrobiologia*, **441**, 173-183.
- Robinson, S.G. 1994. Orbital forcing of bottom-current enhanced sedimentation on Feni Drift, NE Atlantic, during the mid-Pleistocene. *Palaeoceanography*, **9**, 943-972.
- Robinson, S.G. & McCave, I.N. 1994. Orbital forcing of bottom-current enhanced sedimentation on Feni Drift, NE Atlantic, during the mid-Pleistocene. *Paleoceanography*, **9**, 943-972.
- Robinson, S.G., Maslin, M.A. & McCave, I.N. 1995. Magnetic susceptibility variations in Upper Pleistocene deep-sea sediments of the NE Atlantic: Implications for ice rafting and paleocirculation at the last glacial maximum. *Palaeoceanography*, **10**, 221-250.
- Rogalla., E. 1961. Results of Hydrographic investigations between St.Kilda and Rockall Bank in April 1959. *Annales Biologiques*, **16**, 65-66.
- Ruckley, N.A. & Chesher, J.A. 1987. Caithness, Sheet 58N - 04W, 1:250 000 Map Series, Sea-bed Sediments and Quaternary Geology. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.
- Ruckley, N.A. & Evans, D. 1988. Little Minch, Sheet 57N - 08W including part of the Great Glen 57N - 06W, 1:250 000 Map Series, Sea-bed Sediments and Quaternary Geology. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.
- Ruddiman, W.F., Sancetta, C.D. & McIntyre, A. 1977. Glacial/interglacial response rate of subpolar North Atlantic waters to climatic change; the record in oceanic sediments. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, **280**, 119-142.
- Ruddiman, W.F. & Kidd, R.B. 1986. *Initial Reports of the Deep Sea Drilling Project*. US Government Printing Office, Washington DC.
- Ruddiman, W.F. & Kidd, R.B. 1987. Site 610. In: Ruddiman, W.F., Kidd, R.B. & Thomas, E. (eds.). *Initial Report of the Deep Sea Drilling Project*. **94**. US Government Printing Office, Washington, 351-470.

Rushby, J.S.M. & Revie, J. 1975. Long range sonar mapping of the continental shelf. *Marine Geology*, **19**, M41-M52.

Russell, M.J. & Smythe, D.K. 1978. Evidence for an early Permian oceanic rift in the north North Atlantic. *In: Neumann, E.R. & Ramberg, I.B. (eds.). Petrology and geochemistry of continental rifts.* Dordrecht: Reidel Publishing Company, 173-179.

Stoker, M.S. & Fannin, N.G.T. 1984. A Geological Framework for the North West United Kingdom Continental Shelf and Slope. pages 15.

Stoker, M.S., Long, D. & Fyfe, J.A. 1985. A revised Quaternary stratigraphy for the central North Sea.

Stoker, M.S. 1988. Pleistocene ice-proximal glaciomarine sediments in boreholes from the Hebrides Shelf and Wyville-Thomson Ridge, NW UK Continental Shelf. *Scottish Journal of Geology*, **24**, 249-262.

Stoker, M.S. 1988. Sula Sgeir, Sheet 59N - 08W, 1:250 000 Map Series, Sea-bed Sediments. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.

Stoker, M.S. 1988. Pleistocene ice-proximal glaciomarine sediments in boreholes from the Hebrides shelf and Wyville-Thomson Ridge, NW UK continental shelf. *Scottish Journal of Geology*, **24**, Pt 3, 249-262.

Stoker, M.S., Morton, A.C., Evans, D., Hughes, M.J., Harland, R. & Graham, D.K. 1988. Early Tertiary basalts and tuffaceous sandstones from the Hebrides Shelf and Wyville-Thomson Ridge, NE Atlantic. *In: Morton, A.C. & Parson, L.M. (eds.). Early Tertiary volcanism and the opening of the Atlantic.* (Geological Society of London Special Publications). **39**. Geological Society of London, London, 271-282.

Stoker, M.S. and Hitchen, K. 1988. Sula Sgeir, Sheet 59N - 08W, 1:250 000 Map Series, Solid Geology. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.

Stoker, M.S. 1989. British Geological Survey Shallow Drilling Programme 1988.

Stoker, M.S., Harland, R., Morton, A.C. & Graham, D.K. 1989. Late Quaternary stratigraphy of the northern Rockall Trough and Faeroe-Shetland Channel, Northeast Atlantic Ocean. *Journal of Quaternary Science*, **4**, 211-222.

Stoker, M.S. 1990. Judd, Sheet 60N - 06W, 1:250 000 Map Series, Quaternary Geology. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.

Stoker, M.S. 1990. Sula Sgeir, 59N - 08W, 1:250 000 Map Series, Quaternary Geology. Ordnance Survey, Southampton for the British Geological Survey, Natural Environment Research Council.

Stoker, M.S. 1990. Glacially-influenced sedimentation on the Hebridean slope, northwestern United Kingdom continental margin. *In: Dowdeswell, J.A. & Scourse, J.D.*

(eds.). *Glacimarine environments: processes and sediments*. (Geological society of London Special Publications). **53**. Geological Society of London, London, 349-362.

Stoker, M.S., Abraham, D.A., Evans, D., Long, D., Ardu, D.A. & Dobinson, A.D. 1990. The Rockall Continental Margin: a review of existing data and recommendations for a 10-year regional survey programme.

Stoker, M.S. 1991. Glacially-influenced basin plain sedimentation, north-west United Kingdom passive continental margin. A119 in *Program with Abstracts - Geological Association of Canada; Mineralogical Association of Canada; Canadian Geophysical Union, Joint Annual Meeting*.

Stoker, M.S., Harland, R. & Graham, D.K. 1991. Glacially influenced basin plain sedimentation in the Southern Faeroe-Shetland Channel, north-west United Kingdom continental margin. *Marine Geology*, **100**, 185-199.

Stoker, M.S. & Holmes, R. 1991. Submarine end-moraines as indicators of Pleistocene ice-limits off northwest Britain. *Journal of the Geological Society of London*, **148**, 431-434.

Stoker, M.S., Leslie, A.B., Scott, W.D., Briden, J.C., Hine, N.H., Harland, R., Wilkinson, I.P., Mitchell, C.J., Kroon, D., Maher, B., Evans, D. & Ardu, D.A. 1992. A Multidisciplinary study of Mid-Tertiary to Quaternary sediments recovered in BGS borehole 88/7, 7A, Hebrides Slope, Northern Rockall Trough Region. *BGS Technical Report*

Stoker, M.S., Stewart, F.S., Paul, M.A. & Long, D. 1992. Problems associated with seismic facies analysis of Quaternary sediments on the northern UK continental margin. *Journal of Underwater Technology*, **18**, 3-15.

Stoker, M.S., Stewart, F.S., Paul, M.A. and Long, D. 1993. Problems associated with seismic facies analysis of Quaternary sediments on the Northern UK Continental Margin. In: Ardu, D.A., Clare, D., Hill, A., Hobbs, R. & Jardine, R.J.a.S., J.M. (eds.). *Advances in Underwater Technology. Offshore Site Investigation and Foundation Behaviour*. **28**. Kluwer Academic Publishers., 239-262.

Stoker, M.S. 1993. The geology of the Hebrides and West Shetland shelves, and adjacent deep water areas. *British Geological Survey United Kingdom Offshore Regional Report*, pages 149.

Stoker, M.S., Hitchen, K. & Graham, C. 1993. The Geology of the Hebrides and West Shetland Shelves and adjacent deep-water areas. In: Abrahams, D.A. & Ritchie, J.D. (eds.). HMSO for the British Geological Survey, 149pp.

Stoker, M.S., Leslie, A.B., Scott, W.D., Briden, J.C., Hine, N.M., Harland, R., Wilkinson, I.P., Evans, D. & Ardu, D.A. 1994. A record of late Cenozoic stratigraphy, sedimentation and climate change from the Hebrides Slope, NE Atlantic Ocean. *Journal of the Geological Society of London*, **151**, 235-249.

Stoker, M.S. 1995. Glacially-influenced basin-plain sedimentation in the Southern Faeroe-Shetland Channel, Northwest UK passive Continental Margin. *1st Nordic Marine Sciences Meeting. Goeteborg 23-26 Mars. 1995*.

- Stoker, M.S. 1995. The influence of Glacigenic Sedimentation on Slope-Apron development on the Continental Margin off NW Britain. *In: Scrutton, R.S., Stoker, M.S., Shimmiel, G.B. & Tudhope, A.W. (eds.). Geological Society Special Publication. The Tectonics, Sedimentation and Palaeoceanography of the North Atlantic Region. 90.* Geological Society of London, London, 159-177.
- Stoker, M.S. 1996. Sediment-drift development in the northern Rockall Trough, NW Britain. 65 in *Applied Geoscience conference Warwick University 15-18 April 1996.* Geological Society, London.)
- Stoker, M.S. 1996. Glacially-influenced basin-plain sedimentation in the southern Faeroe-Shetland Channel, NW United Kingdom passive continental margin. 72 in *Applied Geoscience conference Warwick University 15-18 April 1996.* Geological Society, London.)
- Stoker, M.S. & Gillespie, E.J. 1996. Sequence Stratigraphy and the Prediction of Sand-Prone Depositional Packages in the Neogene-Quaternary Succession of the West Shetland Margin.
- Stoker, M.S. 1997. Submarine End-Moraines on the West Shetland Shelf, North-West Britain. *In: Davies, T.A., Bell, T., Cooper, A.K., Josenhans, H., Polyak, L.S., A. & Stoker, M.S. and Stravers, J A. (eds.). Glaciated Continental Margins: an atlas of acoustic images.* Chapman and Hall, London. 84-85.
- Stoker, M.S. 1997. Submarine Debris Flows on a Glacially-Influenced Basin Plain, Faeroe-Shetland Channel. *In: Davies, T.A., Bell, T., Cooper, A.K., Josenhans, H., Polyak, L., Solheim, A., Stoker, M.S. & Stravers, J.A. (eds.). Glaciated Continental Margins: an atlas of acoustic images.* Chapman and Hall, London, 126-127.
- Stoker, M.S. 1997. Seismic-stratigraphic record of glaciation on the Hebridean margin, north-west Britain. *In: Davies, T.A., Bell, T., Cooper, A.K., Josenhans, H., Polyak, L., Solheim, A., Stoker, M.S. & Stravers, J.A. (eds.). Glaciated Continental Margins: An Atlas of Acoustic Images.* Chapman & hall, London, 264-267.
- Stoker, M.S. 1997. Mid- to late Cenozoic sedimentation on the continental margin off NW Britain. *Journal of the Geological Society of London, 154,* 509-515.
- Stoker, M.S. and Howe, J A. 1997. Glacially-Influenced Sediment Drifts in the Rockall Trough. *In: Davies, T.A., Bell, T., Cooper, A.K., Josenhans, H., Polyak, L. Solheim, A., Stoker, M.S. and Stravers, J A. (eds.). Glaciated Continental Margins: An Atlas of Acoustic Images.* Chapman and Hall, London., 290-293.
- Stoker, M.S. 1998. Reconstructing Quaternary palaeo-environments on the continental margin of NW Britain. 150 in *GeoScience 98, (Keele University, 14-18 April 1998) Abstracts Volume.* Geological Society of London.)
- Stoker, M.S. 1998. Sediment-drift development on the continental margin off NW Britain. *In: Stoker, M.S., Evans, D. & Cramp, A. (eds.). Geological Processes on Continental Margins: Sedimentation, Mass-Wasting and Stability.* (Geological Society of London Special Publications). **129.** Geological Society of London, London, 229-254.

- Stoker, M.S., Akhurst, M.C., Howe, J.A. & Stow, D.A.V. 1998. Sediment drifts and contourites on the continental margin off northwest Britain. *Sedimentary Geology*, **115**, 35-51.
- Stoker, M.S., Egerton, P., Holmes, R. & Long, D. 1998. Summary of shallow geology west of the Hebrides - notes of presentations given to the Western Frontier Association members, Aberdeen 24th June 1997, and London, 25th June 1997. *Offshore Technology Report*.
- Stoker, M.S., Evans, D. & Cramp, A. (eds.). 1998. *Geological Processes on Continental Margins: Sedimentation, Mass-Wasting and Stability*. (Geological Society of London Special Publications). Geological Society of London, London.
- Stoker, M.S. and Stow, D.A.V. 1998. Seismic-stratigraphic expression of a sediment-drift complex, Rockall Trough. 10 in *GeoScience 98, (Keele University, 14-18 April 1998) Abstracts Volume*. Geological Society of London.)
- Stoker, M.S., van Weering, Tj C E. and Svaerdborg, T. 1999. Towards a unified mid- to late Cenozoic stratigraphic framework for the Rockall Trough. 78 in *North East Atlantic Slope Processes: Multi-Disciplinary Approaches, Abstract Book. Conference 24-27th January 1999*. Friend, P.a.K., N. (editor). Southampton Oceanography Centre, Southampton, UK.)
- Stoker, M.S. (ed.). 1999. *Stratigraphic nomenclature of the UK North West Margin. 3. Mid- to late Cenozoic stratigraphy*. British Geological Survey, Edinburgh.
- Stoker, M.S., Evans, D. & Gillespie, E.J. 1999. A comparison of seismic interpretations on the UK and Faroese flanks in the Faroe-Shetland Channel.
- Stoker, M.S., Nielsen, T., van Weering, Tj C E. and Kuijpers, A. 2001. Towards an understanding of the Neogene tectonostratigraphic framework of the NE Atlantic margin between Ireland and the Faeroe Islands. In Press. *Marine Geology*.
- Stoker, M.S., van Weering, T.C.E. & Svaerdborg, T. 2001. A mid- to late Cenozoic tectonostratigraphic framework for the Rockall Trough. In: Shannon, P.M., Haughton, P. & Corcoran, D. (eds.). *The Petroleum Exploration of Ireland's Offshore Basins*. **188**. Geological Society of London, London, 411-438.
- Stoker, M.S. 2002. Growth and partial destruction of a Neogene sediment drift, western Rockall Trough. In: Meinert, J. & Weaver, P.P.E. (eds.). *European Margin Sediment Dynamics: Sidescan Sonar and Seismic Images*. Springer-Verlag, New York.
- Stoker, M.S. 2002. Central Rockall Basin (56°-58°N, 8°-15°W). Solid Geology: 1:500,000. British Geological Survey and Petroleum Affairs Division, Ireland.
- Stoker, M.S. 2002. Late Neogene development of the UK Atlantic margin. In: Doré, A.G.D., Cartwright, J., Stoker, M.S., Turner, J.P. & White, N. (eds.). *Exhumation of the North Atlantic Margin: Timing, Mechanisms and Implications for Petroleum Exploration*. Geological Society of London, Special Publications, 196, London.
- Stoker, M.S. & Hitchen, K. 2002. The Rockall-Porcupine Margin. In: Mienert, J. & Weaver, P. (eds.). *ENAM/STEAM Atlas of Seismic Images*. Springer, London.

Stoker, M.S., Nielsen, T., van Weering, T.C.E. & Kuijpers, A. 2002. Towards an understanding of the Neogene tectonostratigraphic framework of the NE Atlantic margin between Ireland and the Faroe Islands. *Marine Geology*.

Stoker, M.S. & STRATAGEM partners. 2002. The Neogene stratigraphy of the glaciated European margin from Lofoten to Porcupine. A product of the EC-supported STRATAGEM Project. World Wide Web Address: <http://www.stratagem-europe.org>.

Stoker, M.S. in press. *United Kingdom offshore regional report: the geology of the north-west margin of Scotland*. HMSO for the British Geological Survey, London.

Stow, D.A.V. & Holbrook, J.A. 1984. North Atlantic contourites: an overview. In: Stow, D.A.V. & Piper, D.J.W. (eds.). *Geological Society Special Publication No. 15. Fine-Grained Sediments: Deep Water Processes and Facies*. 245-256.

Stow, D.A.V. & Holbrook, J.A. 1984. Hatton Drift contourites, Northeast Atlantic, Deep Sea Drilling Project Leg 81. In: *Initial reports of the Deep Sea Drilling Project covering Leg 81 of the cruises of the drilling vessel Glomar Challenger; Southampton, United Kingdom, to Ponta Delgada, Azores Islands, July-September, 1981*. **81**; Texas A & M University Ocean Drilling Program, College Station, TX, United States, 695-699.

Stow, D.A.V. & Holbrook, J.A. 1984. Hatton Drift contourites, northwest Atlantic, Deep Sea Drilling Project Leg 81. *Initial Report of the Deep Sea Drilling Project*, **81**, 695-699.

Strachan, P. 1987. Geotechnical data from vibrocores and gravity cores; Geikie and Lewis sheets.

Strachan, P. & Evans, D. 1991. A local deep water sediment failure on the NW slope of the UK. *Scottish Journal of Geology*, **27**, 107-112.

STRATAGEM Partners. 2002. *The Neogene Stratigraphy of the glaciated European margin from Lofoten to Porcupine*.

Stride, A.H., Curray, J.R., Moore, D.G. & Belderson, R.H. 1969. Marine geology of the Atlantic continental margin of Europe. *Philosophical Transactions of the Royal Society of London, Series A: Mathematical and Physical Sciences*, **264**, 31-75.

Suess, E., Kremling, K. & Mienert, J. 1994. Nordatlantik 1993, Cruise No. 26. 24 August - 26 November. *Meteor-Berichte, Universitat Hamburg*, **94/4**, 1-256.

Sutherland, D.G. 1981. The raised shorelines and deglaciation of the Loch Long/Loch Fyne area, western Scotland. Unpublished thesis, University of Edinburgh.

Sutherland, D.G. 1984. The Quaternary deposits and landforms of Scotland and the neighbouring shelves: a review. *Quaternary Science Reviews*, **3**, 157-254.

Sutherland, D.G. 1984. The submerged landforms of the St Kilda archipelago, western Scotland. *Marine Geology*, **58**, 435-442.

Sutherland, D.G. 1987. Submerged rock platforms on the Continental Shelf West of Sula Sgeir. *Scottish Journal of Geology*, **23**, 251-260.

Sutherland, M.W. 1996. User Guide to the Clathrates Project for the Western Frontiers Association.

Svaerdborg, T. 1998. *A study of the Eocene to Recent deposits in the Rockall Trough, continental margin of the northeast Atlantic Ocean. Processing and interpretation of reflection seismic data.* M.Sc. thesis University of Aarhus,

Talbot, L.A. 1992. Some aspects of the sedimentological and geotechnical properties of sediments from the Continental Margin West of Shetland. Unpublished thesis, University of Heriot Watt.

Talbot, L.A., Paul, M.A. & Stoker, M.S. 1994. Geotechnical studies of a Plio-Pleistocene sedimentary sequence from the Hebrides Slope, NW UK Continental Margin. *Geo-Marine Letters*, **14**, 244-251.

Talbot, L.A., Paul, M.A. & Stoker, M.S. 1994. Geotechnical and acoustic characteristics of Plio-Pleistocene sediments from the Hebrides slope. *Geo-Marine Letters*, **14**, 244-251.

Talbot, L.A., Paul, M A., and Stoker, M S. 1996. Geotechnical properties of seabed sediments from the continental margin off northwest Britain. 67 in *Applied Geoscience conference Warwick University 15-18 April 1996*. Geological Society, London.)

Taylor, G.R. 1988. *The geotechnical characterisation of sediment from the north western part of the British continental margin with specific reference to the stability of the continental slope.* (unpublished B.Eng thesis) Heriot-Watt University, Edinburgh,

Thomas, B.W., Miller, J.M. & Malcolm, A. 1984. Radiometric surveys of the seabed. *Oceanology International Conference*. (Brighton: Proceedings of the Oceanology International '84 Conference, Paper No. 10, 11.)

Thomsen, L. & Gust, G. 2000. Sediment erosion thresholds and characteristics of resuspended aggregates on the western European continental margin. *Deep Sea Research Part 1: Oceanographic Research Papers*, **47 Issue 10**, 1881-1897.

Thomson, J., Brown, L., Nixon, S., Cook, G.T. & MacKenzie, A.B. 2000. Bioturbation and Holocene sediment accumulation fluxes in the North-east Atlantic Ocean (Benthic Boundary Layer experiment sites). *Marine Geology*, **169**, 21-39.

Thomson, J., Nixon, S., Croudace, I W., Pedersen, T F., Brown, L., Cook, G T. and MacKenzie, A B. 2001. Redox-sensitive element uptake in north-east Atlantic Ocean sediments (Benthic Boundary Layer Experiment sites). *Earth and Planetary Science Letters.*, **184 Issue 2**, 535-547.

Ting, S. 1937. The coastal configuration of western Scotland. *Geografiska Annaler*, **19**, 62-83.

Trueblood, S. & Morton, N. 1991. Comparative sequence stratigraphy and structural styles of the Slyne Trough and Hebrides Basin. *Journal of the Geological Society of London*, **148**, 197-202.

- Tully, M.C. 1970. North Channel, Irish Sea Cruise Report, Project 69/7. *Marine Geophysics Unit Report*, pages 9.
- Tully, M.C. 1971. Cruise report from MV Researcher, 25th June - 30th July 1971: Regional geophysical surveys off NW Scotland, MGU project No. 71/05. *Marine Geophysics report*.
- Ulrich, J. 1964. Zur Topographie der Rosemary-Bank. *Kieler Meeresforschung*, **20**, 95-100.
- Unnithan, V., Shannon, P.M., McGrane, K., Readman, P.W., Jacob, A.W.B., Keary, R. & Kenyon, N.H. 2001. Slope instability and sediment redistribution in the Rockall Trough: constraints from GLORIA. In: Shannon, P.M., Haughton, P.D.W. & Corcoran, D.V. (eds.). *Special Publication- Geological Society Of London. The Petroleum Exploration of Ireland's Offshore Basins*. **188**. Geological Society of London, London, 439-454.
- Upton, B.G.J. 1988. History of Tertiary igneous activity in the N Atlantic borderlands. In: Morton, A.C. & Parson, L.M. (eds.). *Early Tertiary volcanism and the opening of the NE Atlantic*. **39**. Geological Society of London, London, 429-453.
- van Weering, T.C.E. & de Rijk, S. 1991. Sedimentation and climate-induced sediments on Feni Ridge, Northeast Atlantic Ocean. *Marine Geology*, **101**, 49-69.
- Vanneste, K., Henriot, J.P., Posewang, J. & Theilen, F. 1995. Seismic stratigraphy of the Bill Bailey and Lousy Bank area: implications for subsidence history. In: Scrutton, R.A., Stoker, M.S., Shimmield, G.B. & Tudhope, A.W. (eds.). *The Tectonics, Sedimentation and Palaeoceanography of the North Atlantic Region*. **90**. Geological Society of London, London, 125-139.
- Vanney, J.R. & Stanley, D.J. 1983. Shelfbreak physiography; an overview. In: Stanley, D.J. & Moore, G.T. (eds.). *The shelfbreak: critical interface on continental margins*. **33**. Society of Economic Paleontologists and Mineralogists Special Publication, No. 33, 1-24.
- Vogt, P.R. 1986. Seafloor topography, sediments and paleoenvironments. In: Hurdle, B.G. (ed.). *The Nordic Seas*. Springer-Verlag, New York, 237-412.
- Vogt, U., Makris, J., O'Reilly, B.M., Hauser, F., Readman, P.W., Jacob, A.W.B. & Shannon, P.M. 1998. The Hatton basin and continental margin: Crustal structure from wide-angle seismic and gravity data. *Journal of Geophysical Research - Solid Earth*, **103**, 12545-12566.
- von Weymarn, J.A. 1979. A new concept of glaciation in Lewis and Harris, Outer Hebrides. In: *The natural environment of the Outer Hebrides*. **Section B: Biological Sciences**. **77**; Royal Society of Edinburgh, Edinburgh, United Kingdom, 97-105.
- Waagstein, R., Morton, A.C., Praegel, N.O. & Taylor, P.N. 1989. Highly alkaline lapilli tuffs from Rosemary Bank: evidence for continental lithosphere beneath the northern part of the Rockall Trough. *Terra Abstracts*, **1**, 31-32.

- Waddams, P. & Cordingley, T. 1997. The regional geology and exploration potential of the NE Rockall Basin. pages 379-390 in *Petroleum geology of Northwest Europe; 5th conference*. Fleet, A.J. & Boldy, S.A.R. (editors). (London: Geological Society of London.)
- Wade, I.P. 1995. Subarctic intermediate water in the Eastern North Atlantic. Unpublished thesis, University of East Anglia.
- Weaver, P.P.E., Kenyon, N.H. & Wynn, R. 2001. Sedimentation patterns on the North-East Atlantic margin. *GSA/GSL Joint Conference 'Earth System Processes - a Global Meeting', Edinburgh, 24-28 June 2001*.
- Weaver, P.P.E., Wynn, R.B., Kenyon, N.H. & Evans, J.M. 2001. Sedimentation patterns along the Northeast Atlantic continental margin. *In: Glacier-Influenced Sedimentation on High-Latitude Continental Margins*. University of Bristol, 63.
- Weaver, P.P.E., Wynn, R.B., Kenyon, N.H. & Evans, J. 2001. Continental margin sedimentation, with special reference to the north-east Atlantic margin. *Sedimentology*, **47**, 239-256.
- Westbrook, G.K. 1985. Cruise Report RRS "Charles Darwin" 4/85, 17 May - 11 June 1985. *Cruise Report*, pages 7.
- Weymarn, J.A.V. 1974-75. Coastline development in Lewis and Harris, Outer Hebrides, with particular reference to the effects of glaciation. Unpublished thesis, University of Aberdeen.
- Wheeler, A.J., van Weering, T.C.E., Hayes, C.P., Hall, I.R. & Devoy, R.J.N. 1999. Turbiditic and contouritic controls on sediment flux in the central northern Rockall Trough. 84 in *North East Atlantic Slope Processes: Multi-Disciplinary Approaches, Abstract Book. Conference 24-27th January 1999*. Friend, P. & Kenyon, N. (editors). Southampton Oceanography Centre, Southampton, UK.)
- Wheeler, A.J., Bett, B.J., Billet, D.S.M. & Masson, D.G. 2001. High resolution side-scan mapping of deep-water coral mounds: Surface morphology and processes affecting growth. *Journal of Conference Abstracts*, **6**, 749.
- Wheeler, A.J., Billett, D.S.M., Masson, D.G., Olu-Le, R.K. & Grehan, A. 2001. The impact of benthic trawling on NE Atlantic coral ecosystems. *ICES 2001 Meeting, Oslo*. ICES.)
- White, R.S., Westbrook, G.K., Fowler, S.R., Spence, G.D., Barton, P.J., Joppen, M., Margan, M.J., Bowen, A.N., Prescott, C. & Bott, M.H.P. 1987. Hatton Bank (northwest U.K.) continental margin structure. *Geophysical Journal of the Astronomical Society*, **89**, 265-272.
- White, R.S. 1988. A hot-spot model for early Tertiary volcanism in the N Atlantic. *In: Morton, A.C. & Parson, L.M. (eds.). Early Tertiary volcanism and the opening of the NE Atlantic*. Geological Society of London Special Publication, No. 39, 3-13.
- Whittaker, T.J.T. 1991. Shoreline wave power on the Isle of Islay. *Underwater Technology*, **17**, 9-15.

Whittington, R.J. 1983. Seismic reflection studies of Quaternary sediments in the Hebridean, Malin, Irish and Celtic seas. *In: Anonymous (ed.). The Seventh U.K. geophysical assembly. 73.* Blackwell [for the] Royal Astronomical Society, London, United Kingdom, 288.

Whittington, R.J. & Dobson, M.R. 1986. Seismic investigations of sedimentary structure in the northern Rockall Trough; a synopsis. *Proceedings of the Royal Society of Edinburgh, Section B, 88*, 53-54.

Whittington, R.J. & Omran, M.A. 1996. Late Cenozoic sediments on the continental margin in the area around the Hebrides Terrace Seamount, eastern Rockall Trough. 67 in *Applied Geoscience conference Warwick University 15-18 April 1996.* Geological Society, London.)

Wilson, J.B. & Roberts, D.G. 1971. MV "Surveyor" Cruise 71/1, 26 February - 11 April 1971. Tide gauges : geology and geophysics on the Hebridian Shelf and on the Rockall Plateau. *Cruise Report*, pages 10.

Wilson, J.B. 1975. RRS Challenger Cruise 14/74 Legs 1 & 2, August - September 1974: Geological investigations on the continental shelf to the west and north of Scotland. *Cruise Report*.

Wilson, J.B. 1979. Biogenic carbonate sediments on the Scottish continental shelf and on Rockall Bank. *Marine Geology, 33*, M85-93.

Wilson, J.B. 1979. The distribution of the coral *Lophelia Pertusa*(L.)(*L. prolifera*(Pallas)) in the Northeast Atlantic. *Journal of the Marine Biological Association of the UK, 59*, 149-164.

Wilson, J.B. 1979. The first recorded specimens of the Deep-water coral *Lophelia Pertusa* (Linnaeus 1758) from British Waters. *Bulletin of the British Museum, Natural History, Zoology Series, 36*, 209-215.

Wilson, J.B. 1979. Biogenic carbonate sediments on the Scottish continental shelf and on Rockall Ban. *Marine Geology, 33*, M85-M93.

Wilson, J.B. 1979. 'Patch' development of the deep-water coral *Lophelia Pertusa* (L.) on Rockall Bank. *Journal of the Marine Biological Association of the United Kingdom, 59*, 165-177.

Wilson, J.B. 1981. Biogenic carbonate sediments on the Scottish continental shelf and on Rockall Bank (ABSTRACT). *Journal of the Geological Society of London, 138*.

Wilson, J.B. 1982. Shelly faunas associated with temperate offshore tidal deposits. *In: Stride, A.H. (ed.). Offshore tidal sands; processes and deposits.* Chapman and Hall. London, United Kingdom, Pages 126-17.

Wilson, J.B. 1982. RRS Discovery Cruise 123, 5th August - 10th September 1981: Geological investigations on the continental slope to the west of the British Isles and Norway. *Cruise Report*, pages 10.

- Wilson, J.B. 1982. RRS Discovery Cruise 147, 26th April - 17th May 1984: Biochemical fluxes in the NE Atlantic (42N and 50N transects). *Cruise Report*.
- Wilson, J.B. 1982. RRS "Challenger" Cruise 9/81, 16 June - 1 July 1981. Sediment and faunal investigations on the continental shelf to the west and north of Scotland. *Cruise Report*, pages 22.
- Wilson, J.B. 1983. RRS Challenger Cruise 12/79: 18 August - 7 September 1979. Geological investigations in the northern North Sea and on the continental shelf west and north of Scotland. Recovering and relaying of current meter moorings south and west of the Wyville-Thomson. pages 22.
- Wilson, J.B. 1984. *RRS Frederick Russell cruise 7/ 83; 21 April - 13 May 1983; Sediment and faunal investigations on the continental shelf and upper continental slope west and north of Scotland and in the Irish Sea*. (Cruise Report - Institute of Oceanographic Sciences). Institute of Oceanographic Sciences, Surrey, United Kingdom.
- Wilson, J.B. 1986. Faunas of tidal currents and wave dominated continental shelves and their use in the recognition of storm deposits. *In: Knight, R.J. & McLean, M.R. (eds.). Shelf Sands and Sandstones*. 11. Canadian Society of Petroleum Geologists, 313-326.
- Wilson, J.B. 1988. A model for temporal changes in the faunal composition of shell gravels during a transgression on the continental shelf around the British Isles. *In: Nelson Campbell, S. (ed.). Sedimentary Geology. Non-tropical shelf carbonates; modern and ancient*. 60. Elsevier. Amsterdam, Netherlands., 95-105.
- Wingfield, R. 1980. Cruise Report MV Whitehorn 80.08.1: North Channel and western Irish Sea. *Marine Geology Internal Report*.
- Wood, M.V., Hall, J. & Van-Hoorn, B. 1987. Post-Mesozoic differential subsidence in the north-east Rockall Trough related to volcanicity and sedimentation. *In: Brooks, J. & Glennie, K. (eds.). Petroleum Geology of North West Europe*. Graham and Trotman, London, 677-685.
- Wood, M.V. 1988. Integrated geophysical studies West of the Hebrides. Unpublished thesis, University of Glasgow.
- Wood, M.V., Hall, J. & Doody, J.J. 1988. Distribution of early Tertiary lavas in the NE Rockall Trough. *In: Morton, A.C. & Parson, L.M. (eds.). Early Tertiary volcanism and the opening of the NE Atlantic*. (Geological Society of London Special Publications). 39. Geological Society of London, London, 283-292.
- Wynn, R.B., Weaver, P.P.E., Kenyon, N.H. & Masson, D.G. 2001. Geological hazards on the Northeast Atlantic continental margin. 739 in *EUG XI 'Continental Slope Stability (COSTA) of Ocean Margins', 8-12 April 2001, Strasbourg, France*.
- Yuan, F. 1997. Acoustic and geotechnical properties of the Western Irish Sea sediments (BL). Unpublished thesis, University of Bangor.

Zhang, X., Gao, Z. & Yao, X. 1999. Internal wave deposits in the northeastern Rockall Trough, North Atlantic Ocean; reinterpretation of deep-water sediment wave formation. *Acta Sedimentologica Sinica*, **17**, 464-472.

Ziegler, P.A. 1981. Evolution of Sedimentary Basins in North-West Europe. *In: Illing, L.V. & Hobson, G. (eds.). Petroleum geology of the continental shelf of North-West Europe*. Heyden and Son, London, 3-39.

Ziegler, P.A. 1982. *Geological atlas of Western and Central Europe*. Elsevier for Shell International Petroleum, Maatschappij, BV, Amsterdam.


Ziegler, P.A. 1987. Evolution of the Arctic North Atlantic borderlands. *Petroleum Geology of North West Europe*, 1201-1204.

Ziegler, P.A. 1988. Evolution of the Arctic-North Atlantic and the Western Tethys. *Memoir of the American Association of Petroleum Geologists*, No. 43.

Ziegler, P.A. 1988. Evolution of the Arctic-North Atlantic and the western Tethys.

Appendix 5

OCCURRENCE OF CARBONATE AND OTHER MOUNDS IN THE SEA7 REGION OF THE ATLANTIC MARGIN



VIKRAM UNNITHAN¹, PAT SHANNON²



**MARINE AND PETROLEUM GEOLOGY RESEARCH GROUP
DEPARTMENT OF GEOLOGY
UNIVERSITY COLLEGE DUBLIN
IRELAND**

SEA7 REPORT

JULY 2003

¹ EMAIL: Vikram.Unnithan@ucd.ie

² EMAIL: P.shannon@ucd.ie

TABLE OF CONTENTS

1.2 SUMMARY.....	III
INTRODUCTION.....	1
OBJECTIVES.....	1
DELIVERABLES.....	2
HISTORICAL BACKGROUND TO CARBONATE MOUND RESEARCH.....	2
CARBONATE MOUNDS.....	3
MOUND TYPES.....	3
<i>Hovland Mound Province.....</i>	<i>4</i>
<i>Belgica Mound Province.....</i>	<i>4</i>
<i>Pelagia Mound Province.....</i>	<i>4</i>
<i>Logachev Mound Province.....</i>	<i>4</i>
<i>GSI Mound Province.....</i>	<i>5</i>
<i>Porcupine Bank Canyon Mound Province.....</i>	<i>5</i>
<i>Magellan Mounds.....</i>	<i>5</i>
<i>Darwin Mounds.....</i>	<i>6</i>
<i>Other Mound Structures.....</i>	<i>6</i>
MOUND AGES.....	7
FAUNAL ASSEMBLAGES.....	7
CURRENT CARBONATE MOUND MODELS.....	8
HYDROCARBON AND FLUID SEEP MODELS.....	8
SEDIMENTARY AND OCEANOGRAPHIC CONDITIONING MODEL.....	9
DISCUSSION.....	10
CONCLUSIONS.....	11
BIBLIOGRAPHY.....	13

SUMMARY

In the past decade, a significant amount of new data has been gathered on the geological, ecological and biological aspects of seabed mounds structures in the Atlantic Ocean west of Ireland and the UK. Carbonate mounds are very numerous (>1000), of various sizes and occur as both seabed features and buried mounds. They lie in water depths ranging from 500 to 1500 m and are generally located towards the upper parts of the shelf-slope break along the basin margins, and also on the main basin-bounding banks. Their morphology and shape range from simple cones to complex amalgamated ridge features covering up to 5 km² and standing up to 300 m in height. While the internal structures and composition of the carbonate mounds are poorly constrained, their surfaces are generally covered by reef-building cold-water coral species such as *Lophelia pertusa* and *Madrepora oculata*. In addition to carbonate mounds, volcanic cones and possible mud mounds have also been identified in the region.

This report focuses on the distribution, classification and internal (seismic) structure of carbonate mounds to the west of Ireland and the UK within the SEA7 region and its general environs. It also provides an overview of the main results from ongoing research in EU 5th Framework projects such as GEOMOUND, ECOMOUND and ACES. Various models for the origin and growth of carbonate mounds have been proposed in the literature. These range from hydrocarbon seepage and nutrient models to oceanographic and current influences and are discussed. The report also provides a comprehensive bibliography of mound references, a map of mound locations and a compendium of relevant metadata from the region.

Introduction

Extensive clusters of carbonate build-up and mound structures have been identified at the seabed or in the shallow subsurface in the North Atlantic. Their formation is generally linked to the development and growth of deep, cold-water coral species such as *Lophelia pertusa* and *Madrepora oculata*.

Recent discoveries of reefs and mound structures in the North Atlantic, during various research cruises such as the Training Through Research Cruises (ITR 97), Atlantic Irish Rockall Survey (AIRS96), Belgica Cruises 1997 - 1998, and Pelagia ENAM cruises 1996 - 1998, have generated a great deal of interest in the genesis of deep water bioherms. The possible linkage of modern seabed mounds and their fossil counterparts with hydrocarbon seepage and petroleum plays has intensified the study of these features.

Carbonate mud mounds in shallow waters are defined by Bosence and Bridges (1995) as “*a carbonate build-up having a depositional relief and being composed dominantly of carbonate mud, peloidal and or micrite*”. Classical mounds, as described by various authors (e.g. Monty 1995, Pratt 1995) are restricted to water depths of 50 - 100 m. The most striking characteristic of the mounded structures in the North Atlantic is the water depths at which they occur i.e. 500 - 1000m. These seabed mounds are also unique as regards to their physical size i.e. 10 – 300 m in height and covering an area of a few 100 m² to 3 km² in some cases. In addition to carbonate mound, volcanic cones and possible mud volcanoes (mud mounds) have also been identified using high-resolution swath bathymetry data acquired by the Geological Survey of Ireland as part of its recent Seabed Survey, and high resolution magnetic and gravity data acquired by the Irish Petroleum Affairs Division.

This report summarises data from research on deep, cold-water carbonate mounds, drawing on the wealth of information available from the various scientific cruises, research meetings and recent publications. Current models attempting to shed light on the genesis and evolution of these features are highlighted and discussed.

Objectives

The scope of the present report includes:

- A review of current knowledge on the occurrence of carbonate and other mounds features that support unusual biological communities in the general region of interest (SEA7: SW corner 22.3°W, 52.5°N & NE corner 4°W, 61.7°N) (Fig. 1).
- Collation and inventory of known (carbonate) mound localities in the region of interest.
- To provide a bibliography providing relevant sources of information on carbonate mound work.
- To provide an inventory of meta data for the known areas and general environs.

Deliverables

In addition to a paper copy of this report, a CD-ROM accompanies this report, which includes:

1. This report in Word Format
2. Bibliography in Endnote format
3. Inventory of meta-data
4. ArcView 3.2 Project with GEBCO-CE bathymetry with mound locations superimposed.

Historical Background to Carbonate mound research

The scientific expeditions of Wyville-Thomson, Carpenter and Jeffreys in the early 1870s onboard the HMS Porcupine west of Ireland led to the discovery of numerous hitherto unidentified deep-sea fauna. During these cruises dredge samples of deep-sea coral were obtained. The occurrence of the deep-sea, cold water coral species *Lophelia pertusa* was first reported by Thomson (1874). Scientific work continued during the early 1900's in the Porcupine Seabight during which the Helga and Helga II worked on an extensive series of stations, particularly on the eastern and northern flanks of the Seabight, in water depths of less than 1600 m (Le Danois, 1948).

During the First and Second World wars relatively little work was done in the region. The only exception was Le Danois' synoptic account of the continental slope fauna off the coasts of northwestern Europe with remarkable details on the general facies in the Porcupine Seabight. His facies interpretation of different European continental slope types and showed the sites of coral occurrences. According to Le Danois, these occurred mostly in a narrow bathymetric zone between 500-1000 m along the continental slope and formed huge belts up to several hundreds of meters high and thousands of meters long. In the Porcupine Seabight he reported such a band of coral on the eastern flank and named it "massif de la Baie de Dingle".

After Le Danois's work on corals to the west of Ireland and U.K., the focus of deep-sea coral research shifted to the Norwegian margin. Ever since the first detailed description of *Lophelia pertusa* detailed studies on the biology and anatomy of deep-sea corals especially *Lophelia pertusa*, *Madrepora*, *Desmophyllum* and *Dendroophyllia* were subject of many research projects (Freiwald, 1998; Hovland and Mortensen, 1999; Mortensen, 2000). Also in other areas of the European margin, *Lophelia* was studied but less intensively than in Norway. Coral structures have been found along the northwest continental margin in the fjords and offshore Norway (Freiwald, 1998; Hovland and Mortensen, 1999; Mortensen, 2000), around the British Isles (Wilson, 1975; Wilson, 1979a; Wilson, 1979b), the Faroe Islands (Frederiksen et al., 1992; Jensen and Frederiksen, 1992), the Rockall Bank (Scoffin et al., 1980; Wilson, 1979c), the Rockall Trough (de Haas et al., 2000; Kenyon et al., 1998), the Gollum channels (Tudhope and Scoffin, 1995), the Porcupine Bank (Scoffin and Bowes, 1988) and in the Mediterranean Sea (Zibrowius, 1980). In addition to the northeastern Atlantic, corals and coral structures have also been reported along the margin of West Africa (Zibrowius and Gili, 1990), in the northwest Atlantic Blake Plateau (Reed, 1992; Stetson et al., 1962), near the Bahamas (Messing et al., 1990; Mullins et al., 1981; Neumann et al., 1977; Neumann and Paull, 1998), on Hatteras Slope (Paull et al., 2000), in the Gulf of Mexico (Moore and Bullis, 1960) and from a few scattered records from the Pacific (Squires, 1965) and Indian Ocean (Rogers, 1999).

In the early 1990s the deep-water coral structures to the west of Ireland and U.K. were the focus of academic and commercial hydrocarbon interest. An extensive seismic and sampling surveys for oil exploration in the Porcupine Seabight showed mounded dome-like structures on the seabed. Gravity cores from the mounds yielded *Lophelia pertusa* and mostly muddy sediments. The

publication by Hovland et al. (1994) suggested a relationship between mounds and interpreted deeper faults underneath the mounds and proposed a model linking hydrocarbon seepage and mound occurrence.

These observations and the proposed model resulted in renewed academic interest in the subsurface of the Porcupine Seabight. The first cruise to the mounds in the Porcupine Seabight was organized by the RCMG (Renard Centre of Marine Geology) in the framework of the EU MAST III projects - CORSAIRES and ENAM II to study slope instabilities along the eastern flank of the Porcupine Seabight with high-resolution seismic profiles. This first cruise by the RV Belgica found in addition to the surface mounds reported by Hovland et al. (1994) discovered numerous buried mound features further to the north of the Hovland province. This first successful seismic survey was followed by a Training Through Research- CORSAIRES cruise (Kenyon et al., 1998) during which shallow sediment cores and sidescan sonar data was collected from different mound provinces. During the summer of 1998 in the course of a large-scale regional survey of the seabed by the Atlantic Frontier Environmental Network (AFEN) mounds now called the Darwin mounds were discovered. The findings of these cruises led to the funding of three EU funded projects - GEOMOUND, ECOMOUND and ACES - with the aim of studying the internal and external controls on coral mound build-ups and the biology of the deep-water corals along the NE European margins. The Geological Survey of Ireland has recently completed an extensive swath bathymetry survey across the Irish designated waters west of Ireland and has detected many new mounds.

Carbonate Mounds

Carbonate mounds in the North Atlantic have been identified from their surface and subsurface (seismic) morphology. The carbonate mound catalogue presented in this report has been compiled by the authors from a variety of data sources: the updated seismic catalogue by Croker & O'Laughlin (1998), De Mol (2002), Unnithan (2001), GSI Seabed Survey (pers. comm.) and published sources. The datasets used to identify the mound features include sidescan sonar (TOBI, GLORIA, OKEAN, ORETECH), seismic (both commercial and academic), multibeam (GSI Seabed Survey) and ROV video (Victor6000).

1.3 MOUND TYPES

The carbonate mounds can be broadly categorised into surface and buried structures. A carpet of coral debris, which supports a living coral fauna consisting mainly of *Lophelia pertusa* and *Madrepora Oculata*, covers the surface mounds (Fig. 5). In addition to these colonial species, the solitary coral *Desmophyllum cristagalli* and the octocoral *Stylaster* are also occasionally present. On the basis of ROV, multibeam, sidescan sonar, echosounder and seismic records, a number of mound provinces have been identified. They occur in the northern and eastern Porcupine Seabight, Porcupine, Rockall, Hatton and Fangorn banks, to the west of Ireland (see Figs 1 & 2). The published literature makes reference to a large number of mound provinces with a proliferation of names. The names are based on a variety of characteristics including geographic location, morphology, internal geometry and even the names of either the vessels or scientists involved in their original discovery. Mounds can be either surface or buried features. They are principally located along the banks and shelf-slope breaks (Fig. 2). In addition, many samples or corals have been recovered or reported from dredge samples along the Atlantic margin west of Ireland and the UK (Fig. 3). The main mound provinces include the following:

1.3.1 Hovland Mound Province

This is named after Martin Hovland, lead author of a pioneering paper on mounds in the northern Porcupine Seabight (Hovland et al., 1994) (Fig. 4). These mounds are generally conical to dome shaped structures 1.5 km in diameter and about 150 m in height (Fig. 6). They are situated around the 700 to 800 m isobaths. These mounds are each typically surrounded by a distinct moat (up to 50 m deep). Seismic characteristics include pull-up of parallel reflectors just below the mounds. Characteristic mounds include the Propellor and Perseverance mounds.

1.3.2 Belgica Mound Province

This is named after the ship that discovered these features along the eastern margin of the Porcupine Seabight during a scientific cruise in the summer of 1997. These structures are pinnacle or conical mounds with similar dimensions as the Hovland mounds (Fig. 7). They are found in groups forming echelon NNE-SSW trending barriers or banks. The presence of a seismically transparent layer beneath the mounds distinguishes them from the Hovland mounds. Within this province, proto-mound structures (Moirra mounds) have been identified on the basis of sonar and ROV observations. These structures are 5-10m high, have a diameter of 20-30m and seem to be morphologically similar to the Darwin mounds identified in the northern U.K. Rockall Trough (Masson *et al.* 2001). Mounds and mound family sites where detailed work has concentrated during the GEOMOUND and ECOMOUND projects include the Thérèse Mound and the Challenger Mound.

1.3.3 Pelagia Mound Province

This consists of a cluster of mounds on the northern Porcupine Bank investigated by the RV Pelagia, a Dutch research vessel. These mounds are larger (average 300 m in height, covering a few square km) than the Hovland and Belgica mounds. They are also found in slightly deeper waters (750 to 1200 m). These mounds have complex shapes ranging from conical to elongate (both alongslope and downslope) banks to amalgamated structures (Fig. 8).

1.3.4 Logachev Mound Province

This consists of complex mound structures on the eastern margin of the Rockall Bank, investigated first by the Russian research vessel Professor Logachev in 1997 (Kenyon *et al.* 1998). These mounds, later investigated by the Victor ROV (R2 Site) and TOBI sidescan sonar, show highly complex amalgamated ridge and mound structures (Fig. 9). Seismic profiles across this highlight the downslope trending complex ridge structures. The Rockall Bank mounds, occupying a 100km zone parallel to the bathymetric contours on the Rockall Bank, are smaller and morphologically complex in shape. With respect to the mounds described in previous sections, they are found in shallower waters ranging from 500 - 1000m. The mounds, which are covered by dead and living assemblages of *L. Pertusa* and *M. Oculata*, are perched on what is presumed outcrops of rock, erratics or glacial debris. Sonar and seismic data acquired on the second leg of the TTR 7 cruise, clearly demonstrated this preference for a pre-existing platform to develop on. Seismically, they look similar to the other types of mounds, with a chaotic internal structure and a dominant, high amplitude basal reflector. Along the margins of some of the mounds, there is evidence for velocity pull-up. Strong currents and especially the movement of sand influence the intricate shapes of these mounds. Bedforms due to currents on the SE Rockall Bank show evidence for strong alongslope currents with significant accumulation of coarse sandy material between the mounds being transported and deposited by strong currents. In some cases, there is evidence of sand waves covering the mounds or sand transport across the tops of the mounds. Preliminary results suggests that both the Porcupine Bank and Rockall Bank mounds are covered by a carpet of coral debris which supports a living coral fauna consisting mainly of *L. pertusa* and *M. oculata*. In addition to these colonial species, the solitary coral

Desmophyllum cristagalli and the octocoral *Stylaster* are also occasionally present. Morphologically these mounds appear more complex than the mounds in the Porcupine.

1.3.5 GSI Mound Province

The occurrence of new surface mounds on the Rockall, Hatton and Fangorn banks has been reported by the Geological Survey of Ireland (Geoghegan & Monteys, 2003). These mounds were imaged using multibeam and echosounder data from the Seabed Survey (Fig. 10). Detailed analysis of the extent, size and morphology of these mound structures is ongoing (Geoghegan & Monteys, 2003). Preliminary examination of the Fangorn Bank mounds suggests that they are morphologically similar to the Hovland mounds. The Fangorn Bank mounds have distinct moat structures around the mounds, are generally conical in shape, and are about 350m in height. However, these mounds are found in slightly deeper water (around 1250 m) than the Hovland mounds.

1.3.6 Porcupine Bank Canyon Mound Province

This loosely refers to the mounds found in association with Porcupine Bank Canyon on the southern margin of the Porcupine Ridge. These mounds have been identified on the basis of GLORIA sidescan sonar, echosounder and recently acquired multibeam data (Xavier Monteys pers. comm.). These mounds are conical-shaped structures about 300 m in height and are more frequent on the southern steeper flank of the Porcupine Bank Canyon.

1.3.7 Magellan Mounds

These are numerous (>1000) buried mound-like structures in the northern Porcupine Seabight. The buried mounds or Magellan mounds (Fig. 11), were first observed in 1993 on industrial high resolution seismics and later investigated by Belgica 1997 Cruise in the northern Porcupine Seabight. All the identified Magellan mounds are situated to the north of the Hovland mounds. These mounds have not been found elsewhere in the northern Atlantic and are unique to the Porcupine Seabight. They have no morphological expression at the seafloor and were first identified on the basis of high-resolution seismic data collected by the survey vessel Svitzer Magellan. They are characterised by lung-shaped, ovoid seismically transparent structures, 50 - 100 m in height and up to 250 m in diameter. These mounds are generally found to the north and west of the Hovland Mound Province, in water depths of 500 - 750m. Some of the mounds occur in pairs. Orthogonal seismic lines across some of the mound pairs, shows that these mounds are singular ring shaped mounds around a central kernel. The frequency and size of the Magellan mounds decreases to the north, essentially following the same trend as the Hovland mounds. On seismic sections, the base of the mounds appears to originate from a local unconformable seismic horizon. Surprisingly, there is little or no loss of signal directly below the mounds and all seismic horizons can be traced as continuous horizons. Along the margins of the mounds there is little velocity pull-up and little or no distortion of the seismic horizons. Within the mounds, there is a complete masking or blanketing. No internal structures are observed with the mounds. The seismic horizons directly above the mounds are slightly convex or domed.

1.3.8 Darwin Mounds

These were discovered in the 1998 by scientists from the SOC during the AFEN environmental TOBI survey north and west of Scotland. These mounds are named after the NERC ship RV Charles Darwin. The second group of Darwin mounds were discovered during a DTI survey in 1999. The two fields of mounds cover an area of 13km x 4km and 13km x 9km respectively and are found in water depths ranging from 900 to 1060m. The two surveys discovered in excess of 225 mound structures. Each mound is c. 50-100m in diameter and up to 5m high (Fig. 12). The mounds are found north of a large area of pockmarks and appear to be closely related to these fluid escape

features (Masson et al., 2003). The Darwin mounds have a characteristic teardrop shaped 'tails' of moderately high backscatter on TOBI sidescan sonar images spanning a few hundred meters in length downstream (NE-SW) from the mounds. The tops of the mounds are covered by *Lophelia pertusa* and less abundant *Madrepora oculata* and associated fauna. Average density of coral heads is reported to be 1 colony per 4m² (Bett, 2001). The mound tails are characterised by high-density populations of xenophyophores, which are giant protozoans unique to the deep-sea. The pockmark fauna is similar to the background sediments with higher density of large burrows and echinothurid urchins dominate the megafauna.

1.3.9 Other mound structures

On the AIRS GLORIA sonar images, numerous oval features were observed along the lower slope of the northern Porcupine Bank at 54°N and 14°W (Unnithan, 2001; Kenyon et al., 1998). These are 500m to 1km in diameter and are characterised by very high backscatter. The conical shape is determined from the shape of the shadows, i.e. a conical object has high backscatter on the side facing the transducers and prolonged distinct shadows. While the mounded features bear some resemblances to carbonate mounds, these sharp conical features are interpreted to represent extrusive volcanic cones (Fig. 13). Unfortunately there are no 3.5 kHz echosounder data over these features, but high local magnetic and gravity anomalies support the presence of a volcanic body. Kenyon et al. (1998) describe these conical features using medium range sonar and shallow seismic data. The conical shape and low acoustic penetration of these features (see Fig. 23 of Kenyon et al. 1998) is consistent with volcanic bodies. Grab samples from the features recovered black rocks and manganese encrustations. Kenyon et al. (1998) suggested that the black rocks were igneous/volcanic and their presence may be related to the source of the manganese. Based on seismic and magnetic data Naylor et al. (1999) named these preserved Paleocene volcanic cones as the Droll Igneous Centre. The possibility that some of the igneous cones contain a covering veneer of carbonate coral material cannot be discounted.

There are three oval, 2-10 km in diameter high backscattering areas along the eastern margin of Lorien Bank, south-western Rockall Trough. The boundaries of these regions are well-defined. The surrounding region is characterised by low, homogeneous backscattering. The strong contrast of the oval bodies with the surrounding area suggests an abrupt change in surface texture and hence, lithology. The results from recent swath bathymetry surveys (confidential data PAD) suggest the presence of numerous volcanic seamounts of similar morphology to the Anton Dohrn, but smaller in size, in the vicinity of Lorien Bank (Peter Croker, pers. comm.). Their location, size, shape and backscattering characteristics suggest that these structures are volcanic seamount-type features (Unnithan, 2001). The backscattering characteristics are also similar to those of carbonate mounds. However, individual carbonate mounds are generally an order of magnitude smaller than these features. The recent GSI Seabed survey has brought to light new mound structures of presumed volcanic nature based on their seamount morphology, high magnetic signature. These mounds are found in the southern Rockall Trough, Fangorn Bank and Eriador Seamount (see Fig. 2). In addition, some geophysical evidence has been obtained to suggest the presence of some active mud volcanoes or mud mounds in the Porcupine Seabight (Jean-Pierre Henriët, pers. comm.).

Mound Ages

There is significant debate in academic circles regarding the ages of carbonate mound structures along the margins of the Rockall Trough and Porcupine Seabight. To date, none of the large mounds in the Irish and UK waters has been drilled to the base. Surface box and giant piston cores obtained from the summit and flanks of the mounds provide ages in the order of 200,000 years B.P.

(Andre Freiwald, pers. comm.). Regional seismo-stratigraphic analysis suggests that the base of the mounds in the Porcupine Seabight and Rockall Trough overlies the C10 reflector. This reflector has been dated by the ODP 980/981 drill site in the Rockall Trough as early Pliocene (STRATAGEM Project Report – Atlas, Pat Shannon, pers. comm.). These ages are in general agreement with ages calculated by O'Reilly et al. (2003) based on *Lophelia pertusa* growth rates reported in literature and mound dimensions (height and width) of mounds in the Pelagia province. Seismo-stratigraphic correlation by the RCMG group in Gent (e.g. de Mol, 2001) have proposed a slightly younger Late Pleistocene age based on sample dates from the flanks of the mounds in the eastern Porcupine Seabight.

Faunal Assemblages

There is large faunal variability (nature, distribution and type) between the mounds observed on video footage obtained during the CARACOLE 2001 and Polarstern Victor XIX3a 2003 cruise to the Porcupine Seabight and Rockall Trough mounds. The main species of corals found on all the mounds is *Lophelia pertusa*. Varying amounts of *Madrepora oculata*, *Stylasterid* and *Desmophyllum* are found on the mounds. Mounds such as Perseverance, Propellor and Theresa (see attached meta-dataset Figure CARACOLE for locations) do not have substantial living coral cover and are characterised by a variety of otrehspecies of actinian, antipatharian, gorgonians (soft corals), glass sponges, encrusting (as yet undetermined) yellow and blue sponges, cidaris (sea urchins), arthropods such as *paromola*, *bathynectes* and fish species such as *Lopheus*, *Neocyttus helgae*, *Pisces blanoides*, *chimera*. Mounds on the eastern margin of the Rockall Bank are covered by a thick (1-3m) cover of both living and dead *Lophelia pertusa* with *Madrepora oculata*. *Desmophyllum* is not seen on these mounds. Small stylasters are observed growing on these mounds. Characteristics fauna includes stalkless crinoids, large red gorgonians, antipatharians, cup-shaped sponges and a large variety of crab and fishes.

Current Carbonate Mound Models

Models for carbonate mound growth must address aspects of mound initiation, growth and sustenance. In terms of mound initiation, two main end-member models exist in the published literature: 1) hydrocarbon or fluid seepage migration (e.g. Hovland et al., 1994) and 2) sedimentological and oceanographic conditioning (e.g. de Mol, 2002). These models are described and discussed below.

1.4 HYDROCARBON AND FLUID SEEPAGE MODELS

Three models have been proposed an evolution and triggering of the Hovland and the Magellan mounds by hydrocarbon and fluid seepage processes:

- The first model was proposed by Hovland (1990), Hovland et al. (1994, 1998). This suggested a causal relationship between hydrocarbon seeps and deep-water coral features. Deep-water coral banks could form as a consequence of local fertilization that resulted from focused hydrocarbon seepage. In the case of the Hovland mounds in the Porcupine Basin, Hovland et al. (1994) proposed that the coral mound structures are located immediately above major

faults, which could function as hydrocarbon migration pathways (Fig. 14). Hydrocarbon seeps provided a food source for bacteria, fertilizing the waters around the seep sites. The model suggested that focused hydrocarbon seeps along the faults or fractures led to local eutrophication and enhanced chemosynthetic bacterial growth, which encouraged and supported higher organisms by providing them with nutrients and a suitable substrate on which to attach. This symbiotic relationship would have laid the basis for further growth of the mounds. The validity of this model has been questioned recently by Bailey et al. (2003), whose analysis of the high-resolution 3-D seismic data from the same area suggests that faults and mounds are not spatially related and that many of the features interpreted as faults by Hovland et al. (1994) are seismic artifacts.

- The potential role of methane in triggering mound growth in the Magellan province has been evoked by Henriot et al. (1998; 2001) in a model developed to explain possible causal relationships between methane migration, slope destabilization and mound growth (Fig. 15). Implicit in this model is a possible phase of hydrate build-up and decay resulting from variations in bottom water temperatures during the last glacial and interglacial periods. This model provides a process for focused seeps around which the coral bank develop, hence forming ring-shaped coral banks in the Magellan province. During Quaternary times, with repeated fluctuations from polar (in front of an Irish ice-sheet) to temperate conditions, extreme variations in bottom water temperatures (up to 11°C) may have translated into cycles of local growth and decay of gas hydrates, fuelled by methane from deeper hydrocarbon reservoirs. The authors suggest that the Magellan mound province coincides with an underlying slope failure, where hydrates may have played a role.
- A third model was proposed by McDonnell (2001) based on the coincidence between coral banks and the relation of underlying Oligocene and Miocene contourite deposits. The underlying contourites may have played a role as fluid migration pathway from underlying Tertiary deltaic strata. However, no direct evidence of migration or of hydrocarbon accumulation are reported in the work of Mc Donnell (2001). Seepage of thermogenic hydrocarbons at the seafloor does occur at the present in the northern part of the Porcupine Basin, in the non-commercial Connemara oil field about 65km north of the Magellan mound province. These seeps appear clearly on industrial seismic data and high resolution seismic sections as high-amplitude patches along reflections, vertical disruption of seismic reflections or vertical zones of chaotic facies, interpreted as gas chimneys (Games, 2001). However, no mounds are reported in this region, while in the Magellan, Hovland or Belgica mound provinces there are no such fluid migration features. Most large faults do not extent to the seafloor and there are no clear indications so far for vertical fluid migration pathways neither on industrial seismic data (McDonnell, 2001) nor on the interpreted high-resolution seismic data. Pockmarks in the region of the coral banks are not active structures but might have been active in the recent past. To date no paleo-pockmarks horizons have been observed in the data set, which can link the mound development to the coral mound evolution.

1.5 SEDIMENTARY AND OCEANOGRAPHIC CONDITIONING MODEL

There is ample evidence from sidescan sonar and echosounder data to suggest that the mounds of the Porcupine and Rockall banks are strongly influenced by currents (O'Reilly et al., 2003). The mounds are generally located in areas (Wilson, 1979) with rugged seabed topography or close to the shelf edge. The corals may have settled in these areas to take advantage of the increased currents and possible suspended matter. Along the Porcupine Bank, there is evidence for a relationship between the occurrence of corals and water stratification. Ahermaphyitic corals accumulate below the thermocline which exists at 600-1400m in the Rockall Trough (Rice, 1991).

Changes in currents, water temperature or nutrient supply force the coral colonies to selectively occupy favourable parts of the larger mounds. The dead coral traps pelagic and hemipelagic detritus and when the environmental conditions are favourable for colonisation eventually forms a substrate

for the further growth of corals. This model could explain the growth of carbonate mounds as a biological response to changes in environmental conditions coupled with normal pelagic/hemipelagic sedimentation.

The sedimentary and oceanographic model consists of initiation and development phases (e.g. Unnithan et al., 2003):

- Mound initiation: *Lophelia pertusa* needs a hard substrate for initial settlement. For the initiation of mounds, this could be an erosion surface such as the C10, glacial drop-stones or pre-existing topography. In addition to settlement substrate, the presence of a suitable current regime to deliver food and prevent clogging through sedimentation as well as optimum water temperature and salinity regimes govern the distribution of *Lophelia pertusa*. Once successfully established and mound growth initiated, the growth phase of the mound would be influenced by changing oceanographic conditions which dictate the selective survival of certain coral patches.
- Mound development: This starts with the growth and death of coral thickets, cementation and hard-ground formation followed by erosion which provides hard substrate for the next growth cycle at that location (see Fig. 16). Once the mounds are able to generate their own (micro) topography, growth can be sustained simply because the corals are in an elevated position with enhanced capacity to catch nutrients and are removed from moving sand. Unnithan et al.'s (2003) observations show that lateral growth of mounds occurs when upper and mid-slope terraces are formed by current action and colonised by corals. Dense coral stands or fronds (often with thin fan-like growth composed of basal dead coral with an outer edge of living coral) show preferential growth on the up current vertical face of the terrace. The dead coral matrix may serve to baffle sediments behind the leading living coral edge promoting extension of the terrace over time following cementation.

1.6 DISCUSSION

In the light of new evidence from various cruises in the northern Porcupine, the Hovland model needs to be modified. Recent sediment and core samples from the mounds show no clear evidence of chemosynthetic faunal assemblages. Furthermore, geochemical and isotope analysis reveals the dominance of a marine ^{13}C fingerprint. Moreover, high-resolution seismic profiles show little evidence for faulting directly beneath the mounds. This implies that a direct hydrocarbon fuelled cycle for the growth of the Hovland mounds is unlikely. However sidescan sonar data from the TTR 7 and recently the Pelagia TOBI 2002 cruises imaged pockmarks in the northern and eastern Porcupine, which suggests that fluid expulsion has occurred in the basin.

A vital component in the seepage models is the causal relation between the distribution of the coral banks and structures identified as potential fluid seepage sites. Henriët et al. (2001) related the buried Magellan mounds to the occurrence of the polygonal faults observed in a Neogene slide succession; Hovland et al. (1994) related the coral banks to deeper faults and McDonnell (2001) to Oligocene contourite deposits in the Magellan and Hovland mounds. All three models assume that for the start-up phase the coral banks are preferentially located at sites where fluid seepage is highest.

Fluids need not necessarily have been hydrocarbons. Basinal brine venting, along unconformities, fractures or by diffusion, could have facilitated the formation of authigenic carbonate, forming hard grounds on which ambient benthic fauna could have settled on (Henriët et al., 1998, 2001; Hovland et al., 1994). The formation of local hard grounds by venting may explain the spatial distribution of the coral banks.

Hovland et al. (1994) assume that the shape of the coral banks is the result of hydrocarbon seepage distribution. Henriët et al. (1998, 2001) interpreted a ring shape of a large number of Magellan

mounds, which may argue for a focused venting. If deep-water coral banks were caused by a close, direct link between hydrocarbon seepage along faults, we might expect the coral banks to be strictly lined up along the faults, which is apparently not the case as discussed above. In the detailed 3D and 2D studies of the Magellan mound province (Britsurvey, 1997; Huvenne et al., submitted; McDonnell, 2001) no ring structures has been observed as interpreted by Henriët et al. (1998; 2001) The elongated shape indicates the current direction as illustrated on TOBI sidescan sonar data by O'Reilly et al (2003). These observations argue for an external control of the coral bank development.

In the Belgica mound province some local enhanced reflectors are observed in the top of some of the Neogene-Holocene contourite units. This might indicate a very local and small accumulation of some fluids as gas. However, the gas content in the samples taken on the Thérèse mound, overlying these contourites, is low and in the order of atmospheric values (Jochen Naeth, pers comm.). During the Marion Dufresne coring campaign MD123 of 2001 (Van Rooij et al., 2002) a H₂S smell was reported for one core located on Thérèse Mound. However, follow-up chemical analysis of the core failed to confirm or detect the presence of any H₂S (Jochen Naeth, pers. comm.).

No conclusive evidence is seen in the present day setting between methane seeps and coral mounds formation. Only in the case of the Darwin mounds there seems to be a link between sand extrusion, mound formation and colonisation by deep-sea, cold water corals. However it more than likely that the Darwin mounds are selectively colonised by corals due to their elevated topography and other external factors such as currents and nutrient supply. Nevertheless none of the proposed models provide conclusive evidence regarding the origin of the coral structures because they are based on sparse data. Only further deep drilling will provide a conclusive answer.

The carbonate mounds tend to occur in elongate clusters and are best developed in Irish waters on the banks and along the upper parts of the shelf/slope break. Surprisingly few mound provinces have been located to date in UK waters. However, comparison with the geological/bathymetric setting of the known locations, together with the presence of dredge samples of corals (see Fig. 3) would suggest that mound provinces could be anticipate in the general environs of the Wyville-Thomson Ridge.

What controls coral and mound growth? To date, little is know about the biology and life cycle of *Lophelia pertusa*. For example, when and how often do they spawn? What mechanisms and strategies do they adopt for colonisation? Which of the factors among water temperature, pressure/depth, salinity, turbidity, nutrients or currents are the most important for healthy coral growth? For the limited video footage current direction, erosion (presence/absence), nutrients, turbidity and substrate (hardground/dropstone) seem to be major primary controls. For mound growth, the coral growth pattern (patchy, debris & live) and erosion by deep-sea currents also seem to be the most important parameter.

Conclusions

In the past few years, our knowledge of cold water bioherms has increased significantly. Some of the important conclusions that we draw from this report are:

- Mounds occurring in clusters have been reported from on the eastern, northern Porcupine Seabight, Porcupine Bank, Rockall Bank, northern Rockall Trough, Lorien Bank, Fagorn Bank, Eriador Seamount and Hatton Bank.

- Seabed mounds have been documented on the basis of a variety of seismic, sidescan sonar, video and multibeam bathymetry datasets. Most of the mounds are inferred to be carbonate structures based on seismic data, while a few mounds structures, such as on Lorien Bank, are suggested to be volcanic in origin.
- A greater number and variety of mounds are present, in relatively deep waters of 500 - 1200m, on the continental shelf to the west of Ireland. Surprising, only few large mounds structures have been found to the west of U.K. Further mounds are anticipated in the general vicinity of the Wyville-Thomson Ridge.
- A Plio-Pleistocene age has been suggested based on seismo-stratigraphic correlations for some of the larger mounds in the along the margins of the Porcupine Seabight and Rockall Trough
- Faunal assemblages associated with the large carbonate mounds are primarily composed of cold water corals *Lophelia pertusa*, *Madrepora Oculata*, *Desmophyllum* and stlyasterids.
- During recent video cruises to the carbonate mound sites in the Porcupine Seabight, definite evidence for damage to the mounds and corals due to fishing activity was seen. Lost static fishing gear such as gill nets and trawl marks were observed around the mound sites.
- The nature and origin of the carbonate mounds cannot be described by a simple fault related, hydrocarbon seep and bacterial symbiotic model. Various oceanographic, biological and palaeoclimatic variables and their influence on cold water bioherms and mounds are likely to play a role in the initiation and growth of the carbonate mounds.
- The carbonate mounds and the corals are synonymous with the presence of strong currents. Water stratification and internal tides also appears to be important factors determining the location of the carbonate mounds.

Although we have a better insight of the nature, composition, structure and areal distribution of these seabed mounds, their genesis and formation remains elusive. Primary oceanographic variables such as currents, temperature, salinity and suspended matter most definitely influence the carbonate mounds and cold-water corals. But how have these mounds evolved through time and what initiated their growth in the first place, are all some of the important unanswered questions. Current ongoing research on the samples taken from the mounds should be able to answer some of these questions. Perhaps a more integrated oceanographic, biological and geological approach in addition to drilling is required to fully comprehend the nature and complexity of these seabed features.

BIBLIOGRAPHY

- Abraham, D.A. and Ritchie, J.D., 1991. The Darwin Complex: a Tertiary igneous centre in the northern Rockall Trough. *Scottish Journal of Geology*, 27: 113-125.
- Aken, H.M.V., 2000a. The Hydrography of the Mid-Latitude Northeast Atlantic Ocean {II}: The Intermediate Water Masses. *Deep Sea Research Part I*, 47: 789-824.
- Aken, H.M.V., 2000b. The Hydrography of the Mid-Latitude Northeast Atlantic Ocean I: The Deep Water Masses. *Deep Sea Research Part I*, 47: 757-788.
- Aloisi, G., Pierre, C., Rouchy, J.-M., Foucher, J.-P. and Woodside, J., 2000. Methane-related authigenic carbonates of eastern Mediterranean Sea mud volcanoes and their possible relation to gas hydrate destabilisation. *Earth and Planetary Science Letters*, 184(1): 321-338.
- Andrulleit, H., Freiwald, A. and Schafer, P., 1996. Bioclastic carbonate sediments on the southwestern Svalbard shelf. *Marine Geology*, 134(3-4): 163-182.
- Bailey, W., Shannon, P.M., Walsh, J.J. and Unnithan, V., 2003. The spatial distributions of faults and deep sea carbonate mounds in the Porcupine Basin, offshore Ireland. *Marine and Petroleum Geology* (in press).
- Barton, A.J. and White, R.S., 1997. Volcanism on the Rockall continental margin. *Tectonic, magmatic and depositional processes at passive continental margins*, 154(3): 531-536.
- Belderson, R.H., Kenyon, N.H. and Wilson, J.B., 1973. Iceberg plough marks in the northeast Atlantic. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 13(3): 215-224.
- Belderson, R.H., Kenyon, N.K. and Wilson, J.B., 1985. The exploration of the British continental shelf with side-scan sonar, TV and photography, IOS Annual Report, pp. 70-75.
- Berggren, W.A. and Aubert, J., 1976. Late Paleogene (late Eocene and Oligocene) benthonic foraminiferal biostratigraphy and paleobathymetry of Rockall Bank and Hatton-Rockall Basin. *Micropaleontology*, 22(3): 307-326.
- Bett, B.J., 2001. UK Atlantic Margin Environmental Survey: Introduction and overview of bathyal benthic ecology. *Continental Shelf Research*, 21(8-10): 917-956.
- Beyer, A., Schenke, H.W., Klenke, M. and Niederjasper, F., 2003. High resolution bathymetry of the eastern slope of the Porcupine Seabight. *Marine Geology*, 198(1-2): 27-54.
- Bianchi, G.G. and McCave, I.N., 2000. Hydrography and sedimentation under the deep western boundary current on Bjorn and Gardar Drifts, Iceland Basin. *Marine Geology*, 165: 137-169.
- Blamart, D. et al., 2001. Modern NE Atlantic Ocean Cold Water Coral Characteristics. *Geophysical Research Abstracts*, 3: 9320.
- Booth, D.A., 1988. Eddies in the Rockall Trough. *Oceanologica Acta*, II(3): 213--319.
- Bosence, D.W.J. and Bridges, P.H., 1995. A review of the origin and evolution of Carbonate Mounds. *Special Publication of the International Association of Sedimentologists*, 23: 3--10.
- Boulvain, F., 2001. Facies architecture and diagenesis of Belgian Late Frasnian carbonate mounds. *Sedimentary Geology*, 145(3-4): 269-294.

- Britsurvey, T.O.M.p., 1997. Site survey Irish Block 35/17-1,14/11/96-13/12/96. *Report released by Petroleum Affairs Division, Dublin.*
- Broecker, W.S., 1994. Massive iceberg discharge as triggers for global climate change. *Nature*, 372: 421-424.
- Bull, J.M. and Masson, D.G., 1996. The southern margin of the Rockall Plateau and stratigraphy, Tertiary volcanism and plate tectonic evolution. *Journal of the Geological Society of London*, 153(4): 601-612.
- Bunch, A.W.H., 1979. A detailed seismic structure of Rockall Bank (55 degrees N, 15 degrees W) and a synthetic seismogram analysis. *Earth and Planetary Science Letters*, 45(2): 453-463.
- Campbell, J.M., Keeton, J. and Le Bas, T.P., 1997. Investigation of the performance of the GLORIA swath bathymetry system. *SOC Unpublished Interium Report*, : 25.
- Chapman, M.R. and Shackleton, N.J., 1999. Global ice-volume fluctuations, North Atlantic ice-rafting events, and deep-ocean circulation changes between 130 and 70ka. *Geology*, 27(9): 795-798.
- Cook, D.R., 1987. The Goban Spur and exploration in a deep water frontier basin. In: J. Brooks and K.W. Glennie (Editors), *Petroleum geology of north west Europe*, pp. 623-632.
- Cope, J.C.W., Ingham, J.K. and Rawson, P.F., 1992. *Atlas of Palaeogeography and Lithofacies*. The Geological Society, London.
- Courtney, R. and Shaw, J., 2000. Multibeam bathymetry and backscatter Imaging of the Canadian Continental Shelf. *Geoscience Canada*, 27: 31-42.
- Cremer, M., Faugeres, J.-C., Grousset, F. and Gonthier, E., 1993. Late Quaternary sediment flux on sedimentary drifts in the Northeast Atlantic. *Contourites and bottom currents*, 82(1-4): 89-101.
- Crocker, P.F. and Klemperer, S.L., 1989. Structure and stratigraphy of the Porcupine Basin and relationships to deep crustal structure and the opening of the North Atlantic. *Extensional tectonics and stratigraphy of the North Atlantic margins*, 46: 445-459.
- Crocker, P.F. and Shannon, P.M., 1987. The evolution and hydrocarbon prospectivity of the Porcupine Basin, Offshore Ireland. In: J. Brooks and K.W. Glennie (Editors), *Petroleum Geology of North West Europe*. Graham and Trotham, pp. 633-642.
- Crocker, P.F. and O'Loughlin, O., 1998. A Catalogue of Irish Offshore Carbonate Mud Mounds. *TTR-7 Post Cruise Conference on Carbonate Mud Mounds and Cold Water Reefs*, 1: 13.
- Crocker, P.F. and Shannon, P.M., 1995. The petroleum geology of Ireland's offshore basins; Introduction. In: P.F. Crocker and P.M. Shannon (Editors), *The Petroleum Geology of Ireland's Offshore Basins*, Geological Society London, Special Publications, 93: 1-8.
- Cronin, B.T. and Kidd, R.B., 1998. Heterogeneity and Lithotype Distribution in Ancient Deep-Sea Canyons: Point Lobos Deep-Sea Canyon as a Reservoir Analogue. *Sedimentary Geology*, 115: 315-349.
- Cuif, J.P., Dauphin, Y., Freiwald, A., Gautret, P. and Zibrowius, H., 1999. Biochemical markers of zooxanthellae symbiosis in soluble matrices of skeleton of 24 Scleractinia species. *Comparative Biochemistry and Physiology a-Molecular and Integrative Physiology*, 123(3): 269-278.
- Cunningham, A.P., Barker, P.F. and Tomlinson, J.S., 1998. Tectonic and Sedimentary Environment of the North Scotia Ridge Region Revealed by Side-Scan Sonar. *Journal of the Geological Society, London*, 155: 941-956.

- Damuth, J.E., 1978. Echo characteristics of the Norwegian - Greenland Sea: Relationship to Quaternary Sedimentation. *Marine Geology*, 28: 1-36.
- Damuth, J.E., 1980. Use of High-Frequency (3.5-12kHz) Echograms in the Study of Near-Bottom Sedimentation Processes in the Deep-Sea: A Review. *Marine Geology*, 38: 51-75.
- Davies, M. et al., 1998. Seabed Database Collation Report, North Atlantic. Data Gathering Projects for the STC and ETC, Rockall Studies Group. *Internal Classified Report*, 1: 500+.
- de Graciansky, P.C. et al., 1985a. Site 550. In: P.C. de Graciansky et al. (Editors), Initial reports of the Deep Sea Drilling Project covering Leg 80. U.S. Govt. Printing Office, pp. 251-355.
- de Graciansky, P.C. et al., 1985b. Site 551. In: P.C. de Graciansky et al. (Editors), Initial reports of the Deep Sea Drilling Project covering Leg 80. U.S. Govt. Printing Office, pp. 357-385.
- De Mol, B., 1998. Geosphere-biosphere coupling: Carbonate Mud Mounds and Cold Water Reefs: Workshop Report No. 143. *UNESCO publication*, 143: 60.
- De Mol, B. and RUG. Faculteit Wetenschappen. Vakgroep geologie en bodemkunde, 2002. Development of coral banks in Porcupine seabight (SW Ireland) a multidisciplinary approach. s.n., Gent, XI, 363 + bijlage pp.
- De Mol, B. et al., 2002. Large deep-water coral banks in the Porcupine Basin, southwest of Ireland. *Marine Geology*, 188(1-2): 193-231.
- Deacon, M.B., 1996. How the Science of Oceanography Developed. In: C.P. Summerhays and S.A. Thorpe (Editors), *Oceanography: An Illustrated Guide*. Manson Publishing, pp. 9-26.
- Dickson, R.R. and Kidd, R.B., 1986. Deep circulation in the southern Rockall Trough, the oceanographic setting of Site 610. In: W.F. Ruddiman et al. (Editors), *Initial reports of the Deep Sea Drilling Project covering Leg 94*. U.S. Govt. Printing Office, pp. 1061-1074.
- Dickson, R.R. and McCave, I.N., 1986. Nephloid layers on the continental slope west of Porcupine Bank. *Deep Sea Research*, 33(6): 791 - 818.
- Dingle, R.V., Megson, J.B. and Scrutton, R.A., 1982. Acoustic stratigraphy of the sedimentary succession west of Porcupine Bank, NE Atlantic Ocean and a preliminary account. *Marine Geology*, 47: 17-35.
- Dingle, R.V. and Scrutton, R.A., 1979. Sedimentary succession and tectonic history of a marginal plateau (Goban Spur, southwest of Ireland). *Marine Geology*, 33: 45-69.
1972. Sequence Stratigraphy and Depositional History of the Eastern Mississippi Fan (Pleistocene), Northeastern Deep Gulf of Mexico Sonographs of the Sea Floor: A Picture Atlas. *AAPG Bulletin*, 82: 1207-1232, 185.
- Dons, C., 1935. Die Verbreitung von Steinkorallen in West-Finnmark. *Zoologische Notizen XXIX Det Kongelige Norske Videnskabers Selskab*, VIII(18): 57-60.
- Dons, C., 1944. Über die nördlichsten Korallenriffen der Welt. *Zoologische Notizen XXV. Det Kongelige Norske Videnskabers Selskab*, VI: 206-209.
- Dorschell, B., Hebbeln, D., Rüggeberg, A. and Dullo, C., in press. Carbonate budget of a cold-water coral carbonate mound: Propeller Mound, Porcupine Seabight. In: J.P. Henriot and C. Dullo (Editors), *Geomound Ecomound Book*. Springer Verlag.

- Frederiksen, R., Jensen, A. and Westerberg, H., 1992. The distribution of the scleractinian coral *Lophelia Pertus* around the Faroe Islands and the relation to internal tidal mixing. *Sarsia*, 77: 157-171.
- Freiwald, A., 1995. Bacteria-Induced Carbonate Degradation - a Taphonomic Case- Study of *Cibicides Lobatulus* from a High-Boreal Carbonate Setting. *Palaios*, 10(4): 337-346.
- Freiwald, A., 1998a. Geobiology of *Lophelia Pertusa* (Scleractinina) Reefs in the North Atlantic. Unpublished Habilitationsschrift zur Erlangung der venia legendi am Fachbereich Geowissenschaften der Universitat Bremen Thesis, Bremen.
- Freiwald, A., 1998b. Modern nearshore cold-temperate calcareous sediments in the Troms District, Northern Norway. *Journal of Sedimentary Research*, 68(5): 763-776.
- Freiwald, A. and Henrich, R., 1994. Reefal Coralline Algal Build-Ups within the Arctic-Circle - Morphology and Sedimentary Dynamics under Extreme Environmental Seasonality. *Sedimentology*, 41(5): 963-984.
- Freiwald, A., Huhnerbach, V., Lindberg, B., Wilson, J.B. and Campbell, J., 2002. The Sula Reef Complex, Norwegian shelf. *Facies*, 47: 179-200.
- Freiwald, A. and Mortensen, P.B., 2000. The first record of the deep-water coral *Stenocyathus vermiformis* (Pourtales, 1868) (Scleractinia, Guyniidae) from Norwegian waters. *Sarsia*, 85(3): 275-276.
- Freiwald, A. and Mostafawi, N., 1998. Ostracods in a cold-temperate coastal environment, western Troms, northern Norway: Sedimentary aspects and assemblages. *Facies*, 38: 255-273.
- Freiwald, A., Reitner, J. and Kruttschinna, J., 1997. Microbial alteration of the deepwater coral *Lophelia pertusa*: Early postmortem processes. *Facies*, 36: 223-226.
- Freiwald, A. and Schonfeld, J., 1996. Substrate pitting and boring pattern of *Hydrokkin sarcophaga* Cedhagen, 1994 (Foraminifera) in a modern deep-water coral reef mound. *Marine Micropaleontology*, 28(2): 199-207.
- Freiwald, A., Wilson, J.B. and Henrich, R., 1999. Grounding Pleistocene Icebergs Shape Recent Deep-Water Coral Reefs. *Sedimentary Geology*, 125: 1-8.
- Gage, J.D., 2001. Deep-sea benthic community and environmental impact assessment at the Atlantic Frontier. *Continental Shelf Research*, 21(8-10): 957-986.
- Games, K., 2001. Evidence of shallow gas above the Connemara oil accumulation, Block 26/28, Porcupine Basin, In: P.M. Shannon, P.D.W. Haughton and D.V. Corcoran (Editors), *Petroleum Exploration of Ireland's Offshore Basins*. Geological Society, London, Special Publications, 188, 361-373.
- Gautret, P., Cuif, J.P. and Freiwald, A., 1997. Composition of soluble mineralizing matrices in zooxanthellate and non-zooxanthellate scleractinian corals: Biochemical assessment of photosynthetic metabolism through the study of a skeletal feature. *Facies*, 36: 189-194.
- GEBCO, 1997. General Bathymetric Charts of the Ocean, British Oceanographic Data Centre.
- Geoghegan, M. and Monteys, X., 2003. The Irish National Seabed Survey - Carbonate mounds and deep water coral - new discoveries and known occurrences revisited. *Geophysical Research Abstracts*, 5: 14803.
- Gordon, J.D.M., 2001. Deep-water fisheries at the Atlantic Frontier. *Continental Shelf Research*, 21(8-10): 987-1003.

- Grehan, A.J., Unnithan, V., Olu - Le Roy, K. and Opderbecke, J., 2003. Fishing impacts on Irish deep-water coral reefs: making the case for coral conservation. *American Fisheries Society Publication*, submitted.
- Haq, B.U., 1998. Gas Hydrates: Greenhouse Nightmare ? Energy Panacea or Pipe Dream. *GSA Today*, 8(11): 1-6.
- Harwood, J. and Wilson, B., 2001. The implications of developments on the Atlantic Frontier for marine mammals. *Continental Shelf Research*, 21(8-10): 1073-1093.
- Heggland, R., 1998. Gas seepage as an indicator of deeper prospective reservoirs. A study based on exploration 3D seismic data. *Marine and Petroleum Geology*, 15(1): 1-9.
- Henrich, R. et al., 1995. Controls on Modern Carbonate Sedimentation on Warm-Temperature to Arctic Coasts, Shelves and Seamounts in the Northern- Hemisphere - Implications for Fossil Counterparts. *Facies*, 32: 71-108.
- Henriet, J.P. et al., 1998. Gas Hydrate Crystals May Help Build Reefs. *Nature*, 391: 648-649.
- Henriet, J.P., de Mol, B., Vanneste, M., Huvenne, V., van Rooij, D. and the Porcupine-Belgica '97, 98 and 99 Shipboard Parties., 2001. Carbonate mounds and slope failures in the Porcupine Basin: a development model involving fluid venting. In: P.M. Shannon, P.D.W. Haughton and D.V. Corcoran (Editors), *Petroleum Exploration of Ireland's Offshore Basins*. Geological Society, London, Special Publications, 188, 375-383.
- Hovland, M., 1983. Elongated depressions associated with pockmarks in the Western slope of the Norwegian Trench. *Marine Geology*, 51: 35-46.
- Hovland, M., Croker, P.F. and Martin, M., 1994. Fault-associated seabed mounds (carbonate knolls?) off western Ireland and north-west Australia. *Marine and Petroleum Geology*, 11(2): 232-246.
- Hovland, M., Mortensen, P.B., Brattegard, T., Strass, P. and Rokoengen, K., 1998. Ahermatypic Coral Banks Off Mid-Norway: Evidence for a Link with Seepage of Light Hydrocarbons. *Palaios*, 13: 189-200.
- Hovland, M. and Risk, M., 2003. Do Norwegian deep-water coral reefs rely on seeping fluids? *Marine Geology*, 198(1-2): 83-96.
- Huvenne, V.A.I. et al., in press. The Magellan mound province in the Porcupine Basin. In: J.P. Henriët and C. Dullo (Editors), *Geomound Ecomound Book*. Springer Verlag.
- Huvenne, V.A.I., Blondel, P. and Henriët, J.P., 2002a. Textural analyses of sidescan sonar imagery from two mound provinces in the Porcupine Seabight. *Marine Geology*, 189(3-4): 323-341.
- Huvenne, V.A.I., Croker, P.F. and Henriët, J.P., 2002b. A refreshing 3D view of an ancient sediment collapse and slope failure. *Terra Nova*, 14(1): 33-40.
- Huvenne, V.A.I., De Mol, B. and Henriët, J.-P., 2003. A 3D seismic study of the morphology and spatial distribution of buried coral banks in the Porcupine Basin, SW of Ireland. *Marine Geology*, 198(1-2): 5-25.
- Jensen, A. and Frederiksen, R., 1992. The Fauna associated with the Bank-forming Deepwater Coral *Lophelia Pertusa* (Scleractinaria) on the Faroe Shelf. *Sarsia*, 53-63.
- Jordan, C. and Wilson, J.E., 1998. Reefs: Geologic Considerations for Geophysicists. *Leading Edge*, 17(3): 325--328.
- Judd, A.G. and Hovland, M., 1992. The evidence of shallow gas in marine sediments: *Continental Shelf Research*, 12: 1081-1095.

- Kenyon, N.H. et al., 2003. Giant carbonate mud mounds in the southern Rockall Trough. *Marine Geology*, 195(1-4): 5-30.
- Kenyon, N.H., Ivanov, M.K. and Akhmetzhanov, A.M., 1998. Cold water carbonate mounds and sediment transport on the Northeast Atlantic margin. *UNESCO Intergovernmental Oceanographic Commission Technical Series*, 52: 178.
- Kopaska-Merkel, D.C. and Haywick, D.W., 2001. Carbonate mounds: sedimentation, organismal response, and diagenesis. *Sedimentary Geology*, 145(3-4): 157-159.
- Lance, S. et al., 1998. Submersible study of mud volcanoes seaward of the Barbados accretionary wedge: sedimentology, structure and rheology. *Marine Geology*, 145(3-4): 255-292.
- Le Danois, E., 1948. Les profondeurs de la mer. *Paris, Payot*: 303.
- Lindberg, B., Berndt, C. and Mienert, J., in press. The Fugløy Reefs on the Norwegian-Barents Continental Margin: Cold-water Corals at 70°N, their acoustic signature, geologic, geomorphologic and oceanographic setting. In: J.P. Henriot and C. Dullo (Editors), *Geomound Ecomound Book*. Springer Verlag.
- Long, D., 2001. The Western Frontiers Association - Evaluating seabed conditions west of the UK. *Continental Shelf Research*, 21(8-10): 811-824.
- MacDonald, I.R., Sager, W.W. and Peccini, M.B., 2003. Gas hydrate and chemosynthetic biota in mounded bathymetry at mid-slope hydrocarbon seeps: Northern Gulf of Mexico. *Marine Geology*, 198(1-2): 133-158.
- Masson, D.G., 2001. Sedimentary processes shaping the eastern slope of the Faeroe-Shetland Channel. *Continental Shelf Research*, 21(8-10): 825-857.
- Masson, D.G. et al., 2003. The origin of deep-water, coral-topped mounds in the northern Rockall Trough, Northeast Atlantic. *Marine Geology*, 194: 159-180.
- McDonnell, A., 2001. Comparative Tertiary Basin Development in the Porcupine and Rockall Basins, Unpublished Ph.D. Thesis, University College Dublin pp. 175.
- McDonnell, A. and Shannon, P.M., 2001. Comparative Tertiary stratigraphic evolution of the Porcupine and Rockall basins. In: P.M. Shannon, P.D.W. Haughton and D.V. Corcoran (Editors), *Petroleum Exploration of Ireland's Offshore Basins*. Geological Society, London, Special Publications, 188, 323-344.
- Monty, C.L.V., 1995. The rise and nature of carbonate mud-mounds. *Special Publication of the International Association of Sedimentologists*, 23: 11-48.
- Mortensen, P.B., 2000. *Lophelia pertusa* (Scleractinia) in Norwegian waters. Distribution, growth and associated fauna. PhD Thesis, University of Bergen, Bergen.
- Mortensen, P.B., Hovland, M., Brattegard, T. and Farestveit, R., 1995. Deep Water Bioherms of the Scleractinian Coral *Lophelia pertusa* (L.) at 64 N on the Norwegian Shelf: Structure and Associated Megafauna. *Sarsia*: 145-158.
- Mortensen, P.B., Hovland, M., Fosså, J.H. and Furevik, D., 2001. Distribution, abundance and size of *Lophelia pertusa* coral reefs in mid-Norway in relation to seabed characteristics. *Journal of the Marine Biological Association of the United Kingdom*, 84: 581-597.
- Mortensen, P.B. and Rapp, H.T., 1998. Oxygen- and carbon isotope ratios related to growth line patterns in skeletons of *Lophelia pertusa* (L.) (Anthozoa: Scleractinia): Implications for determination of Lineas extension rates. *Sarsia*, 83: 433-446.

- Morton, A.C. et al., 1988. Early Tertiary volcanic rocks in Well 163/6-1A, Rockall Trough. In: A.C. Morton and L.M. Parson (Editors), *Early Tertiary Volcanism and the Opening of the NE Atlantic*. Geological Society Special Publications, 39: 293-308.
- Naeth, J. et al., in press. From the Connemara Oil Field to the Belgica Mound Province: Modelling hydrocarbon migration in the Porcupine Basin, offshore Ireland. In: J.P. Henriët and C. Dullo (Editors), *Geomound Ecomound Book*. Springer Verlag.
- Naylor, D., Shannon, P. and Murphy, N. 1999. *Irish Rockall Basin region - a standard structural nomenclature system*. Petroleum Affairs Division Special Publication 1/99, 42pp.
- New, A.L. and Smythe-Wright, D., 2001. Aspects of the circulation in the Rockall Trough. *Continental Shelf Research*, 21(8-10): 777-810.
- O'Reilly, B.M., Readman, P.W., Shannon, P.M. and Jacob, A.W.B., 2003. A model for the development of a carbonate mound population in the Rockall Trough based on deep-towed sidescan sonar data. *Marine Geology*, 198(1-2): 55-66.
- Paull, C.K., Neumann, A.C., am Ende, B.A., Ussler III, W. and Rodriguez, N.M., 2000a. Lithoherms on the Florida-Hatteras slope. *Marine Geology*, 166(1-4): 83-101.
- Paull, C.K., Neumann, A.C., Ende, B.A.A., Ussler, W. and Rodriguez, N.M., 2000b. Lithoherms on the Florida--Hatteras Slope. *Marine Geology*, 166: 83-101.
- Peter, J.M., Simoneit, B.R.T., Kawka, O.E. and Scott, S.D., 1990. Liquid hydrocarbon-bearing inclusions in modern hydrothermal chimneys and mounds from the southern trough of Guaymas Basin, Gulf of California. *Applied Geochemistry*, 5(1-2): 51-56.
- Pillen, S., 1998. Detailkartering En Seismische Analyse Van de Magellan-Mounds in Het Porcupine Bekken, Ten Zuidwesten Van Ierland. Vakgroep Geologie en Bodemkunde, Faculteit Wetenschappen, Universiteit Gent, Belgium.
- Pinheiro, L.M. et al., 2003. Mud volcanism in the Gulf of Cadiz: results from the TTR-10 cruise. *Marine Geology*, 195(1-4): 131-151.
- Racki, G., Racka, M., Matyja, H. and Devleeschouwer, X., 2002. The Frasnian/Famennian boundary interval in the South Polish-Moravian shelf basins: integrated event-stratigraphical approach. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 181(1-3): 251-297.
- Rice, A.L., Billitt, D.S.M., Thurston, M.H. and Lampitt, R.S., 1991. The Institute of Oceanographic Sciences biology programme in the Porcupine Seabight: Background and general introduction. *Journal of Marine Biology Letters*, 71: 281-310.
- Rice, A.L., Thurston, M.J. and New, A.L., 1990. Dense aggregations of a hexactinellid sponge, *Pheronema carpenteri*. In: *the Porcupine Seabight (northeast Atlantic Ocean) and possible causes*, *Prog. Oceanog*, 24: 179-196.
- Riding, R., 2002. Structure and composition of organic reefs and carbonate mud mounds: concepts and categories. *Earth-Science Reviews*, 58(1-2): 163-231.
- Riestedberg, D., West, O., Lee, S., McCallum, S. and Phelps, T.J., 2003. Sediment surface effects on methane hydrate formation and dissociation. *Marine Geology*, 198(1-2): 181-190.
- Roberts, D.G., 1975a. Marine geology of the Rockall Plateau and Trough. *Philosophical Transactions of the Royal Society of London, Series A: Mathematical and Physical Sciences*, 278(1285): 447-509.

- Roberts, D.G., 1975b. Tectonic and stratigraphic evolution of the Rockall Plateau and trough. In: A.W. Woodland (Editor), *Petroleum and the Continental Shelf of North West Europe*, Applied Science Publishers, pp. 77-91.
- Roberts, D.G. and Jones, M.T., 1978. A Bathymetric, Magnetic and Gravity Survey of the Rockall Bank, HMS Hecla 1969. *Admiralty Marine Science Publications*, 19: 43.
- Roberts, J.M., 2000. Coral colonies make a home on North Sea oil rigs. *Reef Encounter*, 27: 17-18.
- Roberts, J.M., Long, D., Wilson, J.B., Mortensen, P.B. and Gage, J.D., 2003. The cold-water coral *Lophelia pertusa* (Scleractinia) and enigmatic seabed mounds along the north-east Atlantic margin: are they related? *Marine Pollution Bulletin*, 46(1): 7-20.
- Rogers, A.D., 1999. The biology of *Lophelia pertusa* (Linnaeus 1758) and Other Deep-Water Reef-Forming Corals and Impacts from Human Activities. *International Revue der gesamten Hydrobiologie*, 84: 315-406.
- Rüggeberg, A., Dullo, C., Dorschel, B. and Hebbeln, D., in press. Environmental changes and growth history of Propeller Mound, Porcupine Seabight: Evidence from benthic foraminiferal assemblages. In: J.P. Henriët and C. Dullo (Editors), *Geomound Ecomound Book*. Springer.
- Sager, W.W., MacDonald, I.R. and Hou, R., 2003. Geophysical signatures of mud mounds at hydrocarbon seeps on the Louisiana continental slope, northern Gulf of Mexico. *Marine Geology*, 198(1-2): 97-132.
- Scoffin, T.P. and Bowes, G.E., 1988. The facies distribution of carbonate sediments on the Porcupine Bank, northeast Atlantic. *Sedimentary Geology*, 60: 125-134.
- Shannon, P.M., O'Reilly, B.M., Readman, P.W. and Jacob, A.W.B., 2001. A deep-towed sidescan sonar (TOBI) survey of the margins of the Rockall Trough: environmental aspects. In: N.J. Murphy and M. Davies (Editors), *Ireland's Deepwater Frontier: Results from the Petroleum Infrastructure Programme (PIP)*. Conference Abstracts, p. 20-23.
- Stoker, M.S., Hitchen, K. and Graham, C.C., 1993. United Kingdom Offshore Regional Report: The geology of the Hebrides and West Shetland shelves, and adjacent deep-water areas, BGS report, pp. 149+.
- Thomson, C.W. 1874. *The Depths of the Sea*. Macmillan, London, pp.527.
- Unnithan, V., 2001. Geological and sedimentological analysis of sidescan sonar data in the Rockall Trough and Porcupine Seabight, west of Ireland. Unpublished PhD thesis, University College Dublin, Dublin, 360 pp.
- Unnithan, V., Grehan, A., van Weering, T. and Olu-Le Roy, K., 2003. Biological and geological characteristics of the R1 and R2 coral mounds, Rockall Trough, west of Ireland. *Geophysical Research Abstracts*, 5: 08946.
- Unnithan, V., Henriët, J.P., Shannon, P. and Croker, P., 2000. Carbonate Mound Provinces To The West Of Ireland: An Overview. *EOS Transactions*, 81(48): F635.
- Unnithan, V. et al., 1997. Side-scan sonar features in the Rockall Trough, offshore Ireland. *Annales Geophysicae*, 15: C384.
- Unnithan, V. et al., 2001. Slope instability and sediment re-distribution in the Rockall Trough: constraints from GLORIA. In: P.M. Shannon, P.D.W. Haughton and D.V. Corcoran (Editors), *Petroleum Exploration of Ireland's Offshore Basins*. Geological Society, London, Special Publications, 188, 439-454.

- Upton, B.G.J., 1988. History of Tertiary igneous activity in the N Atlantic borderlands. In: A.C. Morton and L.M. Parson (Editors), *Early Tertiary Volcanism and the Opening of the NE Atlantic*. Geological Society Special Publications, 39, 429-453.
- Van Rooij, D., 1998. Sequentie-Stratigrafie Met Hoog-Resolutie-Seismiekaan de Rand Van de Keltische Zee En Het Porcupine Bekken, Ten Zuidwesten Van Ierland. Vakgroep Geologie en Bodemkunde, Faculteit Wetenschappen, Universiteit Gent, Belgium.
- Van Rooij, D., De Mol, B., Huvenne, V., Ivanov, M. and Henriët, J.-P., 2003. Seismic evidence of current-controlled sedimentation in the Belgica mound province, upper Porcupine slope, southwest of Ireland. *Marine Geology*, 195(1-4): 31-53.
- van Weering, T.C.E., de Haas, H., de Stigter, H.C., Lykke-Andersen, H. and Kouvaev, I., 2003a. Structure and development of giant carbonate mounds at the SW and SE Rockall Trough margins, NE Atlantic Ocean. *Marine Geology*, 198(1-2): 67-81.
- van Weering, T.C.E., Dullo, C. and Henriët, J.-P., 2003b. An introduction to geosphere-biosphere coupling; cold seep related carbonate and mound formation and ecology. *Marine Geology*, 198(1-2): 1-3.
- van Weering, T.C.E., Stoker, M. and Thomsen, L., 1997. ENAM and ALIPOR shipboard cruise report: R.V. Pelagia, May 22 - June 13, 1997, Netherlands Institute for Sea Research (NIOZ): Unpublished cruise report, pp. 36.
- Vermeulen, N.J., 1997. Hydrography, Surface Geology and Geomorphology of the Deep Water Sedimentary Basins to the West of Ireland, Marine Resources Series, Marine Institute, Dublin, pp. 41+.
- Weir, C.R., Pollock, C., Cronin, C. and Taylor, S., 2001. Cetaceans of the Atlantic Frontier, north and west of Scotland. *Continental Shelf Research*, 21(8-10): 1047-1071.
- Wilson, J.B., 1979a. Biogenic carbonate sediments on the Scottish continental shelf and on Rockall Bank. *Marine Geology*, 33(3-4): M85-M93.
- Wilson, J.B., 1979b. The distribution of the coral *Lophelia Pertusa* (L.) in the north-east Atlantic. *Journal of the Marine Biological Association of the U.K.*, 59: 149-164.
- Wilson, J.B., 1979c. The first recorded specimens of the deep-water coral *Lophelia pertusa* (Linnaeus, 1758) from British waters. *Bull. Br. Mus. Nat. Hist. (Zool.)*, 36: 209-215.
- Wilson, J.B., 1979d. Patch development of the deep-water coral *Lophelia Pertusa* on Rockall Bank. *Journal of the Marine Biological Association of the U.K.*, 59: 165-177.
- Wood, R., 2001. Are reefs and mud mounds really so different? *Sedimentary Geology*, 145(3-4): 161-171.
- WWF, 2001. WWF Factsheet: CORAL REEFS THREATENED OFF BRITAIN.
- WWF, 2002. The Darwin Mounds: Out of sight and still under threat, Web.
- Yin, P., Berne, S., Vagner, P., Loubrieu, B. and Liu, Z., 2003. Mud volcanoes at the shelf margin of the East China Sea. *Marine Geology*, 194(3-4): 135-149.
- Zibrowius, H., 1980. Les Scléactiniaires de la Méditerranée et de l'Atlantique nord-oriental. *Memoires de l'Institut Oceanographique, Monaco*, 11: 1-227.

Figures




Figure 1: Overview map of the study area highlighting the main bathymetric and geomorphological features. Bathymetric contours from GEBCO-CE 2003 dataset.




Figure 2: Location of the carbonate mounds in the study area. Mound provinces described in the text are also highlighted.




Figure 3: Localities where *L. pertusa* has been sampled or photographed are shown. Of the trawls hauls, only those <15 km in length have been included as points centred on the mid-point of the trawl track. FB Faeroe Bank; HB Hatton Bank; LB Lousy Bank, RB Rockall Bank; SB Stanton Bank; WTR Wyville-Thomson Ridge. (from: <http://www.sams.ac.uk/dml/projects/benthic/lophmap.htm>)




Figure 4: Map from the northern Porcupine Seabight showing the distribution of Hovland surface mounds (black filled circles) and buried Magellan mounds (red filled circles).



Figure 5: High-resolution digital still images from the CARACOLE cruise showing (left) the typical framework reef-building coral species *Lophelia pertusa* and *Madrepora oculata* found on carbonate mounds. The image on the right shows the less abundant solitary coral species *Desmophyllum* (red). Images are copyright IFREMER 2001




Figure 6: High-resolution seismic profile showing the characteristic shape of the Hovland Mounds in the Porcupine Seabight. Note the presence of a moat or depression surrounding the mounds. Seismic profile from Renard Centre for Marine Geology (RCMG), Gent, de Mol, 2001.




Figure 7: High-resolution seismic profile showing the characteristic shape of the Belgica Mounds on the eastern margin of the Porcupine Seabight. Note the characteristic ponded sediments on the upslope margin. Seismic profile from RCMG, Gent, de Mol, 2001.




Figure 8: High resolution TOBI sidescan imagery from the northern Porcupine Bank showing the characteristic shapes of Pelagia mounds. (Shannon *et al.* 2001). TOBI swath width approx. 6 km




Figure 9: High-resolution multibeam data from the R2 site, Logachev mound province acquired during the CARACOLE 2001 cruise. This image highlights the complex wave structures on the surface of the mound. The mound summit is shown in pink, with sediment wave amplitudes of up to 1m and wavelengths of 2-3m (Unnithan *et al.* 2003)




Figure 10: 3-D image, viewing North, of the GSI mound province showing the Fangorn and Hatton Bank with the Edoras Bank Seamont to the west (left in image). The Fangorn Bank is approximately 60 km in diameter (NE-SW orientation, measured along dark green contour) and the mounds are up to 300 m in height.




Figure 11: High resolution seismic line over the buried Magellan Mounds. Note these mounds seem to originate (root) from a single horizon. Image spans 300m along the profile and 225 ms 'TWT' along the depth axis.




Figure 12: High-resolution sidescan sonar image of the Darwin mounds. Note these mounds are substantially smaller than those in Irish Waters. From Wheeler (2001).

54.21 N




Figure 13: Other mound structures -GLORIA mosaic highlighting high-backscatter patches on the northern margin of the Porcupine Bank (Unnithan, 2001. Commercial high-resolution magnetic data (P. Croker per comms) suggests that these mound structures are volcanic in origin.




Figure 14: Hovland et al. (1994) suggest a close relation between hydrocarbon seeps and coral bank formation. According to their model, the first condition is that hydrocarbons are generated at depth (A) and that some of them find their way to the surface in a focused manner, through faults and fissures (B). The seabed is here locally eroded by seepage and the local seawater is provided with nourishment on which bacteria and microorganisms depend. Over time, organisms and their skeletal remains accumulate, whereas authigenic carbonates precipitate locally, and cement the sediment and skeletal debris (C).




Fig. 15: Henriët et al. (2001) proposed a model for the genesis of the Magellan mounds on a site of episodic fluid migration, under strong varying bottom water temperatures. Under glacial conditions, a horizon of gas hydrates could built up in regions of prolific (even transient) methane flux. Decay of the hydrated horizons could generate slope failure. In renewed methane flux conditions, in warm waters, the disrupted horizon funneled the migrating fluids to the seabed, possibly contributing to venting and mound nucleation.




Figure 16: Cartoon sketch showing the sedimentary and oceanographic conditioning model explaining the initiation, growth and development of the carbonate mounds based on video analysis (Unnithan *et al.* 2003).