




DATA NOTE

The genome sequence of the Triangle-marked Roller, *Ancylis achatana* (Denis & Schiffermüller), 1775

[version 1; peer review: awaiting peer review]

Douglas Boyes¹⁺, David C. Lees² , Clare Boyes³,
University of Oxford and Wytham Woods Genome Acquisition Lab,
Natural History Museum Genome Acquisition Lab,
Darwin Tree of Life Barcoding collective,
Wellcome Sanger Institute Tree of Life Management, Samples and Laboratory
team,
Wellcome Sanger Institute Scientific Operations: Sequencing Operations,
Wellcome Sanger Institute Tree of Life Core Informatics team,
Tree of Life Core Informatics collective, Darwin Tree of Life Consortium

¹UK Centre for Ecology & Hydrology, Wallingford, England, UK

²Natural History Museum, London, England, UK

³Independent researcher, Welshpool, Wales, UK

+ Deceased author

V1 First published: 14 Mar 2025, 10:128
<https://doi.org/10.12688/wellcomeopenres.23897.1>
Latest published: 14 Mar 2025, 10:128
<https://doi.org/10.12688/wellcomeopenres.23897.1>

Abstract

We present a genome assembly from a male specimen of *Ancylis achatana* (Triangle-marked Roller; Arthropoda; Insecta; Lepidoptera; Tortricidae). The genome sequence has a total length of 547.27 megabases. Most of the assembly (99.25%) is scaffolded into 28 chromosomal pseudomolecules, including the Z sex chromosome. The mitochondrial genome has also been assembled, with a length of 16.0 kilobases.

Keywords

Ancylis achatana, Triangle-marked Roller, genome sequence, chromosomal, Lepidoptera



This article is included in the [Tree of Life](#) gateway.

Open Peer Review

Approval Status Awaiting Peer Review

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Darwin Tree of Life Consortium (mark.blaxter@sanger.ac.uk)

Author roles: **Boyes D:** Investigation, Resources; **Lees DC:** Investigation, Resources; **Boyes C:** Writing – Original Draft Preparation;

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by Wellcome through core funding to the Wellcome Sanger Institute (220540) and the Darwin Tree of Life Discretionary Award [218328, <https://doi.org/10.35802/218328>].

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2025 Boyes D *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Boyes D, Lees DC, Boyes C *et al.* **The genome sequence of the Triangle-marked Roller, *Ancylis achatana* (Denis & Schiffermüller), 1775 [version 1; peer review: awaiting peer review]** Wellcome Open Research 2025, 10:128 <https://doi.org/10.12688/wellcomeopenres.23897.1>

First published: 14 Mar 2025, 10:128 <https://doi.org/10.12688/wellcomeopenres.23897.1>

Species taxonomy

Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Protostomia; Ecdysozoa; Panarthropoda; Arthropoda; Mandibulata; Pancrustacea; Hexapoda; Insecta; Dicondylia; Pterygota; Neoptera; Endopterygota; Amphiesmenoptera; Lepidoptera; Glossata; Neolepidoptera; Heteroneura; Ditrysia; Apoditrysia; Tortricoidea; Tortricidae; Olethreutinae; Enarmoniini; *Ancylis*; *Ancylis achatana* (Denis & Schiffermüller), 1775 (NCBI: txid1869464)

Background

Ancylis achatana (common name Triangle-marked Roller or Hedge Hook-wing) is a micro-moth in the family Tortricidae. It is common in England and Wales and local in Ireland. It is found eastwards across Europe to Russia (GBIF Secretariat, 2025).

The small moth (forewing length 6–9 mm) is mottled brown and grey with a distinctive pre-tornal triangular rich brown patch in the middle of the trailing edge of the forewings. Although the colouration is variable, the markings are consistent. The moth is on the wing between May and August and comes to light (Sterling *et al.*, 2023).

Ancylis achatana lays its eggs on hawthorn, blackthorn or occasionally *Cotoneaster* spp., and the larva spins a tubular shelter from leaves attached to a twig, and then feeds close by (Hancock *et al.*, 2015). It pupates in the leaf spinnings during May (Langmaid *et al.*, 2018).

We present a chromosome-level genome sequence for *Ancylis achatana* based on one male specimen from Wytham Woods, Oxfordshire, UK. The genome was sequenced as part of the Darwin Tree of Life Project, a collaborative effort to sequence all named eukaryotic species in the Atlantic Archipelago of Britain and Ireland.

Genome sequence report

Sequencing data

The genome of a specimen of *Ancylis achatana* (Figure 1) was sequenced using Pacific Biosciences single-molecule HiFi long reads, generating 59.18 Gb (gigabases) from 5.55 million reads. GenomeScope analysis of the PacBio HiFi data estimated the haploid genome size at 575.95 Mb, with a heterozygosity of 0.29% and repeat content of 38.29%. These values provide an initial assessment of genome complexity and the challenges anticipated during assembly. Based on this estimated genome size, the sequencing data provided approximately 100.0x coverage of the genome. Chromosome conformation Hi-C sequencing produced 106.37 Gb from 704.46 million reads. Table 1 summarises the specimen and sequencing information.

Assembly statistics

The primary haplotype was assembled, and contigs corresponding to an alternate haplotype were also deposited in INSDC databases. The assembly was improved by manual curation, which corrected 55 misjoins or missing joins. These



Figure 1. Photograph of the *Ancylis achatana* (ilAncAcha1) specimen used for genome sequencing.

interventions decreased the scaffold count by 5.88%. The final assembly has a total length of 547.27 Mb in 111 scaffolds, with 67 gaps, and a scaffold N50 of 20.35 Mb (Table 2).

The snail plot in Figure 2 provides a summary of the assembly statistics, indicating the distribution of scaffold lengths and other assembly metrics. Figure 3 shows the distribution of scaffolds by GC proportion and coverage. Figure 4 presents a cumulative assembly plot, with separate curves representing different scaffold subsets assigned to various phyla, illustrating the completeness of the assembly.

Most of the assembly sequence (99.26%) was assigned to 28 chromosomal-level scaffolds, representing 27 autosomes and the Z sex chromosome. These chromosome-level scaffolds, confirmed by Hi-C data, are named according to size (Figure 5; Table 3). During curation, chromosome Z was assigned by synteny to *Archips crataeganus* (GCA_947859365.1) (Boyes *et al.*, 2024).

The mitochondrial genome was also assembled. This sequence is included as a contig in the multifasta file of the genome submission and as a standalone record.

Assembly quality metrics

The estimated Quality Value (QV) and *k*-mer completeness metrics, along with BUSCO completeness scores, were calculated for each haplotype and the combined assembly. The QV reflects the base-level accuracy of the assembly, while *k*-mer completeness indicates the proportion of expected *k*-mers identified in the assembly. BUSCO scores provide a measure of completeness based on benchmarking universal single-copy orthologues.

The combined primary and alternate assemblies achieve an estimated QV of 61.9. The *k*-mer recovery for the primary haplotype is 91.92%, and for the alternate haplotype

Table 1. Specimen and sequencing data for *Ancylis achatana*.

Project information			
Study title	Ancyliis achatana (triangle-marked roller)		
Umbrella BioProject	PRJEB78334		
Species	<i>Ancyliis achatana</i>		
BioSpecimen	SAMEA7520689		
NCBI taxonomy ID	1869464		
Specimen information			
Technology	ToLID	BioSample accession	Organism part
PacBio long read sequencing	ilAncAcha1	SAMEA7520779	whole organism
Hi-C sequencing	ilAncAcha2	SAMEA114806094	whole organism
Sequencing information			
Platform	Run accession	Read count	Base count (Gb)
Hi-C Illumina NovaSeq X	ERR13389762	7.04e+08	106.37
PacBio Revio	ERR13420778	5.55e+06	59.18

90.69%; the combined primary and alternate assemblies have a k -mer recovery of 99.17%. BUSCO analysis using the lepidoptera_odb10 reference set ($n = 5,286$) identified 98.4% of the expected gene set (single = 98.0%, duplicated = 0.5%).

Table 2 provides assembly metric benchmarks adapted from Rhie *et al.* (2021) and the Earth BioGenome Project (EBP) Report on Assembly Standards September 2024. The assembly achieves the EBP reference standard of **6.C.Q61**.

Methods

Sample acquisition and DNA barcoding

An adult male *Ancyliis achatana* (specimen ID Ox000473, ToLID ilAncAcha1) was collected from Wytham Woods, Oxfordshire, United Kingdom (latitude 51.77, longitude -1.34) on 2020-06-13 by light trap. The specimen was collected and identified by Douglas Boyes (University of Oxford) and preserved on dry ice.

The specimen used for Hi-C sequencing (specimen ID NHMUK014584877, ToLID ilAncAcha2) was collected from Bure Marshes Stables, England, United Kingdom (latitude 52.69, longitude 1.45) on 2022-06-29. The specimen was collected by David Lees (Natural History Museum) and identified by Mark Sterling (Natural History Museum) and preserved by dry freezing (-80 °C).

The initial identification was verified by an additional DNA barcoding process according to the framework developed by Twyford *et al.* (2024). A small sample was dissected from the specimen and stored in ethanol, while the

remaining parts were shipped on dry ice to the Wellcome Sanger Institute (WSI) (Pereira *et al.*, 2022). The tissue was lysed, the COI marker region was amplified by PCR, and amplicons were sequenced and compared to the BOLD database, confirming the species identification (Crowley *et al.*, 2023). Following whole genome sequence generation, the relevant DNA barcode region was also used alongside the initial barcoding data for sample tracking at the WSI (Twyford *et al.*, 2024). The standard operating procedures for Darwin Tree of Life barcoding have been deposited on protocols.io (Beasley *et al.*, 2023).

Metadata collection for samples adhered to the Darwin Tree of Life project standards described by Lawniczak *et al.* (2022).

Nucleic acid extraction

The workflow for high molecular weight (HMW) DNA extraction at the Wellcome Sanger Institute (WSI) Tree of Life Core Laboratory includes a sequence of procedures: sample preparation and homogenisation, DNA extraction, fragmentation and purification. Detailed protocols are available on protocols.io (Denton *et al.*, 2023b). The ilAncAcha1 sample was prepared for DNA extraction by weighing and dissecting it on dry ice (Jay *et al.*, 2023). Tissue from the whole organism was homogenised using a PowerMasher II tissue disruptor (Denton *et al.*, 2023a).

HMW DNA was extracted using the Automated MagAttract v2 protocol (Oatley *et al.*, 2023a). DNA was sheared into an average fragment size of 12–20 kb in a Megaruptor 3 system (Bates *et al.*, 2023). Sheared DNA was purified by solid-phase

Table 2. Genome assembly data for *Ancyliis achatana*.

Genome assembly		
Assembly name	ilAncAcha1.1	
Assembly accession	GCA_964214005.1	
Alternate haplotype accession	GCA_964214015.1	
Assembly level for primary assembly	chromosome	
Span (Mb)	547.27	
Number of contigs	178	
Number of scaffolds	111	
Longest scaffold (Mb)	42.27	
Assembly metric	Measure	Benchmark
Contig N50 length	9.22 Mb	≥ 1 Mb
Scaffold N50 length	20.35 Mb	= chromosome N50
Consensus quality (QV)	Primary: 63.0; alternate: 61.1; combined 61.9	≥ 40
<i>k</i> -mer completeness	Primary: 91.92%; alternate: 90.69%; combined: 99.17%	≥ 95%
BUSCO*	C:98.4%[S:98.0%,D:0.5%], F:0.2%,M:1.3%,n:5,286	<i>S</i> > 90%; <i>D</i> < 5%
Percentage of assembly mapped to chromosomes	99.26%	≥ 90%
Sex chromosomes	Z	localised homologous pairs
Organelles	Mitochondrial genome: 16.0 kb	complete single alleles

* BUSCO scores based on the lepidoptera_odb10 BUSCO set using version 5.5.0. C = complete [S = single copy, D = duplicated], F = fragmented, M = missing, n = number of orthologues in comparison.

reversible immobilisation, using AMPure PB beads to eliminate shorter fragments and concentrate the DNA (Oatley *et al.*, 2023b). The concentration of the sheared and purified DNA was assessed using a Nanodrop spectrophotometer and Qubit Fluorometer using the Qubit dsDNA High Sensitivity Assay kit. Fragment size distribution was evaluated by running the sample on the FemtoPulse system.

Hi-C sample preparation

Tissue from the whole organism of the ilAncAcha2 sample was processed for Hi-C sequencing at the WSI Scientific Operations core, using the Arima-HiC v2 kit. In brief, 20–50 mg of frozen tissue (stored at –80 °C) was fixed, and the DNA crosslinked using a TC buffer with 22% formaldehyde concentration. After crosslinking, the tissue was homogenised using the Diagenode Power Masher-II and BioMasher-II tubes and pestles. Following the Arima-HiC v2 kit manufacturer's instructions, crosslinked DNA was digested using

a restriction enzyme master mix. The 5'-overhangs were filled in and labelled with biotinylated nucleotides and proximally ligated. An overnight incubation was carried out for enzymes to digest remaining proteins and for crosslinks to reverse. A clean up was performed with SPRIselect beads prior to library preparation. Additionally, the biotinylation percentage was estimated using the Qubit Fluorometer v4.0 (Thermo Fisher Scientific) and Qubit HS Assay Kit and Arima-HiC v2 QC beads.

Library preparation and sequencing

Library preparation and sequencing were performed at the WSI Scientific Operations core.

PacBio HiFi

At a minimum, samples were required to have an average fragment size exceeding 8 kb and a total mass over 400 ng to proceed to the low input SMRTbell Prep Kit 3.0




Figure 3. Genome assembly of *Ancylis achatana*, ilAncAcha1.1: BlobToolKit GC-coverage plot. Blob plot showing sequence coverage (vertical axis) and GC content (horizontal axis). The circles represent scaffolds, with the size proportional to scaffold length and the colour representing phylum membership. The histograms along the axes display the total length of sequences distributed across different levels of coverage and GC content. An interactive version of this figure is available at https://blobtoolkit.genomehubs.org/view/GCA_964214005.1/blob.

Biosciences, California, USA). The SMRT link software, a PacBio web-based end-to-end workflow manager, was used to set-up and monitor the run, as well as perform primary and secondary analysis of the data upon completion.

Hi-C

For Hi-C library preparation, DNA was fragmented using the Covaris E220 sonicator (Covaris) and size selected using SPRISelect beads to 400 to 600 bp. The DNA was then enriched using the Arima-HiC v2 kit Enrichment beads. Using the NEBNext Ultra II DNA Library Prep Kit (New England Biolabs) for end repair, a-tailing, and adapter ligation. This uses a custom protocol which resembles the standard NEBNext Ultra II DNA Library Prep protocol but where library preparation occurs while DNA is bound to the Enrichment beads. For library amplification, 10 to 16 PCR cycles were required, determined by the sample

biotinylation percentage. The Hi-C sequencing was performed using paired-end sequencing with a read length of 150 bp on an Illumina NovaSeq X instrument.

Genome assembly, curation and evaluation

Assembly

Prior to assembly of the PacBio HiFi reads, a database of k -mer counts ($k = 31$) was generated from the filtered reads using FastK. GenomeScope2 (Ranallo-Benavidez *et al.*, 2020) was used to analyse the k -mer frequency distributions, providing estimates of genome size, heterozygosity, and repeat content.

The HiFi reads were first assembled using Hifiasm (Cheng *et al.*, 2021) with the --primary option. Haplotypic duplications were identified and removed using purge_dups (Guan *et al.*, 2020). The Hi-C reads were mapped to the primary




Figure 4. Genome assembly of *Ancyliis achatana*, ilAncAcha1.1: BlobToolKit cumulative sequence plot. The grey line shows cumulative length for all scaffolds. Coloured lines show cumulative lengths of scaffolds assigned to each phylum using the buscogenes taxrule. An interactive version of this figure is available at https://blobtoolkit.genomehubs.org/view/GCA_964214005.1/dataset/GCA_964214005.1/cumulative.

contigs using bwa-mem2 (Vasimuddin *et al.*, 2019). The contigs were further scaffolded using the provided Hi-C data (Rao *et al.*, 2014) in YaHS (Zhou *et al.*, 2023) using the --break option for handling potential misassemblies. The scaffolded assemblies were evaluated using Gfastats (Formenti *et al.*, 2022), BUSCO (Manni *et al.*, 2021) and MERQURY.FK (Rhie *et al.*, 2020).

The mitochondrial genome was assembled using MitoHiFi (Uliano-Silva *et al.*, 2023), which runs MitoFinder (Allio *et al.*, 2020) and uses these annotations to select the final

mitochondrial contig and to ensure the general quality of the sequence.

Assembly curation

The assembly was decontaminated using the Assembly Screen for Cobionts and Contaminants (ASCC) pipeline. Flat files and maps used in curation were generated via the TreeVal pipeline (Pointon *et al.*, 2023). Manual curation was conducted primarily in PretextView (Harry, 2022) and HiGlass (Kerpedjiev *et al.*, 2018), with additional insights provided by JBrowse2 (Diesh *et al.*, 2023). Scaffolds were visually inspected and




Figure 5. Genome assembly of *Ancylistis achatana*: Hi-C contact map of the iAncAcha1.1 assembly, visualised using HiGlass. Chromosomes are shown in order of size from left to right and top to bottom. An interactive version of this figure may be viewed at https://genome-note-higlass.tol.sanger.ac.uk/l/?d=fNieo9pLR92pE1Di_iUfgA.

Table 3. Chromosomal pseudomolecules in the genome assembly of *Ancylistis achatana*, iAncAcha1.

INSDC accession	Name	Length (Mb)	GC%
OZ172862.1	1	32.02	38
OZ172863.1	2	29.49	37.5
OZ172864.1	3	22.82	38.5
OZ172865.1	4	22.72	38.5
OZ172866.1	5	21.89	38
OZ172867.1	6	21.81	38.5
OZ172868.1	7	21.7	38
OZ172869.1	8	21.61	38
OZ172870.1	9	20.52	38
OZ172871.1	10	20.35	38.5
OZ172872.1	11	20.04	38.5
OZ172873.1	12	19.98	39
OZ172874.1	13	19.72	38.5

INSDC accession	Name	Length (Mb)	GC%
OZ172875.1	14	19.48	38.5
OZ172876.1	15	19.42	38.5
OZ172877.1	16	18.85	38.5
OZ172878.1	17	17.99	39
OZ172879.1	18	17.9	39
OZ172880.1	19	16.71	39
OZ172881.1	20	14.5	38.5
OZ172882.1	21	13.21	39.5
OZ172883.1	22	13.01	39
OZ172884.1	23	12.66	41
OZ172885.1	24	11.92	39
OZ172886.1	25	10.76	39.5
OZ172887.1	26	9.98	39
OZ172888.1	27	9.87	40.5
OZ172861.1	Z	42.27	38
OZ172889.1	MT	0.02	19

corrected as described by [Howe et al. \(2021\)](#). Any identified contamination, missed joins, and mis-joins were amended, and duplicate sequences were tagged and removed. Sex chromosomes were identified by synteny analysis. The curation process is documented at <https://gitlab.com/wtsi-grit/rapid-curation>.

Assembly quality assessment

The Merqury.FK tool ([Rhie et al., 2020](#)), run in a Singularity container ([Kurtzer et al., 2017](#)), was used to evaluate *k*-mer completeness and assembly quality for the primary and alternate haplotypes using the *k*-mer databases ($k = 31$) that were computed prior to genome assembly. The analysis outputs included assembly QV scores and completeness statistics.

A Hi-C contact map was produced for the final version of the assembly. The Hi-C reads were aligned using bwa-mem2 ([Vasimuddin et al., 2019](#)) and the alignment files were combined using SAMtools ([Danecek et al., 2021](#)). The Hi-C alignments were converted into a contact map using BEDTools ([Quinlan & Hall, 2010](#)) and the Cooler tool suite ([Abdennur & Mirny, 2020](#)). The contact map was visualised in HiGlass ([Kerpedjiev et al., 2018](#)).

The blobtoolkit pipeline is a Nextflow port of the previous Snakemake Blobtoolkit pipeline ([Challis et al., 2020](#)). It aligns the PacBio reads in SAMtools and minimap2 ([Li, 2018](#)) and generates coverage tracks for regions of fixed size. In parallel, it queries the GoT database ([Challis et al., 2023](#)) to

identify all matching BUSCO lineages to run BUSCO ([Manni et al., 2021](#)). For the three domain-level BUSCO lineages, the pipeline aligns the BUSCO genes to the UniProt Reference Proteomes database ([Bateman et al., 2023](#)) with DIAMOND blastp ([Buchfink et al., 2021](#)). The genome is also divided into chunks according to the density of the BUSCO genes from the closest taxonomic lineage, and each chunk is aligned to the UniProt Reference Proteomes database using DIAMOND blastx. Genome sequences without a hit are chunked using seqtk and aligned to the NT database with blastn ([Altschul et al., 1990](#)). The blobtools suite combines all these outputs into a blobdir for visualisation.

The blobtoolkit pipeline was developed using nf-core tooling ([Ewels et al., 2020](#)) and MultiQC ([Ewels et al., 2016](#)), relying on the Conda package manager, the Bioconda initiative ([Grüning et al., 2018](#)), the Biocontainers infrastructure ([da Veiga Leprevost et al., 2017](#)), as well as the Docker ([Merkel, 2014](#)) and Singularity ([Kurtzer et al., 2017](#)) containerisation solutions.

Table 4 contains a list of relevant software tool versions and sources.

Wellcome Sanger Institute – Legal and Governance

The materials that have contributed to this genome note have been supplied by a Darwin Tree of Life Partner. The submission of materials by a Darwin Tree of Life Partner is subject to the **‘Darwin Tree of Life Project Sampling Code**

Table 4. Software tools: versions and sources.

Software tool	Version	Source
BEDTools	2.30.0	https://github.com/arq5x/bedtools2
BLAST	2.14.0	ftp://ftp.ncbi.nlm.nih.gov/blast/executables/blast+/ /
BlobToolKit	4.3.9	https://github.com/blobtoolkit/blobtoolkit
BUSCO	5.5.0	https://gitlab.com/ezlab/busco
bwa-mem2	2.2.1	https://github.com/bwa-mem2/bwa-mem2
Cooler	0.8.11	https://github.com/open2c/cooler
DIAMOND	2.1.8	https://github.com/bbuchfink/diamond
fasta_windows	0.2.4	https://github.com/tolkit/fasta_windows
FastK	666652151335353eef2fcd58880bcef5bc2928e1	https://github.com/thegenemyers/FASTK
Gfastats	1.3.6	https://github.com/vgl-hub/gfastats
GoT CLI	0.2.5	https://github.com/genomehubs/goat-cli
Hifiasm	0.19.8-r603	https://github.com/chhy1p123/hifiasm
HiGlass	44086069ee7d4d3f6f3f0012569789ec138f42b84a a44357826c0b6753eb28de	https://github.com/higlass/higlass
MerquryFK	d00d98157618f4e8d1a9190026b19b471055b22e	https://github.com/thegenemyers/MERQURY.FK
Minimap2	2.24-r1122	https://github.com/lh3/minimap2
MitoHiFi	3	https://github.com/marcelauliano/MitoHiFi

Software tool	Version	Source
MultiQC	1.14, 1.17, and 1.18	https://github.com/MultiQC/MultiQC
NCBI Datasets	15.12.0	https://github.com/ncbi/datasets
Nextflow	23.10.0	https://github.com/nextflow-io/nextflow
PretextView	0.2.5	https://github.com/sanger-tol/PretextView
samtools	1.19.2	https://github.com/samtools/samtools
sanger-tol/ascc	-	https://github.com/sanger-tol/ascc
sanger-tol/blobtoolkit	0.5.1	https://github.com/sanger-tol/blobtoolkit
Seqtk	1.3	https://github.com/lh3/seqtk
Singularity	3.9.0	https://github.com/sylabs/singularity
TreeVal	1.2.0	https://github.com/sanger-tol/treeval
YaHS	1.2a.2	https://github.com/c-zhou/yahs

of Practice⁹, which can be found in full on the Darwin Tree of Life website [here](#). By agreeing with and signing up to the Sampling Code of Practice, the Darwin Tree of Life Partner agrees they will meet the legal and ethical requirements and standards set out within this document in respect of all samples acquired for, and supplied to, the Darwin Tree of Life Project.

Further, the Wellcome Sanger Institute employs a process whereby due diligence is carried out proportionate to the nature of the materials themselves, and the circumstances under which they have been/are to be collected and provided for use. The purpose of this is to address and mitigate any potential legal and/or ethical implications of receipt and use of the materials as part of the research project, and to ensure that in doing so we align with best practice wherever possible. The overarching areas of consideration are:

- Ethical review of provenance and sourcing of the material
- Legality of collection, transfer and use (national and international)

Each transfer of samples is further undertaken according to a Research Collaboration Agreement or Material Transfer Agreement entered into by the Darwin Tree of Life Partner, Genome Research Limited (operating as the Wellcome Sanger Institute), and in some circumstances other Darwin Tree of Life collaborators.

Data availability

European Nucleotide Archive: *Ancylis achatana* (triangle-marked roller). Accession number PRJEB78334; <https://identifiers.org/ena.embl/PRJEB78334>. The genome sequence is released openly for reuse. The *Ancylis achatana* genome

sequencing initiative is part of the Darwin Tree of Life (DTOL) project. All raw sequence data and the assembly have been deposited in INSDC databases. The genome will be annotated using available RNA-Seq data and presented through the **Ensembl** pipeline at the European Bioinformatics Institute. Raw data and assembly accession identifiers are reported in [Table 1](#) and [Table 2](#).

Author information

Members of the University of Oxford and Wytham Woods Genome Acquisition Lab are listed here: <https://doi.org/10.5281/zenodo.12157525>.

Members of the Natural History Museum Genome Acquisition Lab are listed here: <https://doi.org/10.5281/zenodo.12159242>.

Members of the Darwin Tree of Life Barcoding collective are listed here: <https://doi.org/10.5281/zenodo.12158331>.

Members of the Wellcome Sanger Institute Tree of Life Management, Samples and Laboratory team are listed here: <https://doi.org/10.5281/zenodo.12162482>.

Members of Wellcome Sanger Institute Scientific Operations: Sequencing Operations are listed here: <https://doi.org/10.5281/zenodo.12165051>.

Members of the Wellcome Sanger Institute Tree of Life Core Informatics team are listed here: <https://doi.org/10.5281/zenodo.12160324>.

Members of the Tree of Life Core Informatics collective are listed here: <https://doi.org/10.5281/zenodo.12205391>.

Members of the Darwin Tree of Life Consortium are listed here: <https://doi.org/10.5281/zenodo.4783558>.

References

- Abdennur N, Mirny LA: **Cooler: scalable storage for Hi-C data and other genomically labeled arrays.** *Bioinformatics.* 2020; **36**(1): 311–316.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Allio R, Schomaker-Bastos A, Romiguier J, *et al.*: **MitoFinder: efficient automated large-scale extraction of mitogenomic data in target enrichment phylogenomics.** *Mol Ecol Resour.* 2020; **20**(4): 892–905.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Altschul SF, Gish W, Miller W, *et al.*: **Basic Local Alignment Search Tool.** *J Mol Biol.* 1990; **215**(3): 403–410.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bateman A, Martin MJ, Orchard S, *et al.*: **UniProt: the universal protein knowledgebase in 2023.** *Nucleic Acids Res.* 2023; **51**(D1): D523–D531.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bates A, Clayton-Lucey I, Howard C: **Sanger Tree of Life HMW DNA fragmentation: diagenode Megaruptor[®]3 for LI PacBio.** *protocols.io.* 2023.
[Publisher Full Text](#)
- Beasley J, Uhl R, Forrest LL, *et al.*: **DNA barcoding SOPs for the Darwin Tree of Life project.** *protocols.io.* 2023; [Accessed 25 June 2024].
[Publisher Full Text](#)
- Boyes D, Gibbs M, University of Oxford and Wytham Woods Genome Acquisition Lab, *et al.*: **The genome sequence of the Brown Oak Tortrix, *Archips crataeganus* (Hübner, 1796) [version 1; peer review: 3 approved, 1 approved with reservations].** *Wellcome Open Res.* 2024; **9**: 9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Buchfink B, Reuter K, Drost HG: **Sensitive protein alignments at Tree-of-Life scale using DIAMOND.** *Nat Methods.* 2021; **18**(4): 366–368.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Challis R, Kumar S, Sotero-Caio C, *et al.*: **Genomes on a Tree (GoAT): a versatile, scalable search engine for genomic and sequencing project metadata across the eukaryotic Tree of Life [version 1; peer review: 2 approved].** *Wellcome Open Res.* 2023; **8**: 24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Challis R, Richards E, Rajan J, *et al.*: **BlobToolKit – interactive quality assessment of genome assemblies.** *G3 (Bethesda).* 2020; **10**(4): 1361–1374.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cheng H, Concepcion GT, Feng X, *et al.*: **Haplotype-resolved *de novo* assembly using phased assembly graphs with hifiasm.** *Nat Methods.* 2021; **18**(2): 170–175.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Crowley L, Allen H, Barnes I, *et al.*: **A sampling strategy for genome sequencing the British terrestrial arthropod fauna [version 1; peer review: 2 approved].** *Wellcome Open Res.* 2023; **8**: 123.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- da Veiga Leprevost F, Grüning BA, Alves Aflitos S, *et al.*: **BioContainers: an open-source and community-driven framework for software standardization.** *Bioinformatics.* 2017; **33**(16): 2580–2582.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Danecek P, Bonfield JK, Liddle J, *et al.*: **Twelve years of SAMtools and BCFtools.** *GigaScience.* 2021; **10**(2): g1ab008.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Denton A, Oatley G, Cornwell C, *et al.*: **Sanger Tree of Life sample homogenisation: PowerMash.** *protocols.io.* 2023a.
[Publisher Full Text](#)
- Denton A, Yatsenko H, Jay J, *et al.*: **Sanger Tree of Life wet laboratory protocol collection V.1.** *protocols.io.* 2023b.
[Publisher Full Text](#)
- Diesh C, Stevens GJ, Xie P, *et al.*: **JBrowse 2: a modular genome browser with views of synteny and structural variation.** *Genome Biol.* 2023; **24**(1): 74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ewels PA, Peltzer A, Fillinger S, *et al.*: **The nf-core framework for community-curated bioinformatics pipelines.** *Nat Biotechnol.* 2020; **38**(3): 276–278.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ewels P, Magnusson M, Lundin S, *et al.*: **MultiQC: summarize analysis results for multiple tools and samples in a single report.** *Bioinformatics.* 2016; **32**(19): 3047–3048.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Formenti G, Abueg L, Brajuka A, *et al.*: **Gfastats: conversion, evaluation and manipulation of genome sequences using assembly graphs.** *Bioinformatics.* 2022; **38**(17): 4214–4216.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- GBIF Secretariat: ***Ancylis achatana* (Denis & Schiffermüller), 1775 checklist dataset.** *GBIF Backbone Taxonomy.* 2025; [Accessed 8 February 2025].
[Reference Source](#)
- Grüning B, Dale R, Sjödin A, *et al.*: **Bioconda: sustainable and comprehensive software distribution for the life sciences.** *Nat Methods.* 2018; **15**(7): 475–476.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Guan D, McCarthy SA, Wood J, *et al.*: **Identifying and removing haplotypic duplication in primary genome assemblies.** *Bioinformatics.* 2020; **36**(9): 2896–2898.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hancock EF, Bland KP, Razowski J: **The Moths and Butterflies of Great Britain and Ireland Tortricidae Olethreutinae.** Leiden: Brill, 2015.
- Harry E: **PretextView (Paired REad TEXTure Viewer): a desktop application for viewing pretext contact maps.** 2022.
[Reference Source](#)
- Howe K, Chow W, Collins J, *et al.*: **Significantly improving the quality of genome assemblies through curation.** *GigaScience.* 2021; **10**(1): g1aa153.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Jay J, Yatsenko H, Narváez-Gómez JP, *et al.*: **Sanger Tree of Life sample preparation: triage and dissection.** *protocols.io.* 2023.
[Publisher Full Text](#)
- Kerpedjiev P, Abdennur N, Lekschas F, *et al.*: **HiGlass: web-based visual exploration and analysis of genome interaction maps.** *Genome Biol.* 2018; **19**(1): 125.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kurtzer GM, Sochat V, Bauer MW: **Singularity: scientific containers for mobility of compute.** *PLoS One.* 2017; **12**(5): e0177459.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Langmaid JR, Palmer S, Young MR: **A field guide to the smaller moths of Great Britain and Ireland.** 3rd ed., British Entomological and Natural History Society, 2018.
[Reference Source](#)
- Lawniczak MKN, Davey RP, Rajan J, *et al.*: **Specimen and sample metadata standards for biodiversity genomics: a proposal from the Darwin Tree of Life project [version 1; peer review: 2 approved with reservations].** *Wellcome Open Res.* 2022; **7**: 187.
[Publisher Full Text](#)
- Li H: **Minimap2: pairwise alignment for nucleotide sequences.** *Bioinformatics.* 2018; **34**(18): 3094–3100.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Manni M, Berkeley MR, Seppely M, *et al.*: **BUSCO update: novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes.** *Mol Biol Evol.* 2021; **38**(10): 4647–4654.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Merkel D: **Docker: lightweight Linux containers for consistent development and deployment.** *Linux J.* 2014; **2014**(239): 2, [Accessed 2 April 2024].
[Reference Source](#)
- Oatley G, Denton A, Howard C: **Sanger Tree of Life HMW DNA extraction: automated MagAttract v.2.** *protocols.io.* 2023a.
[Publisher Full Text](#)
- Oatley G, Sampaio F, Howard C: **Sanger Tree of Life fragmented DNA clean up: automated SPRI.** *protocols.io.* 2023b.
[Publisher Full Text](#)
- Pereira L, Sivell O, Sivess L, *et al.*: **DTOL: taxon-specific standard operating procedure for the terrestrial and freshwater arthropods working group.** 2022.
[Publisher Full Text](#)
- Pointon DL, Eagles W, Sims Y, *et al.*: **sanger-tol/treeval v1.0.0 – Ancient Atlantis.** 2023.
[Reference Source](#)
- Quinlan AR, Hall IM: **BEDTools: a flexible suite of utilities for comparing genomic features.** *Bioinformatics.* 2010; **26**(6): 841–842.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ranallo-Benavidez TR, Jaron KS, Schatz MC: **GenomeScope 2.0 and smudgeplot for reference-free profiling of polyploid genomes.** *Nat Commun.* 2020; **11**(1): 1432.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rao SSP, Huntley MH, Durand NC, *et al.*: **A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping.** *Cell.* 2014; **159**(7): 1665–1680.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rhie A, McCarthy SA, Fedrigo O, *et al.*: **Towards complete and error-free genome assemblies of all vertebrate species.** *Nature.* 2021; **592**(7856): 737–746.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rhie A, Walenz BP, Koren S, *et al.*: **Mercury: reference-free quality, completeness, and phasing assessment for genome assemblies.** *Genome Biol.* 2020; **21**(1): 245.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sterling P, Parsons M, Lewington R: **Field guide to the micro-moths of Great Britain and Ireland.** Second Edition. London: Bloomsbury Publishing, 2023.
[Reference Source](#)
- Twyford AD, Beasley J, Barnes I, *et al.*: **A DNA barcoding framework for**

taxonomic verification in the Darwin Tree of Life project [version 1; peer review: 2 approved]. *Wellcome Open Res.* 2024; **9**: 339.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Ulliano-Silva M, Ferreira JGRN, Krashennikova K, *et al.*: **MitoHiFi: a python pipeline for mitochondrial genome assembly from PacBio high fidelity reads.** *BMC Bioinformatics.* 2023; **24**(1): 288.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Vasimuddin M, Misra S, Li H, *et al.*: **Efficient architecture-aware acceleration of BWA-MEM for multicore systems.** In: *2019 IEEE International Parallel and Distributed Processing Symposium (IPDPS)*. IEEE, 2019; 314–324.
[Publisher Full Text](#)

Zhou C, McCarthy SA, Durbin R: **YaHS: yet another Hi-C scaffolding tool.** *Bioinformatics.* 2023; **39**(1): btac808.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)