

British Antarctic Survey
NATURAL ENVIRONMENT RESEARCH COUNCIL

NERC SCIENCE OF THE ENVIRONMENT | UK Research and Innovation

1:500 000 Scale

Geological Map of Alexander Island

BAS GEOMAP 2 Series, Sheet 8, Edition 1

Geological interpretation and map compilation by T.R. Riley, M.L. Curtis, J.A. Crame, D.J. Cantrill and D.I.M. Macdonald. Geological cross section compiled by T.R. Riley. Data preparation, digital cartography, design and layout by T.R. Riley and M.L. Curtis.

Base map data for coastlines, rock outcrops and ice shelves from the SCAR Antarctic digital database, accessed 2023.

Preferred reference for map:
Riley, T.R., Curtis, M.L., Crame, J.A., Cantrill, D.J. and Macdonald, D.I.M. (2025). Geological map of Alexander Island, Antarctic Peninsula (1:500 000 scale). BAS GEOMAP 2 Series, sheet 8, British Antarctic Survey, Cambridge, UK.

Published by British Antarctic Survey, High Cross, Madingley Road, Cambridge, CB3 0ET, United Kingdom.

GEOLOGICAL LEGEND

Geological units are dark coloured (exposed geology) and light tints (inferred geology under ice or snow)

PERIOD	GROUP	UNIT	VOLCANIC AND SEDIMENTARY ROCKS	MAGMATIC AND METAMORPHIC ROCKS
QUATERNARY / NEOGENE	Beethoven Peninsula Volcanic Group	20	**Beethoven Peninsula Volcanic Field** c.2.5 Ma - 0.1 Ma. Basaltic lavas, pillow lavas, lapilli tuff and tuff breccia. Lava-fed delta and subaqueous tuff cone lithofacies.	
		19	**Mount Pinafore Volcanic Field** c.7.7 Ma - 5.4 Ma. Basaltic lavas, lapilli tuff, hyaloclastite breccia and tuff breccia.	
PALEOGENE	Alexander Island Volcanic Group	18	**Finlandia Formation** c.50 Ma - 46 Ma. Basaltic and basaltic andesite lavas, and rare high-Mg andesite flows.	
		17	**Elgar Formation** c.54 Ma - 49 Ma. Basaltic and basaltic andesite lavas, rare high-Mg andesite flows. Bedded tuffs and agglomerates. Up to 2000 m thick.	
		15	**Colbert Formation** c.65 Ma - 62 Ma. Rhyolitic to dacitic ignimbrite, crystal-lithic tuffs, breccia and volcaniclastic units. Total thickness is >2000 m.	
		14	**Staccato Magmatic Complex** c.82 Ma - 71 Ma. Massive andesite lavas on the southern coast of Monteverdi Peninsula. Diorite intrusion at Staccato Peaks.	
			Emplacement of allochthonous mélange belt and translation of Charcot Island associated with Late Cretaceous plate reorganisation.	
CRETACEOUS	Fossil Bluff Group	13	**Mars Glacier Member** Sandstone, subordinate mudstone and conglomerate. Fossil forest horizons. Up to 1000 m thickness. c.102 Ma.	
		12	**Triton Peak Member** Coarse sandstone and conglomerate. Fossil forest horizons and rare tuff beds. Up to 900 m thickness. c.105 Ma.	
		11	**Deimos Ridge Member** Sandstone with interbedded mudstone. Bed thickness increases upwards. Up to 700 m thickness. c.114 Ma.	
		8 -8a	**Pluto Glacier Formation** Mudstone-siltstone succession with rare sandstone-conglomerate beds. Up to 800 m thickness. 8a: Rhea Corner Member conglomerates. c. 126 Ma.	
		7	**Spartan Glacier Formation** Monotonous succession of mudstone and siltstone. Up to 1000 m in thickness. c. 130 Ma.	
		6 -6a	**Himalia Ridge Formation** Basal succession of mudstone and upper 1000 m arenite/rudite dominated. Rare rift-related basaltic units. Up to 2600 m in thickness. c. 140 Ma. 6a: Jupiter Glacier Member - 90 m sandstone beds representing an abrupt shallowing event.	
JURASSIC		5	**Ablation Point Formation** Slumped blocks of sandstone and interbedded sandstone/mudstone. rare volcanic beds. Up to 440 m in thickness. c. 155 Ma.	
		4	**Atoll Nunataks Formation** Thinly bedded green and black siliceous mudstone with rare granule conglomerates. Up to 1050 m in thickness. c. 177 Ma.	
		3	**Selene Nunatak Formation** Conglomerate and sandstone units derived from the accretionary complex. Fluvial succession at Zebra Ridge. Up to 150 m in thickness. c. 182 Ma.	
		2	**Mount King Beds** Basal mudstone units transition to abundant sandstone-conglomerate units. Locally, mudstone units preserve a diverse macrofauna. Up to 1000 m. c. 182 Ma.	
PERMIAN	LeMay Group	1a	**LeMay Group (accretionary prism)** Succession of variably deformed trench-fill and trench-slope turbidites. Up to 4000 m in thickness. c. 255 Ma. Accretion at c. 230 Ma.	
				Rouen Intrusive Complex c.56 Ma. At least five mappable intrusions ranging from granite to diorite emplaced at shallow crustal levels. G: granite; Gd: granodiorite; To: tonalite; D: diorite.
		15 -16		**LeMay Group (mélange belt)** Thrust-bound allochthonous slices of blueschist facies oceanic crust, mudstones and radiolarian-bearing cherts. Possible seamount at Lully Foothills (1c). Likely emplacement age of c. 90 Ma.
		10		**Charcot Island Beds** Weakly deformed sandstone-conglomerate-shale beds. uPossible unexposed gabbroic basement (Charcot Anomaly). c. 112 Ma.
		9		**Gannon Nunataks Volcanic Beds** Coarse grained volcaniclastic beds unconformably overlying the LeMay Group accretionary complex. c. 118 Ma.

GEOLOGICAL SYMBOLS

- Inferred contact
- Unconformity
- Normal fault (tick on downthrown side)
- Reverse fault (teeth on hanging wall)
- Strike-slip fault
- Axial trace of antiform
- Axial trace of synform
- Bedding (e.g. 38)
- Cleavage (e.g. 78)
- Overturned bedding (e.g. 15)
- Laterally continuous bedding
- Dyke
- Fossil forest locality
- Significant plant fossil locality
- Significant invertebrate fossil locality
- Radiolaria-bearing cherts

OTHER SYMBOLS

- Coastline (adjacent to ice shelf)
- Coastline (adjacent to ocean)
- Contours (100 m intervals)
- Ice shelf

REFERENCES

Armstrong, D. C. (1993). Geological investigation of the Debussy Heights and Rouen Mountains areas, northern Alexander Island. Geology field report. ADS/2R/1992/GL5. British Antarctic Survey unpublished report.

Butterworth, P. J. (1984). Jurassic/Lower Cretaceous rocks at Ablation Valley and Belemnite Point, Alexander Island. ADS/2R/1983/G11. British Antarctic Survey unpublished report.

Butterworth, P. J., Crame, J. A., Howlett, P. J., & Macdonald, D. I. M. (1988). Lithostratigraphy of Upper Jurassic-Lower Cretaceous strata of eastern Alexander Island. Cretaceous Research, 9, 249–264.

Butterworth, P. J. (1991). The role of eustasy in the development of a regional shallowing event in a tectonically active basin, Fossil Bluff Group (Jurassic–Cretaceous), Alexander Island, Antarctica. In: Macdonald, D. I. M. (ed.) *Sedimentation, Tectonics and Eustasy: Sea Level Changes at Active Margins. Special Publication of the International Association of Sedimentologists* **12**, 307–329.

Burn, R. W. (1984). The geology of the LeMay Group, Alexander Island. British Antarctic Survey Scientific Reports, 109, British Antarctic Survey, Cambridge.

Crame, J. A. (1978). Palaeontological studies in south-eastern Alexander Island, 1977–78. ADS/2R/1977/G6. British Antarctic Survey unpublished report.

Crame, J. A. (1985). Stratigraphical studies in the Calliste Cliffs-Planet Heights area of eastern Alexander Island. ADS/2R/1985/G12. British Antarctic Survey unpublished report.

Crame, J. A., & Howlett, P. J. (1988). Late Jurassic and early Cretaceous biostratigraphy of the Fossil Bluff Formation, Alexander Island. British Antarctic Survey Bulletin, 78, 1–35.

Doubleday, P. A., & Storey, B. C. (1998). Deformation history of a Mesozoic forearc basin sequence on Alexander Island, Antarctic Peninsula. Journal of South American Earth Sciences, 11, 1–21.

Doubleday, P. A., Macdonald, D. I. M., & Nell, P. A. R. (1993). Sedimentology and structure of the trench-slope to fore-arc basin transition in the Mesozoic of Alexander Island, Antarctica, Geological Magazine, 130, 737–754.

Doubleday, P. A., Leat, P. T., Alabaster, T., Nell , P. A. R., & Tranter, T. H. (1994). Allochthonous oceanic basalts within the Mesozoic accretionary complex of Alexander Island, Antarctica: remnants of proto-Pacific oceanic crust. Journal of the Geological Society, London, 151, 65–78.

Edwards, C. W. (1980). The geology of central and eastern Alexander Island. Ph.D. thesis, University of Birmingham. 228 pp.

Hyden G., & Tanner P. W. G. (1981). Late-Palaeozoic–early Mesozoic fore-arc basin sedimentary rocks at the Pacific margin in Western Antarctica. *Geologische Rundshau*, 70, 529–541.

Holdsworth, B. K., & Nell, P. A. R. (1992). Mesozoic radiolarian faunas from the Antarctic Peninsula: age, tectonic and palaeoceanographic significance. *Journal of the Geological Society, London*, 149, 1003-1020.

Hole, M. J. (1986). Alkaline volcanism project 1985/86. Field report. ADS/2R/1985/G9. British Antarctic Survey unpublished report.

Howlett, P. J. (1986). Preliminary geological report 1985/86. Biostratigraphy of the Ablation Valley/Ganymede Heights area, Alexander Island. ADS/2R/1985/G4. British Antarctic Survey unpublished report.

Kelly, S. R. A. (1988). Geological field work on the Fossil Bluff Group in SE Alexander Island, 1988/89. Field report. ADS/2R/1988/G2. British Antarctic Survey unpublished report.

Kelly, S. R. A., Doubleday, P. A., Brunton, C. H. C., Dickins, J. M., Sevastopulo, G. D., & Taylor, P. D. (2001). First Carboniferous and ?Permian marine macrofaunas from Antarctica and their tectonic implications, Journal of the Geological Society, London, 158(2), 219–232.

Larter, R. D., & Barker, P. F. (1991). Effects of ridge crest–trench interaction on Antarctic–Phoenix spreading: forces on a young subducting plate. Journal of Geophysical Research, 96, 19583–19607.

Macdonald, D. I. M., Moncrieff, A .C. M. & Butterworth, P. J. (1993). Giant slide deposits from a Mesozoic fore-arc basin, Alexander Island, Antarctica. Geology, 21(11): 1047-1050.

Macdonald, D. I. M. (1987). Geological field report. Sedimentological studies in north eastern Alexander Island. Field report. ADS/2R/1986/GS. British Antarctic Survey unpublished report.

Macdonald, D. I. M., Butterworth, P. J., Crame, J. A. . (1995) Deep marine slide and channel deposits from the Jurassic-Cretaceous Fossil Bluff Group, Alexander Island, Antarctica. In: Pickering, K.T., Hiscott, R.N., Kenyon, N.H., Ricci-Lucchi, F., Smith, R.D.A. (eds.). Atlas of Deep Water Environments: architectural style in turbidite systems, London, Chapman and Hall, 50–55. https://doi.org/10.1007/978-94-011-1234-5_9.

Macdonald, D. I. M., Leat, P. T., Doubleday, P. A., & Kelly, S. R. A. (1999). On the origin of fore-arc basins: new evidence of formation by rifting from the Jurassic of Alexander Island, Antarctica. Terra Nova, 11, 186–193.

Matthews, K. J., Seton, M., & Müller, R. D. (2012). A global-scale plate reorganization event at 105–100 Ma: Earth and Planetary Science Letters, 355, 283–298.

McCarron, J. J. (1991). The geology of the Colbert Mountains, central Alexander Island. Field report. ADS/2R/1990/GL7. British Antarctic Survey unpublished report.

McCarron, J. J. (1992). Geological investigations in Elgar Uplands and Finlandia Foothills, northern Alexander Island. ADS/2R/1991/GL5. British Antarctic Survey unpublished report.

McCarron, J. J., & Larter, R. D. (1998). Late Cretaceous to early Tertiary subduction history of the Antarctic Peninsula, Journal of the Geological Society, London, 155, 255–268.

Miller, S., & Macdonald, D. I. M. (2004). Metamorphic and thermal history of a fore-arc basin: the Fossil Bluff Group, Alexander Island, Antarctica. *Journal of Petrology*, 45 (7), 1453–1465. https://doi.org/10.1093/petrology/egh025

Moncrieff, A. C. M. (1988). Field report of sedimentological studies in south eastern Alexander Island, 1988/89. ADS/2R/1988/G4. British Antarctic Survey unpublished report.

Moncrieff, A. C. M., & Kelly, S. R. A. (1993). Lithostratigraphy of the uppermost Fossil Bluff Group (Early Cretaceous) of Alexander Island, Antarctica: history of an Albian regression. Cretaceous Research, 14, 1–15.

Nell, P. A. R. (1986). Field report on the geology of parts of the eastern LeMay Range and northern Alexander Island. ADS/2R/1985/G11. British Antarctic Survey unpublished report.

Nell, P. A. R. (1987). Field report on the geology of parts of northern Alexander Island and the Fossil Bluff area. 1986–1987 season. ADS/2R/1986/GS. British Antarctic Survey unpublished report.

Nell, P. A. R. (1990). Deformation in an accretionary melange, Alexander Island, Antarctica. In J. Kriper & H. Ruttere (Eds.) *Deformation Mechanisms, Rheology and Tectonics. Geological Society Special Publications*, 54, pp. 405–416.

Nell, P. A. R., & Storey, B. C. (1991). Strike-slip tectonics within the Antarctic Peninsula fore-arc. In *Geological Evolution of Antarctica* (eds. Thomson, M. R. A., Crame, J. A., & Thomson, J.W.), pp. 443-8. Cambridge: Cambridge University Press.

Nichols, G. J. & Cantrill, D. J. (2003). Tectonic and climatic controls on a Mesozoic forearc basin succession, Alexander Island, Antarctica. Geological Magazine, 139, 313-330.

Riley, T. R., Flowerdew, M. J. & Whitehouse, M. J. (2012). Litho- and chronostratigraphy of a fore- to intra-arc basin: Adelaide Island, Antarctic Peninsula. *Geological Magazine*, 149, 768–782.

Riley, T. R., Millar, I. L., Carter, A., Flowerdew, M. J., Burton-Johnson, A., Bastias, J., Storey, C. D., Castillo, P., Chew, D., & Whitehouse, M. J. (2023). Evolution of an accretionary complex (LeMay Group) and terrane translation in the Antarctic Peninsula. *Tectonics*. https://doi.org/10.1029/2022TC007578

Riley, T. R., Flowerdew, M. J, Carter, A., Curtis, M. L., Millar, I. L., Crame, J. A. & Whitehouse, M. J. (2024). Tracking the tempo of a continental margin arc: insights from a forearc succession in West Antarctica. Geological Society of America Bulletin. https://doi.org/10.1130/B37558.1

Smellie, J.L., & Hole, M.J. (2021). Antarctic Peninsula: volcanology. In: J. L. Smellie J.L., et al.. (Eds.) Volcanism in Antarctica: 200 million years of Subduction, Rifting and Continental Break-up. Geology Society London Memoir, 55, 305-325.

Storey, B. C., Brown, R. W., Carter, A., Doubleday, P. A., Hurford, A. J., Macdonald, D. I. M., & Nell, P. A. R. (1996). Fission-track evidence for the thermotectonic evolution of a Mesozoic – Cenozoic fore-arc, Antarctica. Journal of Geological Society, London, 153, 65–82.

Storey, B. C. & Bradshaw, J. D. (1995). Geology of parts of the LeMay Group, Alexander Island. Field report 1994/95 season. ADS/2R/1994/GL3. British Antarctic Survey unpublished report.

Thomson, M. R. A. (1965). Geological report for East Coast Alexander Island, summer 1964. ADS/27/1964/G1. British Antarctic Survey unpublished report.

Thomson, M. R. A. (1979). Upper Jurassic and Lower Cretaceous ammonite faunas of the Ablation Point area, Alexander Island. British Antarctic Survey Scientific Reports 97, 37 pp.

Thomson, M.R.A., & Tranter, T.H. (1986). Early Jurassic fossils from central Alexander Island and their geological setting: British Antarctic Survey, Bulletin, 70, 23–39.

Thomson, M. R. A. & Pankhurst, R. J. (1983). Age of post-Gondwanian calc-alkaline volcanism in the Antarctic Peninsula region. In *Antarctic Earth Science* (eds. R. L. Oliver, P. R. James and J. B. Jago), pp. 328–33. Canberra: Australian Academy of Science.

Tranter, T. H. (1985). Field report on the geology of parts of the LeMay Group of central Alexander Island. ADS/2R/1984/GS. British Antarctic Survey unpublished report.

Tranter, T. H. (1987). The structural history of the LeMay Group of central Alexander Island, Antarctic Peninsula. British Antarctic Survey Bulletin, 77, 61–80.

Tranter, T. H. (1988). The Tectonostratigraphic history of the LeMay Group if central Alexander Island, Antarctica. Ph.D. Thesis, Council for National Academic Awards, pp. 272.

Tranter, T. H. (1991). Accretion and subduction processes along the Pacific margin of Gondwana, central Alexander Island. In: M. R. A. Thomson et al. (Eds.) Geological Evolution of Antarctica. Cambridge University Press, Cambridge, pp. 437–441.

Twinn, G., Riley, T. R., Fox, M., & Carter, A. (2022). Thermal history of the southern Antarctic Peninsula during Cenozoic oblique subduction. Journal of the Geological Society, London, https://doi.org/10.1144/jgs2022-008.