



DATA NOTE

The genome sequence of the Small Dotted Buff moth,

Photedes minima Haworth, 1809

[version 1; peer review: 2 approved]

Douglas Boyes¹⁺, Liam M. Crowley^{id}², Timilehin Adewumi^{id}³,
University of Oxford and Wytham Woods Genome Acquisition Lab,
Darwin Tree of Life Barcoding collective,
Wellcome Sanger Institute Tree of Life Management, Samples and Laboratory
team,
Wellcome Sanger Institute Scientific Operations: Sequencing Operations,
Wellcome Sanger Institute Tree of Life Core Informatics team,
Tree of Life Core Informatics collective, Darwin Tree of Life Consortium

¹UK Centre for Ecology & Hydrology, Wallingford, England, UK

²Department of Biology, University of Oxford, Oxford, England, UK

³Tree of Life, Wellcome Sanger Institute, Hinxton, England, UK

+ Deceased author

V1 First published: 02 Jun 2025, 10:299
<https://doi.org/10.12688/wellcomeopenres.24264.1>
Latest published: 02 Jun 2025, 10:299
<https://doi.org/10.12688/wellcomeopenres.24264.1>

Abstract

We present a genome assembly from a male specimen of *Photedes minima* (Small Dotted Buff; Arthropoda; Insecta; Lepidoptera; Noctuidae). The genome sequence has a total length of 694.66 megabases. Most of the assembly (99.95%) is scaffolded into 31 chromosomal pseudomolecules, including the Z sex chromosome. The mitochondrial genome has also been assembled, with a length of 15.38 kilobases.

Keywords



Photedes minima, Small Dotted Buff, genome sequence, chromosomal, Lepidoptera



This article is included in the [Tree of Life](#) gateway.

Open Peer Review

Approval Status

	1	2
version 1		
02 Jun 2025	view	view

1. **Željko Tomanović**, Serbia University of Belgrade, Belgrade, Serbia

2. **Sivasankaran Kuppasamy**^{id}, Loyola College, Chennai, India

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Darwin Tree of Life Consortium (mark.blaxter@sanger.ac.uk)

Author roles: Boyes D: Investigation, Resources; Crowley LM: Investigation, Resources; Adewumi T: Writing – Original Draft Preparation;

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by Wellcome through core funding to the Wellcome Sanger Institute (220540) and the Darwin Tree of Life Discretionary Award [218328, <https://doi.org/10.35802/218328>].

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2025 Boyes D *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Boyes D, Crowley LM, Adewumi T *et al.* **The genome sequence of the Small Dotted Buff moth, *Photedes minima* Haworth, 1809 [version 1; peer review: 2 approved]** Wellcome Open Research 2025, 10:299 <https://doi.org/10.12688/wellcomeopenres.24264.1>

First published: 02 Jun 2025, 10:299 <https://doi.org/10.12688/wellcomeopenres.24264.1>

Species taxonomy

Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Protostomia; Ecdysozoa; Panarthropoda; Arthropoda; Mandibulata; Pancrustacea; Hexapoda; Insecta; Dicondylia; Pterygota; Neoptera; Endopterygota; Amphiesmenoptera; Lepidoptera; Glossata; Neolepidoptera; Heteroneura; Ditrysia; Obtectomera; Noctuoidea; Noctuidae; Noctuinae; Apameini; *Photedes*; *Photedes minima* Haworth, 1809 (NCBI:txid988148)

Background

Photedes minima, or the Small Dotted Buff, is a moth from the Noctuidae family with a wingspan of 20–26 mm (Kimber, 2025). Males typically have a plain buff appearance with faint dark dots and a slight sheen, while females are smaller with markings that are more distinct. Their forewings range from pale ochreous to light brown, often dusted with darker shades, and females usually have a darker median area. The hindwings vary from yellow-orange to greyish near the edges (Waring *et al.*, 2017).

Distributed across Europe, from the Iberian Peninsula to western Russia (GBIF Secretariat, 2023), *P. minima* is found in marshes, damp woodlands, and wet meadows (Kimber, 2025). In the UK, it is widespread across Britain and Ireland (Waring *et al.*, 2017). Its reliance on moist environments ties closely to the availability of its larval host plant, tufted hair-grass (*Deschampsia cespitosa*) (Kimber, 2025).

The moth has one generation per year, with adults active from late June to August in a single brood (Waring *et al.*, 2017). After mating, females lay eggs near host plants and the larvae feed on grass stems before entering diapause over winter. They pupate in an underground cocoon.

We present a chromosome-level genome sequence for *Photedes minima*, based on a specimen from Wytham Woods, Oxfordshire, United Kingdom (Figure 1). The genome was sequenced as part of the Darwin Tree of Life Project and will provide insights into its evolutionary biology and inform conservation strategies (Blaxter *et al.*, 2022).

Genome sequence report

Sequencing data

The genome of a specimen of *Photedes minima* (Figure 1) was sequenced using Pacific Biosciences single-molecule HiFi long reads, generating 43.48 Gb (gigabases) from 3.17 million reads, which were used to assemble the genome. GenomeScope analysis estimated the haploid genome size at 683.69 Mb, with a heterozygosity of 0.63% and repeat content of 39.57%. These estimates guided expectations for the assembly. Based on the estimated genome size, the sequencing data provided approximately 60 coverage. Hi-C sequencing produced 125.82 Gb from 833.27 million reads, used to scaffold the assembly. RNA sequencing data were also generated and are available in public sequence repositories. Table 1 summarises the specimen and sequencing details.



Figure 1. Photograph of the *Photedes minima* (ilPhoMini1) specimen used for genome sequencing.

Assembly statistics

The primary haplotype was assembled, and contigs corresponding to an alternate haplotype were also deposited in INSDC databases. The assembly was improved by manual curation, which corrected 12 misjoins or missing joins and removed 5 haplotypic duplications. These interventions decreased the scaffold count by 4.76%. The final assembly has a total length of 694.66 Mb in 39 scaffolds, with 8 gaps, and a scaffold N50 of 24.27 Mb (Table 2).

The snail plot in Figure 2 provides a summary of the assembly statistics, indicating the distribution of scaffold lengths and other assembly metrics. Figure 3 shows the distribution of scaffolds by GC proportion and coverage. Figure 4 presents a cumulative assembly plot, with separate curves representing different scaffold subsets assigned to various phyla, illustrating the completeness of the assembly.

Most of the assembly sequence (99.95%) was assigned to 31 chromosomal-level scaffolds, representing 30 autosomes and the Z sex chromosome. These chromosome-level scaffolds, confirmed by Hi-C data, are named according to size (Figure 5; Table 3). During curation, the Z chromosome was identified based BUSCO gene painting with ancestral Merian elements (Wright *et al.*, 2024).

The mitochondrial genome was also assembled. This sequence is included as a contig in the multifasta file of the genome submission and as a standalone record.

Assembly quality metrics

The estimated Quality Value (QV) and *k*-mer completeness metrics, along with BUSCO completeness scores, were calculated for each haplotype and the combined assembly. The QV reflects the base-level accuracy of the assembly, while

Table 1. Specimen and sequencing data for *Photedes minima*.

Project information			
Study title	Photedes minima (small dotted buff)		
Umbrella BioProject	PRJEB71302		
Species	<i>Photedes minima</i>		
BioSpecimen	SAMEA7701533		
NCBI taxonomy ID	988148		
Specimen information			
Technology	ToLID	BioSample accession	Organism part
PacBio long read sequencing	ilPhoMini1	SAMEA7701719	whole organism
Hi-C sequencing	ilPhoMini2	SAMEA114645394	whole organism
RNA sequencing	ilPhoMini2	SAMEA114645394	whole organism
Sequencing information			
Platform	Run accession	Read count	Base count (Gb)
Hi-C Illumina NovaSeq X	ERR13702755	8.33e+08	125.82
PacBio Sequel IIe	ERR12370428	1.36e+06	16.62
PacBio Sequel IIe	ERR12370427	1.81e+06	26.86
RNA Illumina NovaSeq X	ERR13999093	9.15e+07	13.82

k-mer completeness indicates the proportion of expected *k*-mers identified in the assembly. BUSCO scores provide a measure of completeness based on benchmarking universal single-copy orthologues.

The combined primary and alternate assemblies achieve an estimated QV of 67.1. The *k*-mer completeness is 84.63% for the primary haplotype and 84.13% for the alternate haplotype; and 98.68% for the combined primary and alternate assemblies. BUSCO v.5.5.0 analysis using the lepidoptera_odb10 reference set ($n = 5,286$) identified 99.0% of the expected gene set (single = 98.6%, duplicated = 0.5%).

Table 2 provides assembly metric benchmarks adapted from Rhie *et al.* (2021) and the Earth BioGenome Project Report on Assembly Standards September 2024. The primary assembly achieves the EBP reference standard of 7.C.Q68.

Methods

Sample acquisition and DNA barcoding

The specimen used for genome sequencing was an adult male *Photedes minima* (specimen ID Ox000672, ToLID ilPho-Mini1), collected from Wytham Woods, Oxfordshire, United Kingdom (latitude 51.772, longitude -1.338) on 2020-07-20, using a light trap. The specimen was collected and identified by Douglas Boyes (University of Oxford) and preserved on dry ice. A second specimen was used for Hi-C and RNA

sequencing (specimen ID Ox003713, ToLID ilPhoMini2), was collected from the same location on 2023-06-20. The specimen was collected and identified by Liam Crowley (University of Oxford) and preserved on dry ice.

The initial identification was verified by an additional DNA barcoding process according to the framework developed by Twyford *et al.* (2024). A small sample was dissected from each specimen and stored in ethanol, while the remaining parts were shipped on dry ice to the Wellcome Sanger Institute (WSI) (Pereira *et al.*, 2022). The tissue was lysed, the COI marker region was amplified by PCR, and amplicons were sequenced and compared to the BOLD database, confirming the species identification (Crowley *et al.*, 2023). Following whole genome sequence generation, the relevant DNA barcode region was also used alongside the initial barcoding data for sample tracking at the WSI (Twyford *et al.*, 2024). The standard operating procedures for Darwin Tree of Life barcoding have been deposited on protocols.io (Beasley *et al.*, 2023).

Metadata collection for samples adhered to the Darwin Tree of Life project standards described by Lawniczak *et al.* (2022).

Nucleic acid extraction

The workflow for high molecular weight (HMW) DNA extraction at the Wellcome Sanger Institute (WSI) Tree of Life Core

Table 2. Genome assembly data for *Photodes minima*.

Genome assembly		
Assembly name	ilPhoMini1.1	
Assembly accession	GCA_964662195.1	
Alternate haplotype accession	GCA_964662255.1	
Assembly level for primary assembly	chromosome	
Span (Mb)	694.66	
Number of contigs	47	
Number of scaffolds	39	
Longest scaffold (Mb)	33.31	
Assembly metric	Measure	Benchmark
Contig N50 length	24.18 Mb	≥ 1 Mb
Scaffold N50 length	24.27 Mb	= chromosome N50
Consensus quality (QV)	Primary: 68.6; alternate: 66.5; combined: 67.1	≥ 40
k-mer completeness	Primary: 84.63%; alternate: 84.13%; combined: 98.68%	$\geq 95\%$
BUSCO*	C:99.0%[S:98.6%,D:0.5%], F:0.1%,M:0.9%,n:5,286	$S > 90\%$; $D < 5\%$
Percentage of assembly assigned to chromosomes	99.95%	$\geq 90\%$
Sex chromosomes	Z	localised homologous pairs
Organelles	Mitochondrial genome: 15.38 kb	complete single alleles

* BUSCO scores based on the lepidoptera_odb10 BUSCO set using version 5.5.0. C = complete [S = single copy, D = duplicated], F = fragmented, M = missing, n = number of orthologues in comparison.

Laboratory includes a sequence of procedures: sample preparation and homogenisation, DNA extraction, fragmentation and purification (Howard *et al.*, 2025). Detailed protocols are available on protocols.io (Denton *et al.*, 2023b). The ilPho-Mini1 sample was prepared for DNA extraction by weighing and dissecting it on dry ice (Jay *et al.*, 2023). Tissue from the whole organism was homogenised using a PowerMasher II tissue disruptor (Denton *et al.*, 2023a). HMW DNA was extracted using the Automated MagAttract v1 protocol (Sheerin *et al.*, 2023). DNA was sheared into an average fragment size of 12–20 kb in a Megaruptor 3 system (Todorovic *et al.*, 2023). Sheared DNA was purified by solid-phase reversible immobilisation, using AMPure PB beads to eliminate shorter fragments and concentrate the DNA (Strickland *et al.*, 2023). The concentration of the sheared and purified DNA was assessed using a Nanodrop spectrophotometer and a Qubit Fluorometer using the Qubit dsDNA High Sensitivity Assay kit. The fragment size distribution was evaluated by running the sample on the FemtoPulse system.

RNA was extracted from whole organism tissue of ilPho-Mini2 in the Tree of Life Laboratory at the WSI using the RNA Extraction: Automated MagMax™ mirVana protocol (do Amaral *et al.*, 2023). The RNA concentration was assessed using a Nanodrop spectrophotometer and a Qubit Fluorometer using the Qubit RNA Broad-Range Assay kit. Analysis of the integrity of the RNA was done using the Agilent RNA 6000 Pico Kit and Eukaryotic Total RNA assay.

Hi-C sample preparation and crosslinking

Hi-C data were generated from the whole organism of the ilPhoMini2 sample using the Arima-HiC v2 kit (Arima Genomics) with 20–50 mg of frozen tissue (stored at -80°C). As per manufacturer's instructions, tissue was fixed, and the DNA crosslinked using a TC buffer with 22% formaldehyde concentration, and a final formaldehyde concentration of 2%. The tissue was then homogenised using the Diagnocine Power Masher-II. The crosslinked DNA was digested using a restriction enzyme master mix, then biotinylated




Figure 2. Genome assembly of *Photedes minima*, ilPhoMini1.1: metrics. The BlobToolKit snail plot provides an overview of assembly metrics and BUSCO gene completeness. The circumference represents the length of the whole genome sequence, and the main plot is divided into 1,000 bins around the circumference. The outermost blue tracks display the distribution of GC, AT, and N percentages across the bins. Scaffolds are arranged clockwise from longest to shortest and are depicted in dark grey. The longest scaffold is indicated by the red arc, and the deeper orange and pale orange arcs represent the N50 and N90 lengths. A light grey spiral at the centre shows the cumulative scaffold count on a logarithmic scale. A summary of complete, fragmented, duplicated, and missing BUSCO genes in the lepidoptera_odb10 set is presented at the top right. An interactive version of this figure is available at https://blobtoolkit.genomehubs.org/view/GCA_964662195.1/dataset/GCA_964662195.1/snail.

and ligated. A clean up was performed with SPRIselect beads prior to library preparation. DNA concentration was quantified using the Qubit Fluorometer v4.0 (Thermo Fisher Scientific) and Qubit HS Assay Kit, and sample biotinylation percentage was estimated using the Arima-HiC v2 QC beads.

Library preparation and sequencing

Library preparation and sequencing were performed at the WSI Scientific Operations core.

PacBio HiFi

At a minimum, samples were required to have an average fragment size exceeding 8 kb and a total mass over 400 ng to proceed to the low-input SMRTbell Prep Kit 3.0 protocol (Pacific Biosciences), depending on genome size and sequencing depth required. Libraries were prepared using the SMRTbell Prep Kit 3.0 as per the manufacturer's instructions. The kit

includes the reagents required for end repair/A-tailing, adapter ligation, post-ligation SMRTbell bead cleanup, and nuclease treatment. Size-selection and clean-up were carried out using diluted AMPure PB beads (Pacific Biosciences). DNA concentration was quantified using the Qubit Fluorometer v4.0 (ThermoFisher Scientific) with Qubit 1X dsDNA HS assay kit and the final library fragment size analysis was carried out using the Agilent Femto Pulse Automated Pulsed Field CE Instrument (Agilent Technologies) and the gDNA 55kb BAC analysis kit.

Samples were sequenced using the Sequel IIe system (Pacific Biosciences, California, USA). The concentration of the library loaded onto the Sequel IIe was in the range 40–135 pM. The SMRT link software, a PacBio web-based end-to-end workflow manager, was used to set-up and monitor the run, as well as perform primary and secondary analysis of the data upon completion.




Figure 3. Genome assembly of *Photedes minima*, ilPhoMini1.1: BlobToolKit GC-coverage plot. Blob plot showing sequence coverage (vertical axis) and GC content (horizontal axis). The circles represent scaffolds, with the size proportional to scaffold length and the colour representing phylum membership. The histograms along the axes display the total length of sequences distributed across different levels of coverage and GC content. An interactive version of this figure is available at https://blobtoolkit.genomehubs.org/view/GCA_964662195.1/dataset/GCA_964662195.1/blob.

Hi-C

For Hi-C library preparation, the biotinylated DNA constructs were fragmented using a Covaris E220 sonicator and size-selected to 400–600 bp using SPRISelect beads. DNA was then enriched using Arima-HiC v2 Enrichment beads. The NEBNext Ultra II DNA Library Prep Kit (New England Biolabs) was used for end repair, A-tailing, and adapter ligation, following a modified protocol in which library preparation is carried out while the DNA remains bound to the enrichment beads. PCR amplification was performed using KAPA HiFi HotStart mix and custom dual-indexed adapters (Integrated DNA Technologies) in a 96-well plate format. Depending on sample concentration and biotinylation percentage determined at the crosslinking stage, samples were amplified for 10–16 PCR cycles. Post-PCR clean-up was carried out using SPRISelect beads. The libraries were quantified using the Accuclear Ultra High Sensitivity dsDNA Standards Assay kit (Biotium) and normalised to 10 ng/μL before sequencing. Hi-C sequencing was performed on the Illumina NovaSeq X instrument using 150 bp paired-end reads.

RNA

Poly(A) RNA-Seq libraries were constructed using the NEB Ultra II RNA Library Prep kit, following the manufacturer's

instructions. RNA sequencing was performed on the Illumina NovaSeq X instrument.

Genome assembly, curation and evaluation

Assembly

Prior to assembly of the PacBio HiFi reads, a database of k -mer counts ($k = 31$) was generated from the filtered reads using FastK. GenomeScope2 (Ranallo-Benavidez *et al.*, 2020) was used to analyse the k -mer frequency distributions, providing estimates of genome size, heterozygosity, and repeat content.

The HiFi reads were first assembled using Hifiasm (Cheng *et al.*, 2021) with the --primary option. Haplotypic duplications were identified and removed using purge_dups (Guan *et al.*, 2020). The Hi-C reads (Rao *et al.*, 2014) were mapped to the primary contigs using bwa-mem2 (Vasimuddin *et al.*, 2019), and the contigs were scaffolded in YaHS (Zhou *et al.*, 2023) using the --break option for handling potential misassemblies. The scaffolded assemblies were evaluated using Gfastats (Formenti *et al.*, 2022), BUSCO (Manni *et al.*, 2021) and MERQURY.FK (Rhie *et al.*, 2020).

The mitochondrial genome was assembled using Mito-HiFi (Uliano-Silva *et al.*, 2023), which runs MitoFinder




Figure 4. Genome assembly of *Photedes minima*, ilPhoMini1.1: BlobToolKit cumulative sequence plot. The grey line shows cumulative length for all scaffolds. Coloured lines show cumulative lengths of scaffolds assigned to each phylum using the buscogenes taxrule. An interactive version of this figure is available at https://blobtoolkit.genomehubs.org/view/GCA_964662195.1/dataset/GCA_964662195.1/cumulative.




Figure 5. Genome assembly of *Photedes minima*: Hi-C contact map of the ilPhoMini1.1 assembly, generated using PretextSnapshot. Chromosomes are shown in order of size and labelled with chromosome numbers (top) and chromosome accession numbers (left).

Table 3. Chromosomal pseudomolecules in the genome assembly of *Photodes minima*, ilPhoMini1.

INSDC accession	Name	Length (Mb)	GC%
OZ211931.1	Z	33.31	38.5
OZ211932.1	1	28.32	39
OZ211933.1	2	28.12	38.5
OZ211934.1	3	26.43	38.5
OZ211935.1	4	26.32	38.5
OZ211936.1	5	26.16	38.5
OZ211937.1	6	26.05	38.5
OZ211938.1	7	25.87	38.5
OZ211939.1	8	25.7	38.5
OZ211940.1	9	25.6	38.5
OZ211941.1	10	25.0	39
OZ211942.1	11	24.63	38.5
OZ211943.1	12	24.3	38.5
OZ211944.1	13	24.27	38.5
OZ211945.1	14	24.18	38.5
OZ211946.1	15	23.82	38.5
OZ211947.1	16	23.76	38.5
OZ211948.1	17	22.94	39
OZ211949.1	18	22.92	39
OZ211950.1	19	22.57	38.5
OZ211951.1	20	21.02	39
OZ211952.1	21	20.52	39.5
OZ211953.1	22	20.27	38.5
OZ211954.1	23	20.17	39
OZ211955.1	24	19.33	39.5
OZ211956.1	25	16.36	39
OZ211957.1	26	16.31	39
OZ211958.1	27	13.93	40
OZ211959.1	28	13.54	40
OZ211960.1	29	12.16	39.5
OZ211961.1	30	10.45	41
OZ211962.1	MT	0.02	20.5

(Allio *et al.*, 2020) and uses these annotations to select the final mitochondrial contig and to ensure the general quality of the sequence.

Assembly curation

The assembly was decontaminated using the Assembly Screen for Cobionts and Contaminants (ASCC) pipeline. Flat files and maps used in curation were generated via the TreeVal pipeline (Pointon *et al.*, 2023). Manual curation was conducted primarily in PretextView (Harry, 2022) and HiGlass (Kerpedjiev *et al.*, 2018), with additional insights provided by JBrowse2 (Diesh *et al.*, 2023). Scaffolds were visually inspected and corrected as described by Howe *et al.* (2021). Any identified contamination, missed joins, and mis-joins were amended, and duplicate sequences were tagged and removed. Sex chromosomes were identified by Merian element painting. The curation process is documented at <https://gitlab.com/wtsi-grit/rapid-curation>.

Assembly quality assessment

The Merqury.FK tool (Rhie *et al.*, 2020), run in a Singularity container (Kurtzer *et al.*, 2017), was used to evaluate *k*-mer completeness and assembly quality for the primary and alternate haplotypes using the *k*-mer databases ($k = 31$) computed prior to genome assembly. The analysis outputs included assembly QV scores and completeness statistics.

The genome was analysed in the blobtoolkit pipeline, a Nextflow (Di Tommaso *et al.*, 2017) port of the previous Snake-make Blobtoolkit pipeline (Challis *et al.*, 2020). It aligns the PacBio reads in SAMtools (Danecek *et al.*, 2021) and minimap2 (Li, 2018) and generates coverage tracks for regions of fixed size. In parallel, it queries the GoAT database (Challis *et al.*, 2023) to identify all matching BUSCO lineages to run BUSCO (Manni *et al.*, 2021). For the three domain-level BUSCO lineages, the pipeline aligns the BUSCO genes to the UniProt Reference Proteomes database (Bateman *et al.*, 2023) with DIAMOND blastp (Buchfink *et al.*, 2021). The genome is also divided into chunks according to the density of the BUSCO genes from the closest taxonomic lineage, and each chunk is aligned to the UniProt Reference Proteomes database using DIAMOND blastx. Genome sequences without a hit are chunked using seqtk and aligned to the NT database with blastn (Altschul *et al.*, 1990). The blobtools suite combines all these outputs into a blobdir for visualisation.

The blobtoolkit pipeline was developed using nf-core tooling (Ewels *et al.*, 2020) and MultiQC (Ewels *et al.*, 2016), relying on the Conda package manager, the Bioconda initiative (Grüning *et al.*, 2018), the Biocontainers infrastructure (da Veiga Leprevost *et al.*, 2017), as well as the Docker (Merkel, 2014) and Singularity (Kurtzer *et al.*, 2017) containerisation solutions.

Table 4 contains a list of relevant software tool versions and sources.

Wellcome Sanger Institute – Legal and Governance

The materials that have contributed to this genome note have been supplied by a Darwin Tree of Life Partner. The submission

Table 4. Software tools: versions and sources.

Software tool	Version	Source
BLAST	2.14.0	ftp://ftp.ncbi.nlm.nih.gov/blast/executables/blast+/
BlobToolKit	4.3.9	https://github.com/blobtoolkit/blobtoolkit
BUSCO	5.5.0	https://gitlab.com/ezlab/busco
bwa-mem2	2.2.1	https://github.com/bwa-mem2/bwa-mem2
DIAMOND	2.1.8	https://github.com/bbuchfink/diamond
fasta_windows	0.2.4	https://github.com/tolkit/fasta_windows
FastK	666652151335353eef2fcd58880bcef5bc2928e1	https://github.com/thegenemyers/FASTK
Gfastats	1.3.6	https://github.com/vgl-hub/gfastats
Goat CLI	0.2.5	https://github.com/genomehubs/goat-cli
Hifiasm	0.19.8-r603	https://github.com/chhy123/hifiasm
HiGlass	44086069ee7d4d3f6f3f0012569789ec138f42b84a a44357826c0b6753eb28de	https://github.com/higlass/higlass
MercuryFK	d00d98157618f4e8d1a9190026b19b471055b22e	https://github.com/thegenemyers/MERQURY.FK
Minimap2	2.24-r1122	https://github.com/lh3/minimap2
MitoHiFi	3	https://github.com/marcelauliano/MitoHiFi
MultiQC	1.14, 1.17, and 1.18	https://github.com/MultiQC/MultiQC
Nextflow	23.10.0	https://github.com/nextflow-io/nextflow
PretextView	0.2.5	https://github.com/sanger-tol/PretextView
PretextViewSnapshot	-	https://github.com/sanger-tol/PretextViewSnapshot
samtools	1.19.2	https://github.com/samtools/samtools
sanger-tol/ascc	0.1.0	https://github.com/sanger-tol/ascc
sanger-tol/blobtoolkit	0.6.0	https://github.com/sanger-tol/blobtoolkit
Seqtk	1.3	https://github.com/lh3/seqtk
Singularity	3.9.0	https://github.com/sylabs/singularity
TreeVal	1.2.0	https://github.com/sanger-tol/treeval
YaHS	1.2a.2	https://github.com/c-zhou/yahs
GenomeScope2.0	2.0.1	https://github.com/tbenavi1/genomescope2.0

of materials by a Darwin Tree of Life Partner is subject to the **‘Darwin Tree of Life Project Sampling Code of Practice’**, which can be found in full on the Darwin Tree of Life website [here](#). By agreeing with and signing up to the Sampling Code of Practice, the Darwin Tree of Life Partner agrees they will meet the legal and ethical requirements and standards set out within this document in respect of all samples acquired for, and supplied to, the Darwin Tree of Life Project.

Further, the Wellcome Sanger Institute employs a process whereby due diligence is carried out proportionate to the nature of the materials themselves, and the circumstances under which they have been/are to be collected and provided for use. The purpose of this is to address and mitigate any potential

legal and/or ethical implications of receipt and use of the materials as part of the research project, and to ensure that in doing so we align with best practice wherever possible. The overarching areas of consideration are:

- Ethical review of provenance and sourcing of the material
- Legality of collection, transfer and use (national and international)

Each transfer of samples is further undertaken according to a Research Collaboration Agreement or Material Transfer Agreement entered into by the Darwin Tree of Life Partner, Genome Research Limited (operating as the Wellcome Sanger Institute), and in some circumstances other Darwin Tree of Life collaborators.

Data availability

European Nucleotide Archive: *Photedes minima* (small dotted buff). Accession number PRJEB71302; <https://identifiers.org/ena.embl/PRJEB71302>. The genome sequence is released openly for reuse. The *Photedes minima* genome sequencing initiative is part of the Darwin Tree of Life Project (PRJEB40665), the Sanger Institute Tree of Life Programme (PRJEB43745) and Project Psyche (PRJEB71705). All raw sequence data and the assembly have been deposited in INSDC databases. The genome will be annotated using available RNA-Seq data and presented through the [Ensembl](#) pipeline at the European Bioinformatics Institute. Raw data and assembly accession identifiers are reported in [Table 1](#) and [Table 2](#).

Author information

Members of the University of Oxford and Wytham Woods Genome Acquisition Lab are listed here: <https://doi.org/10.5281/zenodo.12157525>.

Members of the Darwin Tree of Life Barcoding collective are listed here: <https://doi.org/10.5281/zenodo.12158331>.

Members of the Wellcome Sanger Institute Tree of Life Management, Samples and Laboratory team are listed here: <https://doi.org/10.5281/zenodo.12162482>.

Members of Wellcome Sanger Institute Scientific Operations: Sequencing Operations are listed here: <https://doi.org/10.5281/zenodo.12165051>.

Members of the Wellcome Sanger Institute Tree of Life Core Informatics team are listed here: <https://doi.org/10.5281/zenodo.12160324>.

Members of the Tree of Life Core Informatics collective are listed here: <https://doi.org/10.5281/zenodo.12205391>.

Members of the Darwin Tree of Life Consortium are listed here: <https://doi.org/10.5281/zenodo.4783558>.

References

- Allio R, Schomaker-Bastos A, Romiguier J, *et al.*: **MitoFinder: efficient automated large-scale extraction of mitogenomic data in target enrichment phylogenomics**. *Mol Ecol Resour.* 2020; **20**(4): 892–905. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Altschul SF, Gish W, Miller W, *et al.*: **Basic Local Alignment Search Tool**. *J Mol Biol.* 1990; **215**(3): 403–410. [PubMed Abstract](#) | [Publisher Full Text](#)
- Bateman A, Martin MJ, Orchard S, *et al.*: **UniProt: the Universal Protein Knowledgebase in 2023**. *Nucleic Acids Res.* 2023; **51**(D1): D523–D531. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Beasley J, Uhl R, Forrest LL, *et al.*: **DNA barcoding SOPs for the Darwin Tree of Life project**. *protocols.io.* 2023; [Accessed 25 June 2024]. [Publisher Full Text](#)
- Blaxter M, Mieszkowska N, Di Palma F, *et al.*: **Sequence locally, think globally: the Darwin Tree of Life project**. *Proc Natl Acad Sci U S A.* 2022; **119**(4): e2115642118. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Buchfink B, Reuter K, Drost HG: **Sensitive protein alignments at Tree-of-Life scale using DIAMOND**. *Nat Methods.* 2021; **18**(4): 366–368. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Challis R, Kumar S, Sotero-Caio C, *et al.*: **Genomes on a Tree (GoaT): a versatile, scalable search engine for genomic and sequencing project metadata across the eukaryotic Tree of Life [version 1; peer review: 2 approved]**. *Wellcome Open Res.* 2023; **8**: 24. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Challis R, Richards E, Rajan J, *et al.*: **BlobToolKit – interactive quality assessment of genome assemblies**. *G3 (Bethesda).* 2020; **10**(4): 1361–1374. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cheng H, Concepcion GT, Feng X, *et al.*: **Haplotype-resolved *de novo* assembly using phased assembly graphs with hifiasm**. *Nat Methods.* 2021; **18**(2): 170–175. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Crowley L, Allen H, Barnes I, *et al.*: **A sampling strategy for genome sequencing the British terrestrial arthropod fauna [version 1; peer review: 2 approved]**. *Wellcome Open Res.* 2023; **8**: 123. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- da Veiga Leprevost F, Grünig BA, Alves Aflitos S, *et al.*: **BioContainers: an open-source and community-driven framework for software standardization**. *Bioinformatics.* 2017; **33**(16): 2580–2582. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Danecek P, Bonfield JK, Liddle J, *et al.*: **Twelve years of SAMtools and BCFtools**. *GigaScience.* 2021; **10**(2): giab008. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Denton A, Oatley G, Cornwell C, *et al.*: **Sanger Tree of Life sample homogenisation: PowerMash**. *protocols.io.* 2023a. [Publisher Full Text](#)
- Denton A, Yatsenko H, Jay J, *et al.*: **Sanger Tree of Life wet laboratory protocol collection V.1**. *protocols.io.* 2023b. [Publisher Full Text](#)
- Di Tommaso P, Chatzou M, Floden EW, *et al.*: **Nextflow enables reproducible computational workflows**. *Nat Biotechnol.* 2017; **35**(4): 316–319. [PubMed Abstract](#) | [Publisher Full Text](#)
- Diesh C, Stevens GJ, Xie P, *et al.*: **JBrowse 2: a modular genome browser with views of synteny and structural variation**. *Genome Biol.* 2023; **24**(1): 74. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- do Amaral RJV, Bates A, Denton A, *et al.*: **Sanger Tree of Life RNA extraction: automated MagMax™ mirVana**. *protocols.io.* 2023. [Publisher Full Text](#)
- Ewels P, Magnusson M, Lundin S, *et al.*: **MultiQC: summarize analysis results for multiple tools and samples in a single report**. *Bioinformatics.* 2016; **32**(19): 3047–3048. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ewels PA, Peltzer A, Fillinger S, *et al.*: **The nf-core framework for community-curated bioinformatics pipelines**. *Nat Biotechnol.* 2020; **38**(3): 276–278. [PubMed Abstract](#) | [Publisher Full Text](#)
- Formenti G, Abueg L, Brajuka A, *et al.*: **Gfastats: conversion, evaluation and manipulation of genome sequences using assembly graphs**. *Bioinformatics.* 2022; **38**(17): 4214–4216. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- GBIF Secretariat: ***Photedes minima* Haworth, 1809**. *GBIF Backbone Taxonomy*. 2023; [Accessed 20 April 2025]. [Reference Source](#)
- Grünig B, Dale R, Sjödin A, *et al.*: **Bioconda: sustainable and comprehensive software distribution for the life sciences**. *Nat Methods.* 2018; **15**(7): 475–476. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Guan D, McCarthy SA, Wood J, *et al.*: **Identifying and removing haplotypic duplication in primary genome assemblies**. *Bioinformatics.* 2020; **36**(9): 2896–2898. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Harry E: **PretextView (Paired REad TEXTure Viewer): a desktop application for viewing pretext contact maps**. 2022. [Reference Source](#)
- Howard C, Denton A, Jackson B, *et al.*: **On the path to reference genomes for all biodiversity: lessons learned and laboratory protocols created in the Sanger Tree of Life core laboratory over the first 2000 species**. *bioRxiv.* 2025. [Publisher Full Text](#)
- Howe K, Chow W, Collins J, *et al.*: **Significantly improving the quality of**

genome assemblies through curation. *GigaScience*. 2021; **10**(1): g1aa153.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Jay J, Yatsenko H, Narváez-Gómez JP, *et al.*: **Sanger Tree of Life sample preparation: triage and dissection.** *protocols.io*. 2023.
[Publisher Full Text](#)

Kerpedjiev P, Abdennur N, Lekschas F, *et al.*: **HiGlass: web-based visual exploration and analysis of genome interaction maps.** *Genome Biol*. 2018; **19**(1): 125.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Kimber I: **Small Dotted Buff *Photedes minima* (Haworth, 1809).** *UKMoths*. 2025; [Accessed 20 April 2025].

[Reference Source](#)

Kurtzer GM, Sochat V, Bauer MW: **Singularity: scientific containers for mobility of compute.** *PLoS One*. 2017; **12**(5): e0177459.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Lawniczak MKN, Davey RP, Rajan J, *et al.*: **Specimen and sample metadata standards for biodiversity genomics: a proposal from the Darwin Tree of Life project [version 1; peer review: 2 approved with reservations].** *Wellcome Open Res*. 2022; **7**: 187.

[Publisher Full Text](#)

Li H: **Minimap2: pairwise alignment for nucleotide sequences.** *Bioinformatics*. 2018; **34**(18): 3094–3100.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Manni M, Berkeley MR, Seppey M, *et al.*: **BUSCO update: novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes.** *Mol Biol Evol*. 2021; **38**(10): 4647–4654.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Merkel D: **Docker: lightweight Linux containers for consistent development and deployment.** *Linux J*. 2014; **2014**(239): 2. [Accessed 2 April 2024].

[Reference Source](#)

Pereira L, Sivell O, Sivess L, *et al.*: **DTOL Taxon-specific Standard Operating Procedure for the terrestrial and freshwater arthropods working group.** 2022.

[Publisher Full Text](#)

Pointon DL, Eagles W, Sims Y, *et al.*: **sanger-tol/treeval v1.0.0 – Ancient Atlantis.** 2023.

[Publisher Full Text](#)

Ranallo-Benavidez TR, Jaron KS, Schatz MC: **GenomeScope 2.0 and Smudgeplot for reference-free profiling of polyploid genomes.** *Nat Commun*. 2020; **11**(1): 1432.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Rao SSP, Huntley MH, Durand NC, *et al.*: **A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping.** *Cell*. 2014; **159**(7): 1665–1680.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Rhie A, McCarthy SA, Fedrigo O, *et al.*: **Towards complete and error-free genome assemblies of all vertebrate species.** *Nature*. 2021; **592**(7856): 737–746.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Rhie A, Walenz BP, Koren S, *et al.*: **Merquy: reference-free quality, completeness, and phasing assessment for genome assemblies.** *Genome Biol*. 2020; **21**(1): 245.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Sheerin E, Sampaio F, Oatley G, *et al.*: **Sanger Tree of Life HMW DNA extraction: automated MagAttract v.1.** *protocols.io*. 2023.

[Publisher Full Text](#)

Strickland M, Cornwell C, Howard C: **Sanger Tree of Life fragmented DNA clean up: manual SPRI.** *protocols.io*. 2023.

[Publisher Full Text](#)

Todorovic M, Sampaio F, Howard C: **Sanger Tree of Life HMW DNA fragmentation: diagenode Megaruptor[®]3 for PacBio HiFi.** *protocols.io*. 2023.

[Publisher Full Text](#)

Twyford AD, Beasley J, Barnes I, *et al.*: **A DNA barcoding framework for taxonomic verification in the Darwin Tree of Life project [version 1; peer review: 2 approved].** *Wellcome Open Res*. 2024; **9**: 339.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Uliano-Silva M, Ferreira JGRN, Krashenninnikova K, *et al.*: **MitoHiFi: a python pipeline for mitochondrial genome assembly from PacBio high fidelity reads.** *BMC Bioinformatics*. 2023; **24**(1): 288.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Vasimuddin M, Misra S, Li H, *et al.*: **Efficient architecture-aware acceleration of BWA-MEM for multicore systems.** In: *2019 IEEE International Parallel and Distributed Processing Symposium (IPDPS)*. IEEE, 2019; 314–324.

[Publisher Full Text](#)

Waring P, Townsend M, Lewington R: **Field guide to the Moths of Great Britain and Ireland: third edition.** Bloomsbury Wildlife Guides, 2017.

[Reference Source](#)

Wright CJ, Stevens L, Mackintosh A, *et al.*: **Comparative genomics reveals the dynamics of chromosome evolution in Lepidoptera.** *Nat Ecol Evol*. 2024; **8**(4): 777–790.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Zhou C, McCarthy SA, Durbin R: **YaHS: yet another Hi-C scaffolding tool.** *Bioinformatics*. 2023; **39**(1): btac808.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:  

Version 1

Reviewer Report 22 July 2025

<https://doi.org/10.21956/wellcomeopenres.26761.r125566>

© 2025 Kuppusamy S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Sivasankaran Kuppusamy 

Loyola College, Chennai, India

Authors have done the chromosome-level genome assembly of Small Dotted Buff moth, *Photedes minima* Haworth, 1809. They generated 183.12 Gb sequence data through different sequencing platforms. Then assembled the sequences and obtained a total length of 696.66 megabases. Authors have identified 31 chromosomal pseudomolecules. The comprehensive data is useful for the phylogenomic studies of moths.

Minor comments

The first sentence of the Background may include the describer's name for the species i.e., *Photedes minima* Haworth, 1809

The second paragraph of the background first sentence may be written as "*P. minima* is distributed across Europe, from the Iberian Peninsula to western Russia (GBIF Secretariate, 2023) and is usually found in marshes, damp woodlands and wet meadows (Kimber, 2025)".

The first sentence of the second paragraph can be modified as "The moth has an univoltine, with adults active from late June to August....."

We present a chromosome-level genome sequence for *P. minima*.....

Throughout the text, the author consistently used the genus' full name, *Photedes minima*. Start with the full genus name once, if possible, then shorten it to *P. minima*.

Authors have produced 183.12 Gb comprehensive sequences through the Hi-C illumina and PacBio platforms. After the assembly they received 694.66 megabases. Through the genome annotation the authors haven't listed the protein-coding genes, non-coding genes and gene transcripts.

The research article was well prepared, and the manuscript meets the necessary scientific standard and is suitable for indexing.

Is the rationale for creating the dataset(s) clearly described?

Yes

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Phylogenetic analysis of Noctuoidea moths using mitogenome sequence

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 07 July 2025

<https://doi.org/10.21956/wellcomeopenres.26761.r125571>

© 2025 Tomanović Ž. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Željko Tomanović

Serbia University of Belgrade, Belgrade, Serbia

Boyes et al. presented a genome assembly of *Photedes minima* (Small Dotted Buff; Arthropoda; Insecta; Lepidoptera; Noctuidae), a species with a wide European distribution. The authors briefly presented the biology and ecology of the species. The genome sequence has a total length of 694.66 megabases, which are distributed over 31 chromosomes, including 30 autosomes and the sex chromosome Z. The mitochondrial genome was also assembled. The authors have presented the assembly statistics and all the methods used in detail. Ms is very well written.

Is the rationale for creating the dataset(s) clearly described?

Yes

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: biodiversity, phylogenetics, ecology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.
