# nature ecology & evolution

Article

# Spatio-temporal variability in forest biodiversity associated with human well-being across socio-economic deprivation gradients

Accepted: 21 May 2025

Published online: 24 June 2025

Check for updates

J. C. Fisher <sup>©</sup><sup>1</sup>⊠, M. Dallimer <sup>©</sup><sup>2,3</sup>, G. E. Austen <sup>©</sup><sup>1</sup>, K. N. Irvine<sup>4</sup>, S. G. Aizlewood <sup>©</sup><sup>1</sup>, P. M. King <sup>©</sup><sup>1,2</sup>, H. A. Jackson <sup>©</sup><sup>5,6</sup>, R. D. Fish <sup>©</sup><sup>1,3</sup> & Z. G. Davies <sup>©</sup><sup>1</sup>

Biodiversity declines are accelerating globally, impacting ecosystem functioning, with consequences for human health. Interactions with biodiversity can be associated with human well-being benefits at the individual level, leading to substantial gains for society when scaled up across populations. However, existing research has not accounted for the species within ecological communities and their effect traits (for example, colours, sounds) that can elicit well-being responses. Many species' effect traits are seasonal, and spatial variation in exposure to ecosystems by different sectors of society can lead to unequal opportunities to gain well-being. Here we use an interdisciplinary analytical approach to explore how the association between forest biodiversity and well-being fluctuates: (1) temporally, between different seasons and (2) spatially, across socio-economic deprivation gradients at a national scale (England and Wales). Species' effect traits and participant well-being were derived through a series of seasonal participatory workshops and questionnaires that incorporated BIO-WELL (a biodiversity-well-being psychometric scale). By generating spatially explicit data, we could examine variability in forest biodiversity associated with human well-being across socio-economic deprivation gradients. Forest species' effect trait richness was spatially heterogeneous, particularly in autumn, spring and summer. Broadleaf forests had greater species' effect trait richness than other categories of forest. Forests with higher species' effect trait richness and forests that were associated with higher self-reported participant well-being were in areas with the least socio-economic deprivation. Forest creation/restoration and nature-health interventions must recognize this ecological and social diversity to ensure initiatives are equitable and socially just.

<sup>1</sup>Durrell Institute of Conservation and Ecology (DICE), School of Anthropology and Conservation, University of Kent, Canterbury, UK. <sup>2</sup>Sustainability Research Institute, School of Earth and Environment, University of Leeds, Leeds, UK. <sup>3</sup>Centre for Environmental Policy, Imperial College London, London, UK. <sup>4</sup>Social, Economic and Geographical Sciences Department, James Hutton Institute, Aberdeen, UK. <sup>5</sup>Woodland Trust, Kempton Way, Grantham, UK. <sup>6</sup>UK Centre for Ecology and Hydrology, Lancaster Environment Centre, Bailrigg, UK. e-mail: J.C.Fisher@kent.ac.uk Biodiversity declines are accelerating globally<sup>1</sup>. This loss of biodiversity is impacting the stability and functioning of ecosystems, with potentially far-reaching consequences for human health and well-being<sup>2</sup>. Well-being is a multidimensional concept, encompassing different contributions to human quality of life<sup>3</sup>. The World Health Organization conceptualizes well-being as "a state of complete, physical, mental and social well-being and not merely the absence of disease or infirmity<sup>\*4</sup>. The multiple domains of well-being encompassed in this definition (bio-, which is physical; psycho-, which is mental, consisting of both cognitive and emotional; and social) comprise the 'biopsychosocial' model of health that originates from integrative medicine<sup>5</sup>. An expanded version of this model, called the biopsychosocial–spiritual model<sup>6,7</sup>, also includes a spiritual domain, conceived as including a connection to something greater than oneself.

Spending time in ecosystems such as forests and wetlands has been linked to a multitude of benefits such as reduced stress, improved cognition and better quality of life<sup>8-10</sup>. Given that well-being predicts mortality and morbidity<sup>11</sup>, scaling up these individual-level gains across entire populations could support the public health sector through substantial avoided societal and healthcare costs (for example, refs. 12,13). Therefore, improving our understanding of how exposure to biodiversity can promote well-being is likely to have widespread implications for both public health and conservation, via initiatives such as nature-based solutions and social ('green') prescribing interventions<sup>14,15</sup>.

Whereas an extensive literature has established that interactions with nature can generate positive well-being responses, this existing body of research generally takes a simplistic approach that relies on homogeneous measures of exposure to 'greenspace' or 'greenness'<sup>9,16</sup>. The role biodiversity plays in delivering improved health has been largely overlooked<sup>17</sup>. This is despite people's engagement with biodiversity within ecosystems being multisensory<sup>18</sup> and influenced by personal and cultural associations<sup>19,20</sup>. Without accounting for biodiversity, and how it is experienced and/or perceived, we may not be able to conserve, restore or create ecosystems that will also generate greater benefits for human health and well-being.

These complex biodiversity–human health relationships can be examined through a functional ecology lens. Some species traits, known as 'effect traits', underpin ecosystem service delivery<sup>21</sup>. For example, mean diameter at breast height of a tree is linked to carbon storage<sup>22</sup>. Likewise, the species' traits that lead to changes in people's well-being can be considered effect traits (for example, the 'calling' sounds of tawny owls (*Strix aluco*) and 'prickly' texture of brambles (*Rubus fruticosus*) eliciting positive and negative well-being, respectively)<sup>17</sup>. Different ecosystems will thus provide different levels of well-being, based on the array of species that occur within the ecological community and the effect traits they support.

Ecosystem impacts on health and well-being fluctuate over time<sup>23</sup>. For instance, grass pollen causes allergies leading to asthma and rhinitis (hay fever), which can be tracked over the course of the year and spatially<sup>24</sup>. Similarly, bird communities alter intra-annually, influencing the supply of cultural ecosystem service benefits<sup>25</sup>. This reflects the seasonality of biodiversity in any given ecosystem, where variations in temperature and precipitation affect resource availability and, subsequently, the presence, abundance and diversity of species and the effect traits they support. Seasonal phenological events themselves, such as leaf senescence in deciduous trees, have also been shown to stimulate positive emotions<sup>26</sup>. Moreover, seasonality also influences how people use ecosystems (for example, ref. 27), due to weather or cultural activities such as participation in holidays and festivals. Despite this, temporal variability is rarely considered in nature–human health research<sup>23</sup>.

Spatial variation in exposure to ecosystems by different sectors of society can lead to unequal opportunities to gain well-being (often referred to as 'environmental health inequalities'). In Europe, for instance, socio-economically deprived groups are often less exposed to green/blue spaces and have a higher prevalence of poor health outcomes<sup>28</sup>. Consequently, they could benefit disproportionately from access to such ecosystems. For example, Mitchell et al.<sup>29</sup>, demonstrated that access to recreational greenspace was positively associated with improved mental well-being across the United Kingdom and more so for those under greater financial strain. However, the evidence base is inconclusive and contradictory (Schüle et al.<sup>30</sup> provides a review). Having a deeper insight into the distribution of species' effect traits within the ecosystems people visit could help disentangle these equivocal findings.

Here we use a novel analytical approach to explore how associations between biodiversity and well-being fluctuate: (1) temporally, between different seasons and (2) spatially, at a national scale (England and Wales) and across socio-economic gradients (Fig. 1). Specifically, we focus on forest ecosystems, which have declined in global land area by over 30% between 1990 and 2015<sup>31</sup>, yet support 80% of terrestrial biodiversity<sup>32</sup>. Temperate forests cover 16% of global land area and are less intact in regions with high human population density and intensive agriculture<sup>33</sup>. Consequently, they are commonly the focus of restoration and creation initiatives, often with the aim of producing 'triple wins' for climate change mitigation, biodiversity and human well-being<sup>34,35</sup>.

We conducted a large, participatory process with a diverse cross-section of the public from England and Wales (Extended Data Fig. 1) in each of the four seasons (autumn, winter, spring and summer). This enabled us to identify which forest species, and their effect traits (colours, sounds, smells, textures and behaviours), were described by participants in relation to both positive and negative well-being at different times of the year<sup>17</sup>. We examined the five domains that constitute the biopsychosocial-spiritual model of health<sup>5,6</sup>: (1) physical (the body and how someone feels physically); (2) emotional (positive and negative mood); (3) cognitive (state of mind); (4) social (perceived connections with others) and (5) spiritual (relationships with something greater than oneself). Hereafter we use the term 'well-being' in relation to people's biopsychosocial-spiritual responses to forest biodiversity. Using species distribution models (SDMs), we created spatio-temporal distributions of species' effect traits. Additionally, we quantified the spatially explicit self-reported well-being responses people derive from forest biodiversity, using a questionnaire that incorporated the biodiversity-well-being psychometric scale BIO-WELL<sup>18</sup> (https://research. kent.ac.uk/bio-well/). We therefore examined associations between biodiversity and human well-being spatio-temporally in two ways: through the species' effect traits and via BIO-WELL (Fig. 1). To investigate environmental health inequalities, we used government data on income- and employment-related deprivation, mapped at the finest spatial resolution that is publicly available (Extended Data Fig. 2). We then coupled the distributions of species' effect traits and BIO-WELL scores with socio-economic deprivation.

## Results

#### Seasonal species' effect traits

We identified 78 species' effect traits that were described by participants as eliciting some form of positive or negative well-being across autumn, winter, spring and summer (Fisher et al.<sup>17</sup> includes the full list of all species' effect traits), associated with the 131 forest species that we could generate SDMs for (that is, they had sufficient fine-scale resolution spatial presence/absence data available and/or model fit was acceptable). Most of these species' effect traits (69) were linked with positive rather than negative (9) well-being, and four were allied to both. The richness of effect traits increased with the number of species, particularly in autumn and winter (Fig. 2).

#### Spatio-temporal distributions of species' effect traits

We combined the SDMs with the number of species' effect traits per species that were described in relation to positive or negative well-being





responses, to create eight spatio-temporal maps, one per season (Fig. 3). These maps showed high levels of spatial heterogeneity, with cumulative species' effect trait richness (the total number of unique effect trait–well-being incidences across all species) ranging from zero to 888 for positive well-being and from zero to 66 for negative well-being.

Hotspots of species' effect traits that elicit positive well-being were apparent across southeast England, broadly coinciding with where broadleaf forest is predominately located (Extended Data Fig. 3). Indeed, we found significantly different cumulative species' effect trait richness between all forest categories (following National Forest Inventory (NFI) definitions; Supplementary Tables 1 and 2). In summer, forests of all categories contained a significantly higher mean cumulative species' effect trait richness than for the other three seasons but most notably compared to winter. Broadleaf forests had significantly greater mean cumulative richness of species' effect traits that were associated with positive well-being compared to other forest categories in autumn, winter and spring (Fig. 4 and Supplementary Table 2). In summer, coniferous forests had the highest mean cumulative richness of species' effect traits. 'Other' forest categories had the lowest mean cumulative richness of species' effect traits in every season. The patterns for mean cumulative species' effect traits associated with negative well-being were consistent with those found for positive.

#### Spatio-temporal distributions of BIO-WELL scores

In total, 4,197 participants fully completed our online questionnaire, with different participants per season. Each seasonal cohort represented a diverse socio-demographic of the English and Welsh public in terms of gender, age, ethnicity and education (Supplementary Table 3), being distributed across England and Wales in both rural and urban areas (Fig. 5). Overall, participants experienced positive well-being in response to forest biodiversity, with BIO-WELL scores averaging 71.1 (range: 0.2–100 out of a possible 0–100), where values <50 indicate a negative response to biodiversity (12% of participants) and >50 indicate a positive response (88% of participants).

In general, we did not detect any differences in the BIO-WELL scores of participants between forest categories or across seasons (Fig. 5c and Supplementary Table 4). The one exception to this was summer, where participants reported statistically higher BIO-WELL scores associated with coniferous forest.

#### Effect traits, well-being and socio-economic deprivation

There was no association between forest area and level of deprivation ( $\beta = 0.027, 95\%$  CI = -0.996-1.051). However, mean cumulative species' effect trait richness, for both positive and negative well-being and across all four seasons, was greatest in the least socio-economic deprived areas where participants lived (Fig. 6a–d and Supplementary Table 5a). These patterns were consistent when examined for all forests across England and Wales (Extended Data Fig. 4 and Supplementary Table 6). Participant BIO-WELL scores were negatively associated with income-related deprivation in the winter and spring (Fig. 6e, f and Supplementary Table 5b).

#### Discussion

To understand how to create and manage ecosystems to meaningfully improve human health and well-being associated with biodiversity,



Fig. 2| Accumulative curves between species richness and species' effect trait richness associated with positive and negative well-being per season in forests across England and Wales. The curve displays the mean, with the upper and lower bounds of the shaded area representing 95% confidence intervals (mean value ± standard error).

researchers must move beyond coarse measures of 'nature' and 'greenspace' and account for the ecological communities present, which are inherently dynamic and spatially variable. In addition to better representing the complexity of ecosystems, we also need to recognize that human populations are diverse in their socio-economic composition and not distributed in a homogeneous manner. In this paper, we therefore begin the process of teasing apart this intricacy by examining spatio-temporal patterns between mean cumulative forest species' effect trait richness, well-being (both positive and negative) associations with biodiversity and socio-economic deprivation gradients. Where we cannot infer causality from our statistical analyses, our approach demonstrates how evidence derived from participatory processes and quantitative social science research methods<sup>17,18</sup> can initiate a step change in the development of forest restoration initiatives that seek to benefit both people and biodiversity.

We used two complementary ways to examine the potential for human well-being associated with forest biodiversity: species' effect traits and BIO-WELL. The former takes a functional ecology perspective<sup>17,36</sup> and the latter an environmental psychological standpoint<sup>18</sup>. We show that there are higher numbers of species' effect traits where species richness is greater, far more so for positive compared with negative well-being, across all seasons. Moreover, this pattern is more pronounced for autumn and winter for positive well-being, with the implication being that improvements in forest biodiversity would have a relatively larger impact on enhancing positive species' effect traits in these seasons (for example, refs. 37,38). When spatio-temporal distributions of mean cumulative species' effect traits were examined, summer supported the greatest richness of traits associated with positive well-being responses. Regional variation was observed, with the southeast having higher densities of traits. This reflects wider patterns of biodiversity across Britain, where the majority of species are on the northwest edge of their geographic range<sup>39</sup> and where broadleaf forest and ancient woodlands are concentrated<sup>40</sup>. With relatively few negative species' effect traits identified, little spatio-temporal variation was detected. Our findings affirm wider evidence suggesting that diverse forest ecosystems can help to positively enhance people's well-being<sup>8,9</sup>. This is despite the data requirements of the SDMs inevitably limiting the number of species that could be mapped to those that are part of national recording schemes with standardized survey methods, with adequate numbers of presence/absence records and that produced statistically acceptable models<sup>41,42</sup>. Furthermore, SDMs are not an actual representation of the biodiversity that is present in specific forests. Given that only 7% of Britain's forests are in good ecological condition (for example, presence of deadwood, veteran trees and diversity of ecological niches that support biodiversity<sup>40,43</sup>), it is unlikely that the majority are currently delivering their full human well-being potential. This adds further weight to calls for conservation and nature recovery to be at the heart of forest restoration initiatives<sup>44,45</sup>.

Further research is needed to ascertain the degree to which particular species' effect traits are more or less beneficial than others for well-being (for example, the smell of coniferous trees compared to the rough texture of bark), whether the relative 'strength' of each particular effect trait leads to different levels of well-being (for example, light to dark purple, potentially equating to within-species phenotypic variation) or if/how multiple effect traits interact to result in additive or multiplicative well-being responses (the cumulative effect of watching adult birds provisioning their chicks alongside the sound of birdsong from one or more species). Understanding these details could facilitate more targeted public health recommendations and interventions (for example, ref. 46) and ally research in this field with that with investigating how different levels of, and interactions between, multiple effect traits influence regulating and provisioning ecosystem service benefits<sup>21,22</sup>.

We found evidence of environmental health inequalities, with the more deprived sectors of society in England and Wales having less potential to gain positive well-being associated with forest biodiversity in proximity to where they live. For instance, disparities were apparent within southeast England between inland and coastal areas, the latter being typically suffering from more extreme levels of socio-economic deprivation<sup>47</sup>. Such spatial inequalities may be further exacerbated by the fact that the green spaces, where they do exist, are either not accessible to the public or are visited infrequently<sup>48</sup>. The lower use of green spaces can be attributed to a variety of factors that are social (for example, personal safety concerns), individual (for example, confidence in managing children outdoors) and contextual (for example, no free time)<sup>49</sup>. When we examined patterns in seasonality, people living in more deprived areas reported lower BIO-WELL scores in winter and spring. This trend could reflect the reduced species' effect trait richness apparent in winter but also less engagement with forests during the colder seasons of the year (for example, poor weather<sup>50</sup>).

The unequal distribution of forests rich in mean cumulative species' effect traits associated with positive well-being across deprivation gradients could also be explained by the 'luxury effect', predominately characterized as an urban phenomenon, which describes a positive association between higher biodiversity and affluence<sup>51</sup>. It is characterized by wealthier residents being drawn to more biodiverse and/or greener areas, creating a demand that raises property values and rents that effectively 'price out' individuals on lower incomes<sup>51</sup>. Another possible hypothesis could be that local authorities and/or private property owners in deprived areas invest less in forest conservation<sup>52</sup>. At local scales, however, more nuanced spatial patterns could be apparent that would require a finer-resolution analysis to disaggregate (for example, community forests in deprived areas may have relatively high biodiversity).

In areas of England and Wales characterized by higher deprivation, access to forests could be improved by strategically targeting nature recovery through better management of existing ecosystems and the creation of new ones. This is particularly pertinent, given that



Fig. 3 | Spatio-temporal distributions of cumulative forest species' effect trait richness per pixel across England and Wales for positive and negative well-being. Cumulative species' effect trait richness is the total number of unique effect trait-well-being incidences across all species, separated into positive and negative well-being, in autumn, winter, spring and summer. Base maps adapted from GADM.



Forest category

**Fig. 4** | **Mean cumulative species' effect trait richness per pixel for different categories of forest across England and Wales.** Forest categories (*n* = 566,394 forest polygons) follow NFI definitions; Supplementary Table 1. Violin plots display the probability density of mean cumulative species' effect trait richness (the width representing the frequency of data points) for each forest category, associated positive or negative well-being per season. White boxplots within the violin plots show the median, interquartile range, minimum and maximum of the same data. Kruskal Wallis H statistics are given in each panel, used to test for differences between forest categories per season, for positive and negative wellbeing, respectively (Supplementary Table 2 provides post-hoc Dunn-Bonferroni test results). Note: the y axis for negative well-being is a smaller scale.



#### Forest category

**Fig. 5** | **BIO-WELL scores, indicating well-being responses to forest biodiversity for online questionnaire participants across England and Wales. a,b**, Spatial distribution of BIO-WELL scores coloured by value (**a**) and a smooth histogram of BIO-WELL scores for each season (**b**). Dashed grey line represents the midpoint above or below which biodiversity is associated with positive (>50) or negative (<50) well-being responses, respectively. **c**, Violin plots displaying the probability density of BIO-WELL scores (the width representing the frequency of data points) for each forest category. Participants indicated the location of the nearby forest that their BIO-WELL score was relevant to in the questionnaire (n = 4,197). White boxplots within the violin plots show the median, interquartile range, minimum and maximum of the same data. BIO-WELL scores >50 and <50 indicate positive and negative well-being responses, respectively. Kruskal Wallis H statistics are given in each panel, used to test for differences between forest categories per season (Supplementary Table 4 provides post-hoc Dunn–Bonferroni test results). Base map in **a** adapted from GADM.

current restoration and tree-planting regimes that are intended to augment carbon sequestration generally overlook the heterogeneous biodiversity preferences, perspectives and values of people who may interact with the forests<sup>32,34,53,54</sup>. The success of forest creation/ restoration projects relies on recognizing this diversity, ensuring that such initiatives are equitable and socially just. In turn, this means that they are more likely to be supported by, and benefit, the local community<sup>55,56</sup>. Nevertheless, making management decisions within forests to promote species' effect traits that have the potential to enhance human well-being needs to be balanced alongside ecological considerations. For example, removing the species and effect traits associated with negative well-being could have detrimental consequences for ecosystem functioning and ecosystem service delivery (for example, degrading trophic interactions). Likewise, culturally important or charismatic species may not be of conservation interest or could be non-native<sup>57</sup>. Trade-offs may need to be navigated, taking care to ensure that unintended adverse impacts for biodiversity conservation are avoided.

In all seasons, we found that broadleaf forests had a greater richness of mean cumulative species' effect traits compared with other forest categories. In temperate climates, people visiting deciduous forests in autumn are more likely to encounter fruiting fungi or senescence of trees (for example, ref. 58), whereas those visiting in spring may experience forest-floor flowers<sup>40</sup>. Deciduous trees themselves support disproportionately high numbers of species' effect traits that elicit positive well-being responses<sup>17</sup>. This is because effect traits linked to the phenology and longevity of deciduous trees are embedded in people's everyday lives, for instance, large, old charismatic trees in Finland promoted sensory and emotional experiences<sup>59</sup>. Deciduous old-growth trees provide habitat for the highest diversity of animals, plants and fungi compared to other forest types<sup>40,60</sup>. Nonetheless, increasing the biodiversity of all forests has the potential to enhance human well-being and possibly delivering additional benefits across multiple other classes of ecosystem service<sup>22</sup>. BIO-WELL scores did not vary significantly between forest categories or seasons, other than being higher for coniferous forests in summer<sup>19</sup>. This might be a consequence of the uneven distribution of forest categories (81% broadleaf, 8% conifer, 3% mixed broadleaf, 1% mixed conifer, 7% other) in the analysis. On the other hand, coniferous forests in summer also supported the highest richness of species' effect traits. For example, at



**Fig. 6** | **Associations between mean cumulative species' effect trait richness and participant BIO-WELL scores per season across socio-economic deprivation gradients in England and Wales. a**-**f**, The *y* axes represent either the mean cumulative species' effect trait richness of all forests within each area where participants live in England and Wales, eliciting positive (**a**,**d**) or negative (**b**,**e**) well-being or BIO-WELL scores (**c**,**f**) (where >50 denotes positive wellbeing responses to forest biodiversity and <50 is negative). The *x* axes are the

proportion of the public considered to be experiencing income- or employmentrelated deprivation living in an area. Forest area (ha) is included as a covariate. Slopes indicate a general linear model with a 95% confidence interval (coloured shading) (Supplementary Table 5). Note: the *y* axes for negative well-being (**b**,**e**) are a smaller scale than for positive (**a**,**d**) and the *y* axis for BIO-WELL (**c**,**f**) is restricted.

this time of year, coniferous forests tend to be characterized by a pine scent that often carries cultural and personal importance and a unique canopy structure that can create well-defined and sun-lit footpaths<sup>61,62</sup>.

In this Article, we make new advances in biodiversity-well-being research by drawing upon and integrating functional ecology and environmental psychology concepts. Foremost, our granular approach can be operationalized by those planning where, when and for whom forest protection, restoration and creation should be targeted<sup>63</sup>. Likewise, it can inform the design and practice of social 'green' prescribing interventions<sup>14,15</sup>, particularly with the goal of improving well-being from forest biodiversity for certain sectors of society and to ensure that this is done adequately across the seasons. Indeed, moving forwards, a focus on how to improve people's well-being from biodiversity in the colder seasons could prove especially fruitful. Practitioners delivering forest restoration/creation or social prescribing interventions could then use BIO-WELL to monitor changes in human well-being in response to the ecological condition of forests<sup>18</sup>. Replicating our methodology for non-forest ecosystems will provide a more comprehensive picture of the well-being potential of biodiversity at a landscape scale. Areas with poor forest species' effect trait richness may contain other highly

valuable ecosystems (for example, coastal grasslands) that support different ecological communities with their own suite of effect traits. Understanding and accounting for this complexity could create more opportunities to deliver natural environments that underpin healthier individuals and societies.

#### Methods

#### Study system

Britain's forest ecosystems, which are 49% broadleaf and 51% coniferous, provide critical habitat for biodiversity<sup>40</sup>. Across England and Wales, 24.9% of the total land area is forest<sup>43</sup>. Forests are often publicly accessible and are among the most frequently visited types of ecosystem<sup>48</sup>. Here we use the NFI definition of forests (at least 0.5 ha in area, 20 m width and at least 20% canopy cover), using an open access dataset from the UK Government's Forest Research department (www.forestresearch.gov.uk) (Supplementary Table 1 and Extended Data Fig. 3).

#### Participatory workshops

We held four participatory workshops in 2019, to identify how people (n = 194) relate to forest biodiversity for their well-being<sup>17,18,20,53</sup>, during

each of the four seasons (autumn n = 48, winter n = 50, spring n = 46and summer n = 50). The participant cohort was new for each workshop. They represented a diversity of the public across age (18-29 years old n = 60, 30-59 n = 68, 60+n = 66), ethnicity (white British n = 146, other n = 48), gender (female n = 102, male n = 92), social grade (ABC1 n = 114, CDE2 n = 80) and urban-rural resident (urban n = 153, rural n = 41) (Extended Data Fig. 1). Social grade is defined as: AB (higher and intermediate managerial, administrative, professional occupations), C1 (supervisory, clerical and junior managerial, administrative and professional occupations), C2 (skilled manual occupations) and DE (semi-skilled and unskilled manual occupations, unemployed). All participants had to have been living in Britain for at least five years, irrespective of their nationality and were over 18 years old. Financial incentives (£100 per person per weekend) and upfront payment of expenses were used to support inclusive participation. Participants were recruited by a social research company to minimize the potential for self-selection bias (that is, individuals with a keen interest in nature).

We took participants to two forests (one being a mixed-deciduous and coniferous plantation, the other an ancient woodland), geographically located in the centre of Britain<sup>17,18,20</sup>. The forests were chosen to ensure that their objective physical and biological characteristics were diverse, both within and across the two ecosystems. We also made sure that the participants were not 'local' to either forest to minimize the impact that prior experience may have had on the their well-being responses to the objective biodiversity features of the sites. We ran a series of data collection activities designed to prompt discussion about forest biodiversity and what traits participants noticed (for example, smells, colours, textures, sounds, behaviours). These included an in situ scavenger hunt, ex situ focus groups and a series of ex situ image-based Q-methodology exercises (refs. 17,18,20 provide details). Activities were audio recorded and transcribed. Ethical approval was provided by the School of Anthropology and Conservation Research Ethics Committee, University of Kent (Ref: 009-ST-19). All participants provided informed consent before taking part in the research.

#### Seasonal species' effect traits

Workshop transcripts were analysed using NVivo (Version 12, QSR International Ply Ltd). We coded specific traits and how people's well-being responded to these traits, both positively and negatively, using the five domains of the biopsychosocial-spiritual model of health<sup>18</sup> (physical. emotional, cognitive, social, spiritual). For each trait, we then identified the species to which the participant was referring (for example, in the Q-methodology image or named by the participant) (ref. 17 provides details). Species that do not occur in British forests were excluded from the dataset. We made inferences about the species in cases were the participants mentioned specific phenological elements (for example 'acorns' were listed as English oak, Quercus robur). When participants alluded to traits associated with a taxonomic group of organisms (for example 'spots' on birds), we consulted reputable sources (Supplementary Table 7) to derive a list of species with that trait, excluding those that were too generic (for example, 'green' on plants). We then recorded the seasonal occurrence of species and their effect traits (for example, pied flycatchers, Ficedula hypoleuca, are not present in Britain in winter).

All data processing and statistical analyses were conducted in R (Version 4.2.0 (ref. 64)). To explore the relationship between species and effect traits in each season, we plotted accumulation curves of trait and species richness (function 'accumcomp' in package BiodiversityR<sup>65</sup>), for positive and negative well-being separately. Across species, there may be overlap in effect traits, meaning that there can be redundancy (where species delivering the same functions as others become functionally redundant/exchangeable) and complementarity (optimal combinations of species that deliver the maximum services) within ecological communities<sup>17</sup>. We gathered 2019 seasonal occurrence records for England and Wales for species within taxonomic groups (birds, butterflies, fungi, mammals and plants, including trees) that have national recording schemes with standardized survey methods (Supplementary Table 7). We used the UK Meteorological Office (www.metoffice.gov.uk/learning/seasons) definition of each season, which is based on the annual temperature cycle: autumn (1 September–31 November), winter (1 December–28/29 February), spring (1 March–30 May) and summer (1 June–31 August).

To generate the SDMs for each individual species, we selected a suite of biologically meaningful predictor variables (Supplementary Table 8), including elevation, precipitation and temperature data at 0.5 km resolution (30 arcseconds). These data are freely available from the BioClim dataset<sup>66</sup> (function 'getData', package Raster<sup>67</sup>). Elevation data were converted into topographic ruggedness (function 'tri', package spatialEco<sup>68</sup>). Topsoil data for land cover, dominant grain size and mean soil nitrogen concentration were acquired from the Countryside Survey<sup>69</sup>. Topsoil data were resampled using bilinear interpolation for continuous data to match the resolution of precipitation and temperature data. For each species, we tested the full set of environmental predictors for collinearity using a step-wise procedure, where highly correlated variables (VIF > 3) were removed.

We approximated seasonal distributions of individual species across English and Welsh forests using ensemble modelling, following best-practice techniques<sup>41,70-72</sup>. Given that recommendations for a minimum number of records for SDMs varies depending on whether species are common/rare and generalist/specialist<sup>41</sup>, we only retained species with a minimum of 80 survey records (for examples, ref. 41), leaving a total of 131 species (Supplementary Table 7). Survey records were uploaded using functions that minimize spatial autocorrelation while maximizing data availability ('load\_occ' function in the 'SSDM' package<sup>72</sup>). SDMs are sensitive to the type of algorithm applied to the data, so we fitted a suite of algorithms to derive statistical consensus among projections: classification tree analysis, generalized linear model and multivariate adaptive regression splines ('ensemble\_modelling' function in the 'SSDM' package<sup>72</sup>). We generated presence-only models for each species from the occurrence records and pseudo-absences (randomly selected artificial data about where each species cannot be found), using default parameters in the SSDM package<sup>72</sup>. Whereas SDMs can overestimate the distribution of planted species, they are routinely used to predict where non-native species may occur<sup>73</sup>. To boost predictive power while maintaining computational efficiency, we ran ten replicates per model algorithm per species<sup>74</sup> and required that models performed above a threshold value of >0.7 for area under the curve<sup>75</sup>. Model accuracy statistics were produced for all models and evaluated using the true skill statistic, assessing values > 0.4as fair, >0.5 as good, >0.7 as very good, >0.85 as excellent and >0.9 as perfect<sup>76</sup>. We created binary presence-absence maps using the highest true skill statistic threshold available for each species' set of models. Binary maps were subsequently clipped to the NFI shapefile for forests in England and Wales.

Our species' effect trait richness spatio-temporal distributions were constructed via the binary species maps (that is, the colour red was considered to be present wherever a European robin, *Erithacus rubec-ula*, was present). Traits were not treated as substitutable, given that the well-being benefits derived from traits are linked to the species and taxonomic group that the trait is supported by<sup>17</sup>. The spatio-temporal distributions of species' effect trait richness were captured in eight maps (positive and negative well-being separately for each of the four seasons) for which the data represents the overall cumulative effect trait richness within each pixel (0.5 min of a degree). We extracted the mean cumulative species' effect trait richness values across each NFI forest category using the 'extract' function in the package Raster<sup>67</sup>.

We compared mean cumulative species' effect trait richness between forest categories using a Kruskal Wallis H test and post-hoc Dunn–Bonferroni tests.

#### Online seasonal questionnaire

In 2021, we administered an online questionnaire to 4,710 participants across England and Wales using Qualtrics, across the four seasons. Participants could only complete the questionnaire once across all four seasons and were not sent the questionnaire if they had been present at one of the workshops. Once again, participants were recruited using a social research company to ensure there was no self-selection bias by individuals interested in nature or well-being and that a diverse public was represented (Supplementary Table 3). All participants were over 18 years old and had been resident in Britain for at least five years. As part of the questionnaire, we asked participants to provide the full postcode of where they lived. We also requested that they indicate a nearby forest on a map and that their questionnaire answers should relate to that forest. We removed data for 513 participants who did not locate a forest from subsequent analyses. We quantified the well-being people associate with forest biodiversity using BIO-WELL, a biodiversity-well-being psychometric scale<sup>18</sup> (https://research.kent. ac.uk/bio-well/). Participants were asked to record their well-being (physical, emotional, cognitive, spiritual and social) responses to different metrics and attributes of biodiversity (Supplementary Text provides details). For each participant, we calculated mean overall BIO-WELL scores across physical, cognitive, emotional, social and spiritual well-being.

#### Spatio-temporal distributions of BIO-WELL scores

We also examined whether there were differences in participants' BIO-WELL scores between the NFI categories associated with their nearby forests or across the seasons. When determining the NFI category for each participant's nearby forest, we used a 0.5-km buffer to account for potential resolution errors incurred through the coordinate system used in the online questionnaire. The differences were tested statistically using a Kruskal Wallis H and post-hoc Dunn–Bonferroni tests adjusted for multiple comparisons.

#### Effect traits, well-being and socio-economic deprivation

We used a government dataset to assess levels of human socio-economic deprivation<sup>77</sup>, at the smallest possible spatial resolution of Lower Super Output Areas (LSOAs; with an average of 1,700 people per LSOA<sup>78</sup>). Within each LSOA, we used the proportion of the population living in income- or employment-related deprivation (Extended Data Fig. 2). For each season, we then extracted the mean cumulative species' effect trait richness value of all forests within the LSOA, using the 'extract' function in the package Raster<sup>67</sup>. We related the mean cumulative species' effect trait richness of forests to the two measures of deprivation using a general linear model with a negative binomial error distribution. We included forest area (ha) as a covariate, given that larger areas are expected to contain a higher diversity of species<sup>79</sup>. Before this, we checked to ensure variance inflation factors were below 1.7 (ref. 80) and ran a bivariate linear model to investigate whether more deprived areas have smaller forests. This approach was repeated for participants' BIO-WELL scores and deprivation. All models were checked for fit, overdispersion and homoscedasticity<sup>81</sup>.

#### **Reporting summary**

Further information on research design is available in the Nature Portfolio Reporting Summary linked to this article.

## Data availability

The authors confirm that all questionnaire data generated during this study can be accessed at https://doi.org/10.22024/UniKent/01.01.541. All species and environmental data analysed for country-wide

modelling during this study are freely accessible as detailed in the Supplementary Material.

#### References

- 1. Díaz, S. et al. Pervasive human-driven decline of life on Earth points to the need for transformative change. *Science* **366**, eaax3100 (2019).
- Isbell, F. et al. Expert perspectives on global biodiversity loss and its drivers and impacts on people. *Front. Ecol. Environ.* 21, 94–103 (2023).
- Stiglitz, J. E., Sen, A. & Fitoussi, J.-P. Report by the Commission on the Measurement of Economic Performance and Social Progress (Commission on the Measurement of Economic Performance and Social Progress, 2010); https://ec.europa.eu/eurostat/ documents/8131721/8131772/Stiglitz-Sen-Fitoussi-Commissionreport.pdf
- 4. Constitution of the World Health Organisation (WHO, 1948); https://www.who.int/about/governance/constitution
- 5. Engel, G. The need for a new medical model. *Science* **196**, 129–136 (1977).
- Linton, M. J., Dieppe, P. & Medina-Lara, A. Review of 99 self-report measures for assessing well-being in adults: exploring dimensions of well-being and developments over time. *BMJ Open* 6, e010641 (2016).
- Irvine, K., Hoesly, D., Bell-Williams, R. & Warber, S. in *Biodiversity* and *Health in the Face of Climate Change* (eds Marselle, M. et al.) 213–247 (Springer Cham, 2019).
- Ferraro, D. M. et al. The phantom chorus: birdsong boosts human well-being in protected areas. *Proc. R. Soc. B* 287, 20201811 (2020).
- 9. Maes, M. J. A. et al. Benefit of woodland and other natural environments for adolescents' cognition and mental health. *Nat. Sustain.* **4**, 851–858 (2021).
- Maund, P. R. et al. Wetlands for wellbeing: piloting a nature-based health intervention for the management of anxiety and depression. *Int. J. Environ. Res. Public Health* 16, 4413 (2019).
- Steptoe, A. Happiness and health. Annu. Rev. Public Health 40, 339–359 (2019); https://doi.org/10.1146/annurev-publhealth-040218-044150
- 12. Valuing the Mental Health Benefits of Woodlands (Forest Research, 2021); https://www.forestresearch.gov.uk/publications/ valuing-the-mental-health-benefits-of-woodlands/
- Grellier, J. et al. Valuing the health benefits of nature-based recreational physical activity in England. *Environ. Int.* 187, 108667 (2024).
- Garside, R. et al. Therapeutic Nature: Nature-Based Social Prescribing for Diagnosed Mental Health Conditions in the UK (DEFRA, 2020); https://arc-swp.nihr.ac.uk/wp/wp-content/ uploads/2021/06/15138\_TherapeuticNature-Finalreport.pdf
- Shanahan, D. F. et al. Nature-based interventions for improving health and wellbeing: the purpose, the people and the outcomes. Sports 7, 141 (2019).
- Helbich, M., de Beurs, D., Kwan, M. P., O'Connor, R. C. & Groenewegen, P. P. Natural environments and suicide mortality in the Netherlands: a cross-sectional, ecological study. *Lancet Planet. Health* 2, e134–e139 (2018).
- 17. Fisher, J. C. et al. Human well-being responses to species' traits. Nat. Sustain. https://doi.org/10.1038/s41893-023-01151-3 (2023).
- Irvine, K. N. et al. BIO-WELL: the development and validation of a human wellbeing scale that measures responses to biodiversity. J. Environ. Psychol. 85, 101921 (2023).
- 19. Austen, G. E. et al. Exploring shared public perspectives on biodiversity attributes. *People Nat.* **3**, 901–913 (2021).
- 20. Fish, R. D. et al. Language matters for biodiversity. *BioScience* https://doi.org/10.1093/biosci/biae014 (2024).

### Article

- Lavorel, S. et al. A novel framework for linking functional diversity of plants with other trophic levels for the quantification of ecosystem services. J. Veg. Sci. 24, 942–948 (2013).
- 22. Felipe-Lucia, M. R. et al. Multiple forest attributes underpin the supply of multiple ecosystem services. *Nat. Commun.* https://doi.org/10.1038/s41467-018-07082-4 (2018).
- 23. Kwan, M. P. The stationarity bias in research on the environmental determinants of health. *Health Place* **70**, 102609 (2021).
- 24. Rowney, F. M. et al. Environmental DNA reveals links between abundance and composition of airborne grass pollen and respiratory health. *Curr. Biol.* **31**, 1995–2003.e4 (2021).
- 25. Graves, R. A., Pearson, S. M. & Turner, M. G. Effects of bird community dynamics on the seasonal distribution of cultural ecosystem services. *Ambio* **48**, 280–292 (2019).
- 26. Paraskevopoulou, A. T. et al. The impact of seasonal colour change in planting on patients with psychotic disorders using biosensors. *Urban For. Urban Green.* **36**, 50–56 (2018).
- 27. Young, M. T. et al. Quantifying urban park use in the USA at scale: empirical estimates of realised park usage using smartphone location data. *Lancet Planet. Health* **8**, e564–e573 (2024).
- Environmental Health Inequalities in Europe (WHO, 2019); https://iris.who.int/handle/10665/325176
- Mitchell, R. J., Richardson, E. A., Shortt, N. K. & Pearce, J. R. Neighborhood environments and socioeconomic inequalities in mental well-being. *Am. J. Prev. Med.* **49**, 80–84 (2015).
- Schüle, S. A., Hilz, L. K., Dreger, S. & Bolte, G. Social inequalities in environmental resources of green and blue spaces: a review of evidence in the WHO European region. *Int. J. Environ. Res. Public Health* 16, 1216 (2019).
- 31. Global Forest Resources Assessment 2015: How Are the World's Forests Changing? (FAO, 2016); https://openknowledge.fao.org/server/api/ core/bitstreams/29b8ae23-99f9-4a05-b796-9a35d02af29d/content
- 32. Augustynczik, A. L. D. et al. Socially optimal forest management and biodiversity conservation in temperate forests under climate change. *Ecol. Econ.* **169**, 106504 (2020).
- 33. The State of the World's Forests: Forests, Biodiversity, and People (FAO & UNEP, 2020); https://doi.org/10.4060/ca8642en
- Pritchard, R. Politics, power and planting trees. Nat. Sustain. 4, 932 (2021).
- Zhang, J., Fu, B., Stafford-Smith, M., Wang, S. & Zhao, W. Improve forest restoration initiatives to meet Sustainable Development Goal 15. Nat. Ecol. Evol. 5, 10–13 (2021).
- Richards, D. & Lavorel, S. Niche theory improves understanding of associations between ecosystem services. *One Earth* 6, 811–823 (2023).
- Song, C. et al. Physiological and psychological effects of walking on young males in urban parks in winter. *J. Physiol. Anthropol.* 32, 18 (2013).
- Song, C., Ikei, H., Igarashi, M., Takagaki, M. & Miyazaki, Y. Physiological and psychological effects of a walk in Urban parks in fall. Int. J. Environ. Res. Public Health 12, 14216–14228 (2015).
- Mason, S. C. et al. Geographical range margins of many taxonomic groups continue to shift polewards. *Biol. J. Linn. Soc.* 115, 586–597 (2015).
- 40. Reid, C. et al. State of the UK's Woods and Trees (Woodland Trust, 2021); https://www.woodlandtrust.org.uk/media/51705/ state-of-the-uks-woods-and-trees-2021-thewoodlandtrust.pdf
- Leroy, B. Choosing presence-only species distribution models. J. Biogeogr. https://doi.org/10.1111/jbi.14505 (2022).
- Soultan, A. & Safi, K. The interplay of various sources of noise on reliability of species distribution models hinges on ecological specialisation. *PLoS ONE* **12**, e0187906 (2017).
- 43. Forestry Statistics. Chapter 1: Woodland Area and Planting (Forest Research, 2020); https://cdn.forestresearch.gov.uk/2022/02/ ch1\_woodland\_fs2020\_cgadfu3.pdf

- 44. Bradbury, R. B. et al. The economic consequences of conserving or restoring sites for nature. *Nat. Sustain.* **4**, 602–608 (2021).
- 45. Cook-Patton, S. C. et al. Protect, manage and then restore lands for climate mitigation. *Nat. Clim. Chang.* **11**, 1027–1034 (2021).
- White, M. P. et al. Spending at least 120 minutes a week in nature is associated with good health and wellbeing. *Sci. Rep.* 9, 7730 (2019).
- 47. Chief Medical Officer Health in Coastal Communities (UK Government Department of Health and Social Care, 2021); https://www.gov.uk/government/publications/chief-medicalofficers-annual-report-2021-health-in-coastal-communities
- Monitor of Engagement with the Natural Environment: The National Survey on People and the Natural Environment—Headline Report 2019 (Natural England, 2019); https://assets.publishing. service.gov.uk/media/5d6cd601e5274a170c435365/Monitor\_ Engagement\_Natural\_Environment\_2018\_2019\_v2.pdf
- 49. Cronin-de-chavez, A., Islam, S. & McEachan, R. R. C. Not a level playing field: a qualitative study exploring structural, community and individual determinants of greenspace use amongst low-income multi-ethnic families. *Health Place* **56**, 118–126 (2019).
- 50. Boyd, F., White, M. P., Bell, S. L. & Burt, J. Who doesn't visit natural environments for recreation and why: a population representative analysis of spatial, individual and temporal factors among adults in England. *Landsc. Urban Plan.* **175**, 102–113 (2018).
- Leong, M., Dunn, R. R. & Trautwein, M. D. Biodiversity and socioeconomics in the city: a review of the luxury effect. *Biol. Lett.* 14, 20180082 (2018).
- 52. Lawrence, A. & Dandy, N. Private landowners' approaches to planting and managing forests in the UK: What's the evidence? *Land Use Policy* **36**, 351–360 (2014).
- 53. Austen, G. E. et al. The diversity of people's relationships with biodiversity should inform forest restoration and creation. *Conserv. Lett.* https://doi.org/10.1111/conl.12930 (2022).
- 54. O'Brien, L. et al. Exploring the social and cultural values of trees and woodlands in England: a new composite measure. *People Nat.* **6**, 1334–1354 (2024).
- Coleman, E. A. et al. Limited effects of tree planting on forest canopy cover and rural livelihoods in Northern India. *Nat. Sustain.* 4, 997–1004 (2021).
- 56. Erbaugh, J. T. et al. Global forest restoration and the importance of prioritizing local communities. *Nat. Ecol. Evol.* **4**, 1472–1476 (2020).
- 57. Langlois, J. et al. The aesthetic value of reef fishes is globally mismatched to their conservation priorities. *PLoS Biol.* **20**, e3001640 (2022).
- Andrew, C. et al. Explaining European fungal fruiting phenology with climate variability. *Ecology* 99, 1306–1315 (2018).
- 59. Vainio, K. et al. Do you have a tree friend? Human-tree relationships in Finland. *People Nat.* **6**, 646–659 (2024).
- 60. Gibson, L. et al. Primary forests are irreplaceable for sustaining tropical biodiversity. *Nature* **478**, 378–381 (2011).
- 61. Bentley, P. R. et al. Nature, smells, and human wellbeing. *Ambio* https://doi.org/10.1007/s13280-022-01760-w (2022).
- 62. Hickman, C. Pine fresh: the cultural and medical context of pine scent in relation to health from the forest to the home. *Med. Humanit.* **48**, 104–113 (2022).
- 63. The State of the World's Forests 2022. Forest Pathways for Green Recovery and Building Inclusive, Resilient and Sustainable Economies (FAO, 2022); https://doi.org/10.4060/cb9360en (2022).
- 64. R Core Team R: A Language and Environment for Statistical Computing (R Foundation for Statistical Computing, 2022).
- 65. Kindt, R. & Coe, R. Tree Diversity Analysis. A Manual and Software for Common Statistical Methods for Ecological and Biodiversity Studies (ICRAF, 2005); http://www.worldagroforestry.org/output/ tree-diversity-analysis

## Article

- Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G. & Jarvis, A. Very high resolution interpolated climate surfaces for global land areas. *Int. J. Climatol.* 25, 1965–1978 (2005).
- 67. Hijmans, R. J. raster: geographic data analysis and modeling. R package version 3.6-20 (2022).
- Riley, S. J., DeGloria, S. D. & Elliot, R. A terrain ruggedness index that quantifies topographic heterogeneity. *Intermt. J. Sci.* 5, 23–27 (1999).
- Emmett, B. A. et al. Countryside Survey: Soils Report from 2007 (Centre for Ecology and Hydrology, 2010); https://nora.nerc. ac.uk/id/eprint/9354/1/CS\_UK\_2007\_TR9.pdf
- Araújo, M. B. et al. Standards for distribution models in biodiversity assessments. Sci. Adv. 5, eaat4858 (2019).
- Booth, T. H. Checking bioclimatic variables that combine temperature and precipitation data before their use in species distribution models. *Austral Ecol.* https://doi.org/10.1111/aec.13234 (2022).
- Schmitt, S., Pouteau, R., Justeau, D., de Boissieu, F. & Birnbaum, P. ssdm: an R package to predict distribution of species richness and composition based on stacked species distribution models. *Methods Ecol. Evol.* 8, 1795–1803 (2017).
- 73. Jarnevich, C. et al. Invaders at the doorstep: using species distribution modeling to enhance invasive plant watch lists. *Ecol. Inform.* **75**, 101997 (2023).
- Baumbach, L., Warren, D. L., Yousefpour, R. & Hanewinkel, M. Climate change may induce connectivity loss and mountaintop extinction in Central American forests. *Commun. Biol.* 4, 869 (2021).
- Hosmer, D. W., Lemeshow, S. & Sturdivant, R. X. Applied Logistic Regression (John Wiley & Sons, 2013); https://doi.org/10.1002/ 9781118548387
- 76. Landis, J. R. & Koch, G. G. The measurement of observer agreement for categorical data. *Biometrics* **33**, 159–174 (1977).
- 77. Indices of Deprivation 2019: Income and Employment Domains Combined for England and Wales (Ministry of Housing, Communities, & Local Government, 2020); https://www.gov.uk/ government/statistics/indices-of-deprivation-2019-incomeand-employment-domains-combined-for-england-and-wales
- 78. Lower Layer Super Output Area (LSOA) Boundaries (ONS, 2016); https://www.data.gov.uk/dataset/fa883558-22fb-4a1a-8529cffdee47d500/lower-layer-super-output-area-lsoa-boundaries
- Kunin, W. E. et al. Upscaling biodiversity: estimating the speciesarea relationship from small samples. *Ecol. Monogr.* 88, 170–187 (2018).
- Zuur, A. F. & Ieno, E. N. A protocol for conducting and presenting results of regression-type analyses. *Methods Ecol. Evol.* 7, 636–645 (2016).
- Harrison, X. A. et al. A brief introduction to mixed effects modelling and multi-model inference in ecology. *PeerJ* 6, e4794 (2018).

# Acknowledgements

We would like to thank all the participants who took park in both the workshops and questionnaires. We are grateful to N.J. Deere, J.E. Bicknell and L. Santiago for useful methodological discussions. The analyses were powered by the specialist and high-performance computing systems provided by Information Services at the University of Kent. Additional thanks go to the organizations who provided species presence/absence data: Mammal Society, British Trust for Ornithology (BTO), Joint Nature Conservation Committee (JNCC), Royal Society for the Protection of Birds (RSPB), UK Butterfly Monitoring Scheme (UKBMS), Fungal Records Database of Britain and Ireland (FRDBI), The Mammal Society and Botanical Society of Britain and Ireland (BSBI). This research was funded by the European Research Council (ERC) Horizon 2020 Research and Innovation Programme (consolidator grant number 726104, held by Z.G.D.) and Woodland Trust (grant number GC22SRG076, held by Z.G.D.). Additionally, J.C.F. and Z.G.D. were supported by Research England's 'Expanding Excellence in England' fund.

# **Author contributions**

Z.G.D. and M.D. conceptualized the study. Z.G.D., M.D., K.N.I. and R.D.F. acquired the funding. Z.G.D. and G.E.A. were responsible for project administration. J.C.F., M.D. and Z.G.D. developed the methodology. J.C.F., S.G.A. and P.M.K. curated and interpreted the data. J.C.F. was responsible for the software and undertook the visualization, formal analysis and writing of the original draft paper. All authors contributed to the investigation and reviewing/editing of the paper.

# **Competing interests**

The authors declare no competing interests.

# **Additional information**

**Extended data** is available for this paper at https://doi.org/10.1038/s41559-025-02765-w.

**Supplementary information** The online version contains supplementary material available at https://doi.org/10.1038/s41559-025-02765-w.

**Correspondence and requests for materials** should be addressed to J. C. Fisher.

**Peer review information** *Nature Ecology & Evolution* thanks Triin Reitalu and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

**Reprints and permissions information** is available at www.nature.com/reprints.

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons. org/licenses/by/4.0/.

© The Author(s) 2025



**Extended Data Fig. 1** | **Residential location of workshop participants across England and Wales.** Blue shading represents the number of participants sampled from each region. Base map adapted from GADM.



**Extended Data Fig. 2** | **Spatial distribution of income- and employment-related deprivation in England and Wales.** The gradient shows the proportion of the public considered to be experiencing (**a**) income- or (**b**) employment-related deprivation living in an area. Base maps adapted from GADM.



**Extended Data Fig. 3** | **Distribution of National Forest Inventory forest categories across England and Wales.** Colours represent forest categories (green = broadleaf, 60%, blue = conifer, 14%, light green = mixed mainly broadleaved, 5%, light blue = mixed mainly conifer, 5%, dark grey = other, 17%). Base map adapted from GADM.

9



Extended Data Fig. 4 | Associations between mean cumulative species' effect trait richness, per season, across socioeconomic deprivation gradients in England and Wales. The y axes represent the mean cumulative species' effect trait richness of all forests across England and Wales, eliciting positive (**a**, **c**) or negative (**b**, **d**) wellbeing. The x axes are the proportion of the public considered to be experiencing income- or employment-related deprivation living in an area. Forest area (hectares) is included as a covariate. Slopes indicate a general linear model with a 95% confidence interval (coloured shading) (Supplementary Table 6). NB: the y axis for negative wellbeing (**b**, **d**) is a smaller scale.

# nature portfolio

Corresponding author(s): Fisher JC

Last updated by author(s): Apr 22, 2025

# **Reporting Summary**

Nature Portfolio wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Portfolio policies, see our <u>Editorial Policies</u> and the <u>Editorial Policy Checklist</u>.

# **Statistics**

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.				
n/a	Cor	firmed		
	$\boxtimes$	The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement		
	$\boxtimes$	A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly		
	$\boxtimes$	The statistical test(s) used AND whether they are one- or two-sided Only common tests should be described solely by name; describe more complex techniques in the Methods section.		
	$\boxtimes$	A description of all covariates tested		
	$\boxtimes$	A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons		
	$\boxtimes$	A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)		
		For null hypothesis testing, the test statistic (e.g. <i>F</i> , <i>t</i> , <i>r</i> ) with confidence intervals, effect sizes, degrees of freedom and <i>P</i> value noted Give <i>P</i> values as exact values whenever suitable.		
$\boxtimes$		For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings		
$\boxtimes$		For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes		
	$\boxtimes$	Estimates of effect sizes (e.g. Cohen's d, Pearson's r), indicating how they were calculated		
		Our web collection on <u>statistics for biologists</u> contains articles on many of the points above.		

# Software and code

Policy information about <u>availability of computer code</u>			
Data collection	Questionnaire data collection was conducted using Qualtrics.		
Data analysis	Qualitative analyses were conducted in NVIVO version 12 (QSR International). Statistical analyses were conducted in R version 4.3.0 (https://www.r-project.org/)		

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Portfolio guidelines for submitting code & software for further information.

# Data

Policy information about availability of data

All manuscripts must include a data availability statement. This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A description of any restrictions on data availability
- For clinical datasets or third party data, please ensure that the statement adheres to our policy

The authors confirm that all questionnaire data generated during this study can be accessed from the following repository: https://doi.org/10.22024/ UniKent/01.01.541. All species and environmental data analysed for country-wide modelling during this study are freely accessible as detailed in the Supplementary Material.

# Human research participants

Policy information about studies involving human research participants and Sex and Gender in Research.

Reporting on sex and gender	Data were collected for gender (shaped by social and cultural circumstances), as determined by participants' self-reporting. Individuals were selected to ensure diversity of perspectives from the public across gender (male = 90, female = 103, prefer not to say = 1), but testing for differences between gender per se was not within the scope of this study. Participants provided written informed consent prior to data collection.
Population characteristics	See above
Recruitment	Workshop participants (n = 194) were recruited via a social research agency between February and October 2019. Individuals were selected to ensure diversity of perspectives from the public. To encourage workshop attendance and inclusivity, participants were incentivised by travel reimbursement and financial renumeration (£100 per person per weekend). Questionnaire participants (n = 4710) were recruited via the same social research agency, between February and October 2021. All participants were over 18 years old and had been resident in Britain for at least five years, and provided informed consent prior to taking part in the research.
Ethics oversight	Ethics approval was provided by the School of Anthropology and Conservation Research Ethics Committee, University of Kent (Ref: 009-ST-19).

Note that full information on the approval of the study protocol must also be provided in the manuscript.

# Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

└── Life sciences └── Behavioural & social sciences └── Ecological, evolutionary & environmental scie
---

For a reference copy of the document with all sections, see <u>nature.com/documents/nr-reporting-summary-flat.pdf</u>

# Life sciences study design

All studies must disclose on these points even when the disclosure is negative.

Sample size	Describe how sample size was determined, detailing any statistical methods used to predetermine sample size OR if no sample-size calculation was performed, describe how sample sizes were chosen and provide a rationale for why these sample sizes are sufficient.
Data exclusions	Describe any data exclusions. If no data were excluded from the analyses, state so OR if data were excluded, describe the exclusions and the rationale behind them, indicating whether exclusion criteria were pre-established.
Replication	Describe the measures taken to verify the reproducibility of the experimental findings. If all attempts at replication were successful, confirm this OR if there are any findings that were not replicated or cannot be reproduced, note this and describe why.
Randomization	Describe how samples/organisms/participants were allocated into experimental groups. If allocation was not random, describe how covariates were controlled OR if this is not relevant to your study, explain why.
Blinding	Describe whether the investigators were blinded to group allocation during data collection and/or analysis. If blinding was not possible, describe why OR explain why blinding was not relevant to your study.

# Behavioural & social sciences study design

All studies must disclose on these points even when the disclosure is negative.

Study description	The study type is mixed methods (qualitative and quantitative cross-sectional, and ecological analyses)
Research sample	Workshop participants were selected to ensure diversity of perspectives from the British public across gender (male = 90, female = 103, prefer not to say = 1), ethnicity (white British = 131, other = 63), and age (18-29 years = 59, 30-59 years = 70, 60+ years = 59, prefer not to say = 6), social grade (AB = 56, C1 = 58, C2 = 42, DE = 38). This dataset oversampled on unrepresented groups (e.g. non-white British). Likewise, questionnaire participants represented a diversity of the British public, oversampled on unrepresented groups. These included gender (male = 2220, female = 1977), ethnicity (white British = 3207, other = 990), and age (range = 18 - 93).
Sampling strategy	For both the workshops and questionnaires, the sampling procedure was stratified, based on simple quotas provided to the social research agency. No sample-size calculation was performed as the dataset was originally qualitative, but the sample sizes (n = 194; n = 4197) were deemed sufficient to represent a diverse set of responses from across the British public.

Data collection	During workshops, participants took part in a 1-hour scavenger hunt in-situ and given paper, pen, and a clipboard. Following these visits, participants were divided into focus groups to discuss their impressions of the forest, recorded using a Dictaphone device. On the second day participants undertook multiple image-based Q-methodology activities (see Austen et al. 2021) using paper and pens. Facilitators were present during the data collection, but were not blind to the study hypotheses. For questionnaires, participants completed questions about where they lived, and their wellbeing responses to attributes of forest biodiversity (BIO-WELL scale, Irvine et al. 2023).	
Timing	During workshops, participants were split over four weekends (n = 46-50 per workshop) across the year (winter = February, Spring = May, Summer = July, Autumn = October). For questionnaires, delivering took place during these same months (winter = 1115, spring = 1021, summer = 1041, autumn = 1020).	
Data exclusions	No participants were excluded from the analyses.	
Non-participation	For workshops, 6 participants were unable to attend due to personal circumstances or weather inhibiting the ability to travel. For questionnaires, 513 participants did not locate a forest when considering the BIO-WELL scale.	
Randomization	Participants were randomly allocated into focus groups.	

# Ecological, evolutionary & environmental sciences study design

All studies must disclose on these points even when the disclosure is negative.

Study description	Briefly describe the study. For quantitative data include treatment factors and interactions, design structure (e.g. factorial, nested, hierarchical), nature and number of experimental units and replicates.
Research sample	Describe the research sample (e.g. a group of tagged Passer domesticus, all Stenocereus thurberi within Organ Pipe Cactus National Monument), and provide a rationale for the sample choice. When relevant, describe the organism taxa, source, sex, age range and any manipulations. State what population the sample is meant to represent when applicable. For studies involving existing datasets, describe the data and its source.
Sampling strategy	Note the sampling procedure. Describe the statistical methods that were used to predetermine sample size OR if no sample-size calculation was performed, describe how sample sizes were chosen and provide a rationale for why these sample sizes are sufficient.
Data collection	Describe the data collection procedure, including who recorded the data and how.
Timing and spatial scale	Indicate the start and stop dates of data collection, noting the frequency and periodicity of sampling and providing a rationale for these choices. If there is a gap between collection periods, state the dates for each sample cohort. Specify the spatial scale from which the data are taken
Data exclusions	If no data were excluded from the analyses, state so OR if data were excluded, describe the exclusions and the rationale behind them, indicating whether exclusion criteria were pre-established.
Reproducibility	Describe the measures taken to verify the reproducibility of experimental findings. For each experiment, note whether any attempts to repeat the experiment failed OR state that all attempts to repeat the experiment were successful.
Randomization	Describe how samples/organisms/participants were allocated into groups. If allocation was not random, describe how covariates were controlled. If this is not relevant to your study, explain why.
Blinding	Describe the extent of blinding used during data acquisition and analysis. If blinding was not possible, describe why OR explain why blinding was not relevant to your study.
Did the study involve field	d work? Yes No

# Field work, collection and transport

Field conditions	Describe the study conditions for field work, providing relevant parameters (e.g. temperature, rainfall).
Location	State the location of the sampling or experiment, providing relevant parameters (e.g. latitude and longitude, elevation, water depth).
Access & import/export	Describe the efforts you have made to access habitats and to collect and import/export your samples in a responsible manner and in compliance with local, national and international laws, noting any permits that were obtained (give the name of the issuing authority, the date of issue, and any identifying information).
Disturbance	Describe any disturbance caused by the study and how it was minimized.

# Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

Materials & experimental systems		Methods	
n/a	Involved in the study	n/a	Involved in the study
$\boxtimes$	Antibodies	$\boxtimes$	ChIP-seq
$\boxtimes$	Eukaryotic cell lines	$\boxtimes$	Flow cytometry
$\boxtimes$	Palaeontology and archaeology	$\boxtimes$	MRI-based neuroimaging
$\boxtimes$	Animals and other organisms		
$\boxtimes$	Clinical data		
$\boxtimes$	Dual use research of concern		

# Antibodies

Antibodies used	Describe all antibodies used in the study; as applicable, provide supplier name, catalog number, clone name, and lot number.
Validation	Describe the validation of each primary antibody for the species and application, noting any validation statements on the manufacturer's website, relevant citations, antibody profiles in online databases, or data provided in the manuscript.

# Eukaryotic cell lines

#### Policy information about cell lines and Sex and Gender in Research

Cell line source(s)	State the source of each cell line used and the sex of all primary cell lines and cells derived from human participants or vertebrate models.
Authentication	Describe the authentication procedures for each cell line used OR declare that none of the cell lines used were authenticated.
Mycoplasma contamination	Confirm that all cell lines tested negative for mycoplasma contamination OR describe the results of the testing for mycoplasma contamination OR declare that the cell lines were not tested for mycoplasma contamination.
Commonly misidentified lines (See <u>ICLAC</u> register)	Name any commonly misidentified cell lines used in the study and provide a rationale for their use.

# Palaeontology and Archaeology

Specimen provenance	Provide provenance information for specimens and describe permits that were obtained for the work (including the name of the issuing authority, the date of issue, and any identifying information). Permits should encompass collection and, where applicable, export.
Specimen deposition	Indicate where the specimens have been deposited to permit free access by other researchers.
Dating methods	If new dates are provided, describe how they were obtained (e.g. collection, storage, sample pretreatment and measurement), where they were obtained (i.e. lab name), the calibration program and the protocol for quality assurance OR state that no new dates are provided.
Tick this box to confi	rm that the raw and calibrated dates are available in the paper or in Supplementary Information.
Ethics oversight	Identify the organization(s) that approved or provided guidance on the study protocol, OR state that no ethical approval or guidance was required and explain why not

Note that full information on the approval of the study protocol must also be provided in the manuscript.

# Animals and other research organisms

Policy information about studies involving animals; ARRIVE guidelines recommended for reporting animal research, and Sex and Gender in Research

Laboratory animals (For laboratory animals, report species, strain and age OR state that the study did not involve laboratory animals.

Wild animals	Provide details on animals observed in or captured in the field; report species and age where possible. Describe how animals were caught and transported and what happened to captive animals after the study (if killed, explain why and describe method; if released, say where and when) OR state that the study did not involve wild animals.
Reporting on sex	Indicate if findings apply to only one sex; describe whether sex was considered in study design, methods used for assigning sex. Provide data disaggregated for sex where this information has been collected in the source data as appropriate; provide overall numbers in this Reporting Summary. Please state if this information has not been collected. Report sex-based analyses where performed, justify reasons for lack of sex-based analysis.
Field-collected samples	For laboratory work with field-collected samples, describe all relevant parameters such as housing, maintenance, temperature, photoperiod and end-of-experiment protocol OR state that the study did not involve samples collected from the field.
Ethics oversight	Identify the organization(s) that approved or provided guidance on the study protocol, OR state that no ethical approval or guidance was required and explain why not.

Note that full information on the approval of the study protocol must also be provided in the manuscript.

# Clinical data

Policy information about clinical studies

All manuscripts should comply with the ICMJE guidelines for publication of clinical research and a completed CONSORT checklist must be included with all submissions.

Clinical trial registration	Provide the trial registration number from ClinicalTrials.gov or an equivalent agency.
Study protocol	Note where the full trial protocol can be accessed OR if not available, explain why.
Data collection	Describe the settings and locales of data collection, noting the time periods of recruitment and data collection.
Outcomes	Describe how you pre-defined primary and secondary outcome measures and how you assessed these measures.

# Dual use research of concern

Policy information about dual use research of concern

#### Hazards

Could the accidental, deliberate or reckless misuse of agents or technologies generated in the work, or the application of information presented in the manuscript, pose a threat to:

 No
 Yes

 Image: Constraint of the security
 Image: Constraint of the security

 Image: Constraint of the security
 Image: Constraint of the security

 Image: Constraint of the security
 Image: Constraint of the security

 Image: Constraint of the security
 Image: Constraint of the security

 Image: Constraint of the security
 Image: Constraint of the security

 Image: Constraint of the security
 Image: Constraint of the security

 Image: Constraint of the security
 Image: Constraint of the security

## Experiments of concern

Does the work involve any of these experiments of concern:

- No Yes
- Demonstrate how to render a vaccine ineffective
   Confer resistance to therapeutically useful antibiotics or antiviral agents
   Enhance the virulence of a pathogen or render a nonpathogen virulent
   Increase transmissibility of a pathogen
   Alter the host range of a pathogen
   Enable evasion of diagnostic/detection modalities
   Enable the weaponization of a biological agent or toxin
- Any other potentially harmful combination of experiments and agents

# ChIP-seq

#### Data deposition

Confirm that both raw and final processed data have been deposited in a public database such as GEO.

Confirm that you have deposited or provided access to graph files (e.g. BED files) for the called peaks.

Data access links May remain private before publication.	For "Initial submission" or "Revised version" documents, provide reviewer access links. For your "Final submission" document, provide a link to the deposited data.
Files in database submission	Provide a list of all files available in the database submission.
Genome browser session (e.g. <u>UCSC</u> )	Provide a link to an anonymized genome browser session for "Initial submission" and "Revised version" documents only, to enable peer review. Write "no longer applicable" for "Final submission" documents.

## Methodology

Replicates	Describe the experimental replicates, specifying number, type and replicate agreement.
Sequencing depth	Describe the sequencing depth for each experiment, providing the total number of reads, uniquely mapped reads, length of reads and whether they were paired- or single-end.
Antibodies	Describe the antibodies used for the ChIP-seq experiments; as applicable, provide supplier name, catalog number, clone name, and lot number.
Peak calling parameters	Specify the command line program and parameters used for read mapping and peak calling, including the ChIP, control and index files used.
Data quality	Describe the methods used to ensure data quality in full detail, including how many peaks are at FDR 5% and above 5-fold enrichment.
Software	Describe the software used to collect and analyze the ChIP-seq data. For custom code that has been deposited into a community repository, provide accession details.

# Flow Cytometry

#### Plots

Confirm that:

The axis labels state the marker and fluorochrome used (e.g. CD4-FITC).

The axis scales are clearly visible. Include numbers along axes only for bottom left plot of group (a 'group' is an analysis of identical markers).

All plots are contour plots with outliers or pseudocolor plots.

A numerical value for number of cells or percentage (with statistics) is provided.

## Methodology

Sample preparation	Describe the sample preparation, detailing the biological source of the cells and any tissue processing steps used.
Instrument	Identify the instrument used for data collection, specifying make and model number.
Software	Describe the software used to collect and analyze the flow cytometry data. For custom code that has been deposited into a community repository, provide accession details.
Cell population abundance	Describe the abundance of the relevant cell populations within post-sort fractions, providing details on the purity of the samples and how it was determined.
Gating strategy	Describe the gating strategy used for all relevant experiments, specifying the preliminary FSC/SSC gates of the starting cell population, indicating where boundaries between "positive" and "negative" staining cell populations are defined.

Tick this box to confirm that a figure exemplifying the gating strategy is provided in the Supplementary Information.

# Magnetic resonance imaging

#### Experimental design

Design type

Indicate task or resting state; event-related or block design.

Design specifications	Specify the number of blocks, trials or experimental units per session and/or subject, and specify the length of each trial or block (if trials are blocked) and interval between trials.	
Behavioral performance measure	State number and/or type of variables recorded (e.g. correct button press, response time) and what statistics were used to establish that the subjects were performing the task as expected (e.g. mean, range, and/or standard deviation across subjects).	
Acquisition		
Imaging type(s)	Specify: functional, structural, diffusion, perfusion.	
Field strength	Specify in Tesla	
Sequence & imaging parameters	Specify the pulse sequence type (gradient echo, spin echo, etc.), imaging type (EPI, spiral, etc.), field of view, matrix size, slice thickness, orientation and TE/TR/flip angle.	
Area of acquisition	State whether a whole brain scan was used OR define the area of acquisition, describing how the region was determined	
Diffusion MRI 📃 Used	Not used	
Preprocessing		
Preprocessing software	Provide detail on software version and revision number and on specific parameters (model/functions, brain extraction, segmentation, smoothing kernel size, etc.).	
Normalization	If data were normalized/standardized, describe the approach(es): specify linear or non-linear and define image types used for transformation OR indicate that data were not normalized and explain rationale for lack of normalization.	
Normalization template	Describe the template used for normalization/transformation, specifying subject space or group standardized space (e.g. original Talairach, MNI305, ICBM152) OR indicate that the data were not normalized.	
Noise and artifact removal	Describe your procedure(s) for artifact and structured noise removal, specifying motion parameters, tissue signals and physiological signals (heart rate, respiration).	
Volume censoring	Define your software and/or method and criteria for volume censoring, and state the extent of such censoring.	

# Statistical modeling & inference

Model type and settings	Specify type (mass univariate, multivariate, RSA, predictive, etc.) and describe essential details of the model at the first and second levels (e.g. fixed, random or mixed effects; drift or auto-correlation).	
Effect(s) tested	Define precise effect in terms of the task or stimulus conditions instead of psychological concepts and indicate whether ANOVA or factorial designs were used.	
Specify type of analysis: 🗌 Whole brain 📄 ROI-based 📄 Both		
Statistic type for inference (See <u>Eklund et al. 2016</u> )	Specify voxel-wise or cluster-wise and report all relevant parameters for cluster-wise methods.	
Correction	Describe the type of correction and how it is obtained for multiple comparisons (e.g. FWE, FDR, permutation or Monte Carlo).	

# Models & analysis

n/a       Involved in the study         Image: State of the study       Functional and/or effective connectivity         Image: State of the study       Graph analysis         Image: State of the study       Multivariate modeling or predictive analysis		
Functional and/or effective connectivity	Report the measures of dependence used and the model details (e.g. Pearson correlation, partial correlation, mutual information).	
Graph analysis	Report the dependent variable and connectivity measure, specifying weighted graph or binarized graph, subject- or group-level, and the global and/or node summaries used (e.g. clustering coefficient, efficiency, etc.).	
Multivariate modeling and predictive analysis	Specify independent variables, features extraction and dimension reduction, model, training and evaluation metrics.	