



Disease surveillance in albatrosses and petrels from the Southwest Atlantic and Southern Ocean

Research Article

We would like very much that it is made explicit in the paper that Patricia Pereira Serafini and Annelise Zabel Sgarioni share the first co-authorship and that these authors contributed equally to this work.

Cite this article: Pereira Serafini P *et al.* (2025) Disease surveillance in albatrosses and petrels from the Southwest Atlantic and Southern Ocean. *Parasitology* **152**, 1172–1178. <https://doi.org/10.1017/S0031182025100590>

Received: 5 April 2025

Revised: 10 June 2025

Accepted: 12 July 2025

First published online: 11 August 2025

Keywords:

climate-driven risks; disease surveillance; haemosporidian parasites; seabirds health; vector-borne pathogens

Corresponding author:

Patricia Pereira Serafini;

Email: patricia.serafini@icmbio.gov.br

Patricia Pereira Serafini^{1,2} , Annelise Zabel Sgarioni³, Richard A. Phillips⁴, Alice Pereira⁵, Tiffany Emmerich⁶, Thamires P. Pontes⁶, Derek B. Amorim⁷, Cristiane K. M. Kolesnikovas⁸, André O. S. Lima³, Guilherme Klafke⁹, José Reck⁹, Afonso C. D. Bainy¹, Karim H. Lüchmann¹⁰ and Camille Bonneaud¹¹

¹Laboratório de Biomarcadores de Contaminação Aquática e Imunoquímica, Universidade Federal de Santa Catarina – UFSC, Florianópolis, SC, Brazil; ²Centro Nacional de Pesquisa e Conservação de Aves Silvestres, Instituto Chico Mendes de Conservação da Biodiversidade – ICMBio, Florianópolis, SC, Brazil; ³Laboratório de Genética Molecular, Escola Politécnica, Universidade do Vale do Itajaí (UNIVALI), Itajaí, SC, Brazil; ⁴British Antarctic Survey, Natural Environment Research Council, Cambridge, Cambridgeshire, UK; ⁵Projeto Albatroz, Florianópolis, SC, Brazil; ⁶Unidade de Estabilização de Animais Marinhos, Universidade do Vale de Itajaí (UNIVALI), Penha, SC, Brazil; ⁷Centro de Estudos Costeiros, Limnológicos e Marinhos (CECLIMAR), Universidade Federal do Rio Grande do Sul (UFRGS), Imbé, RS, Brazil; ⁸Associação R3 Animal, Florianópolis, SC, Brazil; ⁹Laboratório de Parasitologia, Instituto de Pesquisas Veterinárias Desidério Finamor (IPVDF), Eldorado do Sul, RS, Brazil; ¹⁰Departamento de Educação Científica e Tecnológica, Universidade do Estado de Santa Catarina – UDESC, Florianópolis, SC, Brazil and ¹¹Centre for Ecology and Conservation, University of Exeter, Penryn, UK

Abstract

Emerging infectious diseases pose threats to wildlife, particularly in geographically isolated populations where hosts may lack prior exposure and immunity. Seabirds inhabiting remote islands in the southwest Atlantic and Southern Ocean, including threatened albatrosses and petrels, are increasingly affected by infectious pathogens. However, baseline data on vector-borne infections in these species remain scarce. This study assessed the presence of vector-borne haemosporidian parasites (*Plasmodium*, *Haemoproteus* and *Leucocytozoon*) and bacterial pathogens (*Borrelia burgdorferi sensu lato*, *Anaplasma* and *Ehrlichia*) in albatrosses and petrels, providing insights into disease prevalence and potential threats to these populations. We analysed blood and tissue samples from 269 individuals of 5 albatross and 12 petrel species, collected over an 11-year period (2013–2023) from South Georgia and multiple sites along the Brazilian coastline. Molecular assays, including nested Polymerase Chain Reaction (PCR), were used for pathogen screening. Blood smears from birds sampled in South Georgia were also examined for haemoparasites via light microscopy. We found no molecular or microscopy evidence of infection with haemosporidian parasites, *Borrelia*, *Anaplasma* or *Ehrlichia* in any of the samples. These findings suggest that vector-borne pathogens are either absent or at low prevalence, possibly because of limited vector presence, natural resistance or historical isolation from infection. Continuous monitoring is critical given current environmental changes and risks of pathogen introduction via climate-driven shifts in vector distribution. Our study establishes an essential baseline for future disease surveillance, prevention and mitigation in albatrosses and petrels, underscoring the importance of long-term monitoring to detect emerging pathogens in vulnerable seabird populations.

Introduction

Emerging infectious diseases can have devastating impacts on hosts, sometimes with consequences for entire ecosystems (Machalaba *et al.*, 2020; Nicholson *et al.*, 2020; Ferreyra *et al.*, 2022; Jain, 2023). For example, the pandemic of high pathogenicity avian influenza (HPAI H5Nx) resulted in the death of over half a billion wild birds and poultry worldwide within just a few years of emergence in the early 2020s (Gamarra-Toledo *et al.*, 2023; Roberts *et al.*, 2023; Puryear and Runstadler, 2024). Diseases can be particularly severe in areas where hosts are shielded geographically from pathogen exposure, particularly those on remote islands (Wyatt *et al.*, 2008; Atkinson and LaPointe, 2009; Vanstreels *et al.*, 2023). Such threats to isolated populations are often exacerbated by anthropogenic effects, such as global connectivity and climate change, in part through effects on pathogen and vector survival, distribution and spread (Steig *et al.*, 2009; Lynton-Jenkins *et al.*, 2021). Monitoring the health and infection status of vulnerable species inhabiting remote locations can therefore be critical for their conservation.

© The Author(s), 2025. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

The islands of the southwest Atlantic and Southern Ocean are home to a rich diversity of bird species, including many endemics or globally important breeding populations (Poncet et al., 2020; Favero et al., 2024; Poncet et al., 2017). This includes highly threatened albatrosses and petrels, many of which are declining because of incidental mortality (bycatch) in fisheries, predation by invasive species, climate change, or degradation/loss of nesting habitat (Dias et al., 2019; Phillips et al., 2023; Baker et al., 2024). Until recently, there have been few documented cases of mass mortality on islands where albatrosses and petrels breed, but it is unclear whether this is because these populations are largely shielded from infection (Vanstreels et al., 2023), or because of a lack of attention to their health and disease status (Uhart et al., 2018). Avian cholera (*Pasteurella multocida*) is, however, thought to have caused mortality in albatrosses and petrels in the 1980s, with more recent infections in asymptomatic individuals raising questions around the reservoir potential of these species (Gamble et al., 2019). Overall, outbreaks of infectious diseases in albatrosses and petrels appear to be on the rise, e.g. avian cholera and *Erysipelothrix rhusiopathiae* cause yearly recurrent die-offs of the Indian yellow-nosed albatross (*Thalassarche carteri*) on Amsterdam Island (Jaeger et al., 2020). In 2024, an outbreak of HPAI H5N1 on South Georgia caused mortality in multiple taxa, including wandering albatrosses (*Diomedea exulans*), which are listed as Vulnerable by the International Union for the Conservation of Nature, as well as brown skuas (*Stercorarius antarcticus*), gentoo penguins (*Pygoscelis papua*) and Antarctic fur seals (*Arctocephalus gazella*) (Bennison et al., 2024). Given increasing likelihood of disease outbreaks at breeding sites (Banyard et al., 2024), the establishment of continuous monitoring programmes of albatross and petrel health is integral to the rapid detection of emerging diseases and effective predictions of disease spread based on prior infection history (Vanstreels et al., 2023).

Infectious pathogens at risk of emergence in seabirds breeding on remote islands include vector-borne haemosporidian parasites in the genera *Plasmodium*, *Haemoproteus* and *Leucocytozoon* (Quillfeldt et al., 2011; Parsons et al., 2017; Vanstreels et al., 2018; Muñoz-Leal et al., 2019), which cause avian malaria or malaria-like diseases (Valkiunas, 2005; Palinauskas et al., 2008; Bensch et al., 2009; Pacheco and Escalante, 2023). Infections with these parasites have been associated with increased mortality, population declines and even extinctions (Warner, 1968; Hill et al., 2010). Sublethal effects include reductions in body condition, sexual ornamentation or reproductive success, as well as prolonged stopovers or delayed migration (Martínez-Abraín et al., 2004; Marzal et al., 2005, 2008; Barbosa and Palacios, 2009; Quillfeldt et al., 2011; Hegemann et al., 2018). In seabirds, infection with *Plasmodium relictum*, *P. circumflexum* and *P. vaughani* has been associated with increased mortality in captive Humboldt and Magellanic penguins (*Spheniscus demersus* and *S. magellanicus*), little penguins (*Eudyptula minor*) and Atlantic puffins (*Fratercula arctica*) (Sallaberry-Pincheira et al., 2015; Sijbranda et al., 2017; Meister et al., 2021).

Other haemoparasites isolated recently from seabirds at risk of emergence include the bacteria *Borrelia* (Dietrich et al., 2011; Schramm et al., 2014; Parsons et al., 2017; Vanstreels et al., 2023), *Anaplasma* and *Ehrlichia* (Anaplasmataceae) (Vanstreels et al., 2018; Muñoz-Leal et al., 2019). These parasites are transmitted through ticks such as *Ixodes* spp. and *Argasidae* spp., which are common in many seabird colonies (Dietrich et al., 2011, 2014; Vanstreels et al., 2018; Muñoz-Leal et al., 2019). *Borrelia* parasites have been isolated from *Ixodes uriae* ticks collected in colonies

of razorbills (*Alca torda*), Atlantic puffins (*F. arctica*) and black-browed albatrosses (*Thalassarche melanophris*) (Olsen et al., 1993, 1995; Gylfe et al., 1999; Munro et al., 2019). Such infections can be pathogenic in seabirds (Yabsley et al., 2012; Parsons et al., 2018; Vanstreels et al., 2019), as evidenced by a *Borrelia*-positive African penguin (*S. demersus*) that presented antemortem neurological signs and lesions similar to those reported in an owl fatally infected with *B. hermsii* and a domestic fowl infected with *B. anserina* (Dickie and Barrera, 1964; Thomas et al., 2002; Bunikis et al., 2004; Yabsley et al., 2012). *Anaplasma phagocytophilum* has been detected in blood samples from passerines, suggesting their potential role in transmitting the bacterium to ticks; however, the significance of wild birds in the infectious cycle of this parasite remains unclear (Pedersen et al., 2020). Although only mammals have so far been confirmed as competent hosts and reservoirs for *Ehrlichia* bacteria (Rar and Golovljova, 2011), there is growing evidence that the host range may extend to other vertebrates after the detection in wild birds in Brazil of *Ehrlichiae* typically associated with ungulates and carnivores (Machado et al., 2012).

The goal of this study was to survey vector-borne infectious diseases in albatrosses and petrels sampled in the southwest Atlantic and Southern Ocean, with the aim of detecting any novel infection, as well as establishing baselines for future surveillance. We screened 5 species of albatrosses and 12 species of petrels for the presence of haemosporidian parasites (*Plasmodium*, *Haemoproteus* and *Leucocytozoon*), and the bacterial pathogens *Anaplasma*, *Ehrlichia* and *B. burgdorferi sensu lato* (s.l.). Understanding host-pathogen interactions in these species will be key to informing conservation strategies of these threatened species and mitigating potential disease threats in a rapidly changing environment.

Materials and methods

Sampling

The samples were obtained between 2013 and 2023 from 4 different sources: (i) breeding adults and chicks at colonies on Bird Island, South Georgia (54°00'S, 38°03'W); (ii) seabirds bycaught in Brazilian fisheries; (iii) birds found in a weakened state on the Brazilian coast and taken to 2 rehabilitation centres; and (iv) seabirds monitored in stranding networks in Brazil (Table 1). Detailed fieldwork procedures, methods of sampling and locations of seabirds surveyed are provided in Supplementary Table S1.

DNA extraction and molecular analysis

A total of 96 DNA samples from the seabirds sampled in South Georgia were extracted using a QIAamp DNA Micro Kit (Qiagen, Hilden, Germany) following the manufacturer's protocol. For all other samples (Table 1), DNA was extracted using the phenol-chloroform method (Sambrook et al., 1989), with the exception of blood samples preserved in ethanol (Supplementary Table S1), which were extracted using a slightly modified approach (Maia et al., 2017). First, we tested for the presence of haemosporidians (*Plasmodium*, *Haemoproteus* and *Leucocytozoon*) in a subset of samples ($N = 96$ blood samples from South Georgia and 30 liver samples), using a nested PCR targeting a 617-bp fragment of the cytochrome b gene (Hellgreen et al., 2004) and including DNA from *Plasmodium gallinaceum* strain A8 (MalAvi lineage GALLUS01), *Haemoproteus majoris* (MalAvi lineage PARUS1) and *Leucocytozoon* (MalAvi lineage PARUS22) as positive controls. Second, we tested for the presence of *Borrelia burgdorferi sensu lato*

Table 1. Albatrosses and petrels screened for vector-borne parasites from 2013 to 2023 from the southern Brazilian Coast, Brazilian fisheries and from colonies in Bird Island, South Georgia

Family	Species	Sampling location	No. of individuals tested	Haemoparasites investigated
Diomedidae	Wandering albatross <i>Diomedea exulans</i>	Bird Island, South Georgia	50	<i>Plasmodium</i> , <i>Haemoproteus</i> , <i>Leucocytozoon</i> , <i>Borrelia burgdorferi</i> s.l.
	Tristan albatross <i>Diomedea dabbenena</i>	Rehabilitation centres and fisheries bycaught birds, Brazil	1	<i>Anaplasma</i> , <i>Ehrlichia</i>
	Southern royal albatross <i>Diomedea epomophora</i>	Rehabilitation centres and fisheries bycaught birds, Brazil	1	<i>Anaplasma</i> , <i>Ehrlichia</i>
	Atlantic yellow-nosed albatross <i>Thalassarche chlororhynchos</i>	Rehabilitation centres and fisheries bycaught birds, Brazil	15	<i>Anaplasma</i> , <i>Ehrlichia</i>
	Black-browed albatross <i>Thalassarche melanophris</i>	Rehabilitation centres, stranding network and fisheries bycaught birds, Brazil	29	<i>Plasmodium</i> , <i>Haemoproteus</i> , <i>Anaplasma</i> , <i>Ehrlichia</i>
Procellariidae	Great shearwater <i>Ardenna gravis</i>	Rehabilitation centres and fisheries bycaught birds, Brazil	6	<i>Anaplasma</i> , <i>Ehrlichia</i>
	Sooty shearwater <i>Ardenna grisea</i>	Rehabilitation centres and fisheries bycaught birds, Brazil	2	<i>Anaplasma</i> , <i>Ehrlichia</i>
	Cory's shearwater <i>Calonectris borealis</i>	Rehabilitation centres and fisheries bycaught birds, Brazil	10	<i>Anaplasma</i> , <i>Ehrlichia</i>
	Cape petrel <i>Daption capense</i>	Rehabilitation centres and fisheries bycaught birds, Brazil	3	<i>Anaplasma</i> , <i>Ehrlichia</i>
	Southern fulmar <i>Fulmarus glacialis</i>	Rehabilitation centres and fisheries bycaught birds, Brazil	1	<i>Anaplasma</i> , <i>Ehrlichia</i>
	Southern giant petrel <i>Macronectes giganteus</i>	Rehabilitation centres and fisheries bycaught birds, Brazil	8	<i>Anaplasma</i> , <i>Ehrlichia</i>
	Northern giant petrel	Rehabilitation centres and fisheries bycaught birds, Brazil; Bird Island, South Georgia	33	<i>Anaplasma</i> , <i>Ehrlichia</i> , <i>Plasmodium</i> , <i>Haemoproteus</i> , <i>Leucocytozoon</i> , <i>Borrelia burgdorferi</i> s.l.
	White-chinned petrel <i>Procellaria aequinoctialis</i>	Rehabilitation centres and fisheries bycaught birds, Brazil; Bird Island, South Georgia	61	<i>Anaplasma</i> , <i>Ehrlichia</i> , <i>Plasmodium</i> , <i>Haemoproteus</i> , <i>Leucocytozoon</i> , <i>Borrelia burgdorferi</i> s.l.
	Atlantic petrel <i>Pterodroma incerta</i>	Rehabilitation centres and fisheries bycaught birds, Brazil	1	<i>Anaplasma</i> , <i>Ehrlichia</i>
	Soft-plumaged petrel <i>Pterodroma mollis</i>	Rehabilitation centres and fisheries bycaught birds, Brazil	1	<i>Anaplasma</i> , <i>Ehrlichia</i>
Manx shearwater <i>Puffinus puffinus</i>	Rehabilitation centres and fisheries bycaught birds, Brazil	42	<i>Anaplasma</i> , <i>Ehrlichia</i>	
Oceanitidae	Wilson's storm petrel <i>Oceanites oceanicus</i>	Rehabilitation centres and fisheries bycaught birds, Brazil	5	<i>Anaplasma</i> , <i>Ehrlichia</i>

(s.l.) using a nested PCR, which targets the 5S–23S rRNA spacer region of all *Borrelia* species and the same positive controls mentioned by Newman et al. (2015). Finally, we tested for the presence of *Ehrlichia* and *Anaplasma* in 219 tissue samples using a PCR protocol specifically designed to amplify a 345-bp fragment of the 16S rRNA gene (Parola et al., 2000); *Anaplasma marginale* DNA was used as a positive control. In all assays, we included ultrapure water as negative controls.

Light microscopy of blood smears

The 96 blood smears taken from the live seabirds at South Georgia were analysed by light microscopy, following Merino et al. (1997) and Quillfeldt et al. (2010). In brief, one half of the slide was scanned at $\times 200$ magnification to look for haemoparasites, and at least 20 fields in the other half at $\times 400$ to look for intracellular stages of haematozoa. A minimum of 2000 to 10 000 erythrocytes were also checked using the oil immersion objective ($1000\times$).

Results

None of the parasites of interest were detected in the 269 sampled seabirds (Table 1). This included no evidence of infection with either *Plasmodium*, *Haemoproteus* or *Leucocytozoon* parasites in the 96 samples from live albatrosses and petrels obtained in South Georgia, nor of infection with *Plasmodium* or *Haemoproteus* in the liver samples from wandering albatross chicks (18) or black-browed albatrosses (12) found dead at South Georgia or in southern Brazil. We found no evidence of infection with *Borrelia* spp. (Table 1) in any of the 96 samples from live albatrosses and petrels obtained in South Georgia, or with *Anaplasmataceae* parasites in any of the 219 tissue samples from seabirds found at the Brazilian coast or obtained as bycatch. Microscopic observations were consistent with molecular findings in that we did not detect the presence of parasitized erythrocytes in any of the 96 blood smears obtained from live wandering albatrosses, northern giant petrels and white-chinned petrels at South Georgia.

Discussion

Molecular and microscopy methods were used to screen for the presence of key blood parasites in a total of 269 birds from 5 species of albatrosses and 12 species of petrels sampled in the southwest Atlantic and Southern Ocean over a period of 11 years (2013–2023). None of the samples were found to be infected with parasites of the genera *Plasmodium*, *Leucocytozoon*, *Haemoproteus*, *Borrelia*, *Anaplasma* or *Ehrlichia*. Although detecting parasites in blood smears through microscopy is challenging when infection intensity is low (Valkiunas, 2005), the high sensitivity of molecular methods increases our confidence that none of the birds were infected. These results are consistent with other investigations of blood parasite infections conducted in seabirds of the Antarctic region (Laird, 1961; Quillfeldt et al., 2010; Llanos et al., 2018). Screening of 455 birds from 14 species sampled between 1975 and 1978 in South Georgia found that none of the samples tested positive for blood parasites, except a small proportion of wandering, black-browed and grey-headed albatrosses, which were infected with the previously undescribed *Hepatozoon albatrossi* (Peirce and Prince, 1980). Documented cases of *Hepatozoon* in the Antarctic and subantarctic regions are limited to *Hepatozoon albatrossi*, reported in albatrosses and storm petrels (Merino et al., 2014; Parsons et al., 2017). Our study aimed to screen for other parasites with poorly known occurrence, and excluded *Hepatozoon* because its presence is already established. Our findings provide further evidence that most seabird species are likely relatively free of blood parasites (Quillfeldt et al., 2011).

The lack of blood parasites in the samples may be explained partly by the harsh environmental conditions at the breeding grounds, which may preclude vector persistence (Martínez-Abraín et al., 2004). To date, there are no records of vectors such as hematophagous ceratopogonid or hippoboscid biting flies, or *Aedes* mosquitoes, in subantarctic or Antarctic islands where most of the study species nest (Quillfeldt et al., 2011; Ferreira et al., 2020). Such vectors may, however, occur in feeding areas, making disease transmission possible from infected coastal seabird or land bird species. Infection at non-breeding grounds is thought to drive the incidence of *Leucocytozoon* and *Plasmodium* lineages in Caspian gulls (*Larus cachinnans*) wintering on the coast of Poland (Zagalska-Neubauer and Bensch, 2016), and of *Haemoproteus* spp. in Manx shearwaters and black-browed albatrosses off the coast of Brazil (Sgarioni et al., 2024), since these species tested negative

at their breeding colonies (Quillfeldt et al., 2011). The petrel and albatross species that were sampled in South Georgia, including white-chinned petrels, wandering albatrosses and northern giant petrels, can travel to the Patagonian Shelf or shelf-break to feed during the breeding season, and stay there for part or all of the non-breeding season (Phillips et al., 2006; González-Solís et al., 2007; Froy et al., 2015; Clay et al., 2019; Granroth-Wilding and Phillips, 2019). The lower latitudes of the Patagonian Shelf and shelf-break, compared to the Antarctic and sub-Antarctic regions, increase the likelihood of vector presence and infection risk, with the potential for subsequent transmission by returning migrants to colonies during the breeding season (Quillfeldt et al., 2011; Sallaberry-Pincheira et al., 2015). Indeed, haemosporidian parasites, such as *P. relictum*, *P. circumflexum* and *P. vaughani*, as well as *Borrelia* parasites, can be found in migratory birds (McDiarmid, 1969; Wolcott et al., 2021; Bennett et al., 2024), and seabirds are thought to play a role in the dispersal of *Anaplasma* and *Ehrlichia* bacteria (Vanstreels et al., 2018; Muñoz-Leal et al., 2019). As such, the routine surveillance for blood parasite infections in seabirds in the southwest Atlantic and Southern Ocean is essential for detecting any shifts of disease vectors to higher latitudes, thereby increasing risks of range expansions for vector-borne parasites.

While the lack of detectable blood parasites is reassuring, seabirds of the southwest Atlantic and Southern Ocean remain at risk of future infections. *Ehrlichia* spp. was recorded recently in southern Chile in an *I. uriae* tick (Muñoz-Leal et al., 2019), which is a common parasite of seabirds on Antarctic and subantarctic islands (Vanstreels et al., 2020), and new *Plasmodium* infections were recently recorded in seabirds in New Zealand, including in a procellariiform species, the Westland petrel (*Procellaria westlandica*) (Bennett et al., 2024). Many albatrosses and petrels already face numerous other threats, so novel infections are potentially serious additional risks (Heard et al., 2013; Dias et al., 2019; Phillips et al., 2023). As a result, developing conservation and management plans for seabirds of the southwest Atlantic and Southern Ocean will require the implementation of proactive risk assessments, biosecurity measures and long-term disease surveillance, particularly at important breeding and non-breeding areas (Uhart et al., 2018). We recommend that surveillance programmes of blood parasite infections go beyond relying only on visual observations of diseased birds, and incorporate active screenings of seabird samples.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0031182025100590>.

Acknowledgements. We are very grateful to the field assistants who helped with the sampling at Bird Island. The authors are also grateful to all teams and volunteers dedicated to work with monitoring sick and dead seabirds stranded along the Brazilian coast as part of the Beach Monitoring Project of Petrobras, as well as for all teams engaged providing samples and data for the biobank 'Banco Nacional de Amostras Biológicas de Albatrozes e Petreus' (BAAP), which holds samples of birds bycaught in Brazilian fisheries and from seabirds stranding networks in Brazil.

Author contributions. P.P.S., A.Z.S., C.B. and R.A.P. conceptualized the study. R.A.P., A.P., P.P.S., T.E., T.P.P., D.B.A., C.M.K., A.C.D.B. and K.H.L. were responsible for samples and data curation. P.P.S. and A.Z.S. conducted laboratory screening and diagnosis. P.P.S. conducted the literature searches, performed formal analysis and wrote the original draft of the manuscript. R.A.P., A.O.S.L., G.K., J.R., A.C.D.B., K.H.L. and C.B. were responsible for validation, supervision and final review & editing. All authors revised and edited the manuscript providing input for the manuscript structure and topics. P.P.S. and A.Z.S. constructed the table and graphical abstract.

Financial support. This work represents a contribution to the Ecosystems component of the British Antarctic Survey Polar Science for a Sustainable Planet Programme, funded by the Natural Environment Research Council. Part of this work was carried out during a secondment funded by the Agreement on the Conservation of Albatrosses and Petrels. The studied birds were mainly collected and necropsied as part of the Beach Monitoring Project of the Santos Basin (Projeto de Monitoramento de Praias – PMP/BS). PMP is one of the monitoring programs required by Brazil's federal environmental agency, the Institute of the Environment and Renewable Natural Resources (IBAMA), for the environmental licensing process of oil production and transport by Petrobras. Bairy and K.H. Lückmann are recipients of a productivity fellowship from CNPq (311725/2021-0 and 313843/2023-6, respectively). We are thankful to CEMAVE/ICMBio, Petrobras, IBAMA, LABCAI/UFSC and R3 Animal for their continued support, and the Wildlife Conservation Network for providing a veterinary scholarship to A. Z. Sgarioni.

Competing interests. The authors declare there are no conflicts of interest.

Ethical standards. Sampling of live birds at Bird Island was approved by the British Antarctic Survey Animal Welfare and Ethics Committee and carried out with the permission of the Government of South Georgia and the South Sandwich Islands. Sampling from stranded and deceased seabirds in this project was undertaken with all the necessary permits issued by SISBIO/ICMBio and IBAMA, environmental agencies of Brazil.

References

- Atkinson CT and LaPointe DA (2009) Introduced avian diseases, climate change, and the future of Hawaiian honeycreepers. *Journal of Avian Medicine and Surgery* 23(1), 53–63. doi:10.1647/2008-059.1
- Baker GB, Komyakova V, Wellbelove A, Beynon N and Haward M (2024) The implementation of ACAP Best Practice Advice to mitigate seabird bycatch in fisheries: Issues and options. *Marine Policy* 160, 105879. doi:10.1016/j.marpol.2023.105879
- Banyard AC, Bennison A, Byrne AMP, et al. (2024) Detection and spread of high pathogenicity avian influenza virus H5N1 in the Antarctic Region. *Nature Communications* 15, 7433. doi:10.1038/s41467-024-51490-8.
- Barbosa A and Palacios MJ (2009) Health of Antarctic birds: A review of their parasites, pathogens and diseases. *Polar Biology* 32, 1095. doi:10.1007/s00300-009-0640-3
- Bennett J, Niebuhr CN, Lagrue C, Middlemiss KL, Webster T and Filion A (2024) New insights into avian malaria infections in New Zealand seabirds. *Parasitology Research* 123(4), 184. doi:10.1007/s00436-024-08210-4
- Bennison A, Adlard S, Banyard AC, Blockley F, Blyth M, Browne E and Phillips RA (2024) A case study of highly pathogenic avian influenza (HPAI) H5N1 at Bird Island, South Georgia: The first documented outbreak in the subantarctic region. *Bird Study*, 1–12. doi:10.1080/00063657.2024.2396563
- Bensch S, Hellgren O and Pérez-Tris J (2009) MalAvi: A public database of malaria parasites and related haemosporidians in avian hosts based on mitochondrial cytochrome b lineages. *Molecular Ecology Resources* 9(5), 1353–1358. doi:10.1111/j.1755-0998.2009.02692.x
- Bunikis J, Garpmo U, Tsao J, Berglund J, Fish D and Barbour AG (2004) Sequence typing reveals extensive strain diversity of the Lyme borreliosis agents *Borrelia burgdorferi* in North America and *Borrelia afzelii* in Europe. *Microbiology* 150(Pt 6), 1741–1755. doi:10.1099/mic.0.26944-0
- Clay TC, Small C, Tuck GN, Pardo D, Carneiro APB, Wood AG, Croxall JP, Crossin GT and Phillips RA (2019) A comprehensive large-scale assessment of fisheries bycatch risk to threatened seabird populations. *Journal of Applied Ecology* 56, 1882–1893. doi:10.1111/1365-2664.13407
- Dias MP, Martin R, Pearmain EJ, Burfield IJ, Small C, Phillips RA, Yates O, Lascelles B, Borborolu PG and Croxall JP (2019) Threats to seabirds: A global assessment. *Biological Conservation* 237, 525–537. doi:10.1016/j.biocon.2019.06.033
- Dickie CW and Barrera J (1964) A study of the carrier state of avian spirochetosis in the chicken. *Avian Diseases* 8(2), 191–195. doi:10.2307/1588059
- Dietrich M, Gómez-Díaz E and McCoy KD (2011) Worldwide distribution and diversity of seabird ticks: Implications for the ecology and epidemiology of tick-borne pathogens. *Vector Borne and Zoonotic Diseases* 11(5), 453–470. doi:10.1089/vbz.2010.0009
- Dietrich M, Lebarbenchon C, Jaeger A, Le Rouzic C, Bastien M, Lagadec E, McCoy KD, Pascalis H, Le Corre M, Dellagi K and Tortosa P (2014) Rickettsia spp. in Seabird Ticks from Western Indian Ocean Islands, 2011–2012. *Emerging Infectious Diseases* 20(5), 838–842. doi:10.3201/eid2005.131088
- Favero M, Seco Pon JP, Paz J, Hernandez M and Copello S (2024) Seabirds in the Argentine Continental Shelf and Shelf-Break. Acha EM, Iribarne OO and Piola A, (eds.) *The Patagonian Shelfbreak Front. Aquatic Ecology Series*. Vol. 13. Cham: Springer, 185–209. doi:10.1007/978-3-031-71190-9_8
- Ferreira FC, Santiago-Alarcon D and Braga EM (2020) Diptera vectors of avian Haemosporidians: With emphasis on tropical regions. In Santiago-Alarcon D and Marzal A (eds), *Avian Malaria and Related Parasites in the Tropics: Ecology, Evolution and Systematics*. Cham: Springer, 185–250.
- Ferreira HDV, Rudd J, Foley J, Vanstreels RE, Martín AM, Donadio E and Uhart MM (2022) Sarcocystis mangle outbreak decimates South American wild camelid populations in San Guillermo National Park, Argentina. *PLoS ONE* 17(1), e0256616. doi:10.1371/journal.pone.0256616
- Froy H, Lewis S, Catry P, Bishop CM, Forster IP, Fukuda A, Higuchi H, Phalan B, Xavier JC, Nussey DH and Phillips RA (2015) Age-related variation in foraging behaviour in the wandering albatross at South Georgia: No evidence for senescence. *PLoS ONE* 10, e0116415. doi:10.1371/journal.pone.0116415
- Gamarrá-Toledo V, Plaza PI, Angulo F, Gutiérrez R, García-Tello O, Saravia-Guevara P, Mejía-Vargas F, Epiquién-Rivera M, Quiroz-Jiménez G, Martínez P, Huamán-Mendoza D, Inga-Díaz G, La Madrid LE, Luyo P, Ventura S and Lambertucci SA (2023) Highly pathogenic avian influenza (HPAI) strongly impacts wild birds in Peru. *Biological Conservation* 286, 110272. doi:10.1016/j.biocon.2023.110272
- Gamble A, Garnier R, Jaeger A, Gantelet H, Thibault E, Tortosa P, Bourret V, Thiebot JB, Delord K, Weimerskirch H, Tornos J, Barbraud C and Boulignier T (2019) Exposure of breeding albatrosses to the agent of avian cholera: Dynamics of antibody levels and ecological implications. *Oecologia* 189, 939–949. doi:10.1007/s00442-019-04369-1
- González-Solis J, Croxall JP and Afanasyev V (2007) Offshore spatial segregation in giant petrels *Macronectes* spp.: Differences between species, sexes and seasons. *Aquatic Conservation: Marine and Freshwater Ecosystems* 17(S1), 22–36. doi:10.1002/aqc.911
- Granroth-Wilding HMV and Phillips RA (2019) Segregation in space and time explains the coexistence of two sympatric sub-Antarctic petrels. *Ibis* 161, 101–116. doi:10.1111/ibi.12584
- Gylfe Á, Olsen B, Straševićius D, Marti Ras N, Weihe P, Noppa L and Bergström S (1999) Isolation of Lyme disease *Borrelia* from puffins (*Fratercula arctica*) and seabird ticks (*Ixodes uriae*) on the Faeroe Islands. *Journal of Clinical Microbiology* 37(4), 890–896. doi:10.1128/jcm.37.4.890-896.1999
- Heard MJ, Smith KF, Ripp KJ, Berger M, Chen J, Dittmeier J, Goter M, Mccarvey ST and Ryan E (2013) The threat of disease increases as species move toward extinction. *Conservation Biology* 27, 1378–1388. doi:10.1111/cobi.12143
- Hegemann A, Abril PA, Muheim R, Sjöberg S, Alerstam T, Nilsson J and Hasselquist D (2018) Immune function and blood parasite infections impact stopover ecology in passerine birds. *Oecologia* 188, 1011–1024. doi:10.1007/s00442-018-4291-3
- Hellgreen O, Waldenstrom J and Bensch S (2004) A New PCR assay for simultaneous studies of *Leucocytozoon*, *Plasmodium*, and *Haemoproteus* from avian blood. *Journal of Parasitology* 90, 797–802. doi:10.1645/GE-184R1
- Hill AG, Howe L, Gartrell BD and Alley MR (2010) Prevalence of *Leucocytozoon* spp. in the endangered yellow-eyed penguin *Megadyptes antipodes*. *Parasitology* 137, 1477–1485. doi:10.1017/S0031820099919190
- Jaeger A, Gamble A, Lagadec E, Lebarbenchon C, Bourret V, Tornos J, Barbraud C, Lemberger K, Delord K, Weimerskirch H, Thiebot JB, Boulignier T and Tortosa P (2020) Impact of annual bacterial epizootics on albatross population on a remote island. *EcoHealth* 17, 194–202. doi:10.1007/s10393-020-01487-8

- Jain NR (2023) The first animal disease to be eradicated in the history of pandemic-rinderpest. *mLAC Journal for Arts, Commerce and Sciences* 1(1), 2–8. doi:10.59415/mjacs.v1i1.77
- Laird M (1961) A lack of avian and mammalian haematozoa in the Antarctic and Canadian Arctic. *Canadian Journal of Zoology* 39(2), 209–213. doi:10.1139/z61-025
- Llanos S, Suazo CG, Quillfeldt P, Cursach JA and Salas LM (2018) Ectoparasite abundance and apparent absence of hemoparasites in two albatross species in Sub-Antarctic Chile. *Polar Biology* 41, 143–147. doi:10.1007/s00300-017-2177-1
- Lynton-Jenkins JG, Russell AF, Chaves J and Bonneaud C (2021) Avian disease surveillance on the island of San Cristobal, Galapagos. *Ecology and Evolution* 00, 1–12. doi:10.1002/ece3.8431
- Machado RZ, André MR, Werther K, de Sousa E, Gavioli FA and Alves Junior JRF (2012) Migratory and carnivorous birds in Brazil: Reservoirs for *Anaplasma* and *Ehrlichia* species? *Vector-Borne and Zoonotic Diseases* 12(8), 705–708. doi:10.1089/vbz.2011.0803
- Machalaba C, Feferholtz Y, Uhart MM and Karesh WB (2020) Wildlife conservation status and disease trends: Ten years of reports to the worldwide monitoring system for wild animal diseases. *Revue Scientifique et Technique (International Office of Epizootics)* 39(3), 991–1001. doi:10.20506/rst.39.3.3191
- Maia TA, Vilaça ST, Silva LRD, Santos FR and Dantas GPDM (2017) DNA sampling from eggshells and microsatellite genotyping in rare tropical birds: Case study on Brazilian Merganser. *Genetics and Molecular Biology*, 40:808–812. doi:10.1590/1678-4685-GMB-2016-0297
- Martínez-Abraín A, Esparza B and Oro D (2004) Lack of blood parasites in bird species: Does absence of blood parasite vectors explain it all? *Ardeola* 51(1), 225–232.
- Marzal A, Bensch S, Reviriego MI, Balbontin J and De Lope F (2008) Effects of malaria double infection in birds: One plus one is not two. *Journal of Evolutionary Biology*. 21, 979–987. doi:10.1111/j.1420-9101.2008.01545.x
- Marzal A, de Lope F, Navarro C and Møller AP (2005) Malarial parasites decrease reproductive success: An experimental study in a passerine bird. *Oecologia* 142, 541–545. doi:10.1007/s00442-004-1757-2
- McDiarmid A (1969) In *Diseases in Free-living Wild Animals*. London: Academic Press, 332.
- Meister SL, Richard OK, Hoby S, Gurtner C and Basso WU (2021) Fatal avian malaria in captive Atlantic puffins (*Fratercula arctica*) in Switzerland. *International Journal for Parasitology: Parasites and Wildlife* 14, 97–106. doi:10.1016/j.ijppaw.2020.12.007
- Merino S, Martínez J, Masello JF, Bedolla Y and Quillfeldt P (2014) First molecular characterization of a *Hepatoozon* species (Apicomplexa: Hepatozoidae) infecting birds and description of a new species infecting storm petrels (Aves: Hydrobatidae). *The Journal of Parasitology* 100(3), 338–343. doi:10.1645/13-325.1
- Merino S, Potti J and Fargallo JA (1997) Blood parasites of passerine birds from central Spain. *Journal of Wildlife Diseases* 33(3), 638–641. doi:10.7589/0090-3558-33.3.638
- Muñoz-Leal S, Clemes YS, Lopes MG, Acosta ICL, Serpa MCA, Mayorga LFSP, Gennari SM, González-Acuña D and Labruna MB (2019) Novel *Ehrlichia* sp. detected in Magellanic penguins (*Spheniscus magellanicus*) and in the seabird tick *Ixodes uriae* from Magdalena Island, southern Chile. *Ticks and Tick-borne Diseases* 10(6), 101256. doi:10.1016/j.ttbdis.2019.06.015
- Munro HJ, Ogden NH, Mechai S, Lindsay LR, Robertson GJ, Whitney H and Lang AS (2019) Genetic diversity of *Borrelia garinii* from *Ixodes uriae* collected in seabird colonies of the northwestern Atlantic Ocean. *Ticks and Tick-borne Diseases* 10, 101255. doi:10.1016/j.ttbdis.2019.06.014
- Newman EA, Eisen L, Eisen RJ, Fedorova N, Hasty JM, Vaughn C and Lane RS (2015) *Borrelia burgdorferi* sensu lato spirochetes in wild birds in northwestern California: Associations with ecological factors, bird behavior and tick infestation. *PLoS ONE* 10(2), e0118146. doi:10.1371/journal.pone.0118146
- Nicholson SK, Marneweck DG, Lindsey PA, Marnewick K and Davies-Mostert HT (2020) A 20-year review of the status and distribution of African wild dogs (*Lycaon pictus*) in South Africa. *African Journal of Wildlife Research* 50(1), 8–19. doi:10.3957/056.050.0008
- Olsen B, Duffy DC, Jaenson TG, Gylfe A, Bonnedahl J and Bergström S (1995) Transhemispheric exchange of Lyme disease spirochetes by seabirds. *Journal of Clinical Microbiology* 33(12), 3270–3274. doi:10.1128/jcm.33.12.3270-3274.1995
- Olsen B, Jaenson TG, Noppa L, Bunikis J and Bergström S (1993) A Lyme borreliosis cycle in seabirds and *Ixodes uriae* ticks. *Nature* 362(6418), 340–342. doi:10.1038/362340a0
- Pacheco MA and Escalante AA (2023) Origin and diversity of malaria parasites and other Haemosporida. *Trends in Parasitology* 39(7), 501–516. doi:10.1016/j.pt.2023.04.004
- Palinauskas V, Valkiunas G, Bolshakov CV and Bensch S (2008) *Plasmodium relictum* (lineage P- SGS1): Effects on experimentally infected passerine birds. *Experimental Parasitology* 120, 372–380. doi:10.1016/j.exppara.2008.09.001
- Parola P, Roux V, Camicas JL, Baradji I, Brouqui P and Raoult D (2000) Detection of ehrlichiae in African ticks by polymerase chain reaction. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 94(6), 707–708. doi:10.1016/s0035-9203(00)90243-8
- Parsons NJ, Gous TA, Cranfield MR, Cheng LI, Schultz A, Horne E, Last RP, Lampen F, Ludynia K, Bousfield B, Strauss V, Peirce MA and Vanstreels RET (2018) Novel vagrant records and occurrence of vector-borne pathogens in King Penguins (*Aptenodytes patagonicus*) in South Africa. *Polar Biology* 41(1), 79–86. doi:10.1007/s00300-017-2171-7
- Parsons NJ, Voogt NM, Schaefer AM, Peirce MA and Vanstreels RET (2017) Occurrence of blood parasites in seabirds admitted for rehabilitation in the Western Cape, South Africa, 2001–2013. *Veterinary Parasitology* 233, 52–61. doi:10.1016/j.vetpar.2016.12.001
- Pedersen BN, Jenkins A and Kjelland V (2020) Tick-borne pathogens in *Ixodes ricinus* ticks collected from migratory birds in southern Norway. *PLoS ONE* 15(4), e0230579. doi:10.1371/journal.pone.0230579
- Peirce MA and Prince PA (1980) *Hepatoozon albatrossi* sp. nov. (Eucoccidia: Hepatozoidae) from *Diomedea* spp. in the Antarctic. *Journal of Natural History* 14, 447–452. doi:10.1080/00222938000770381
- Phillips RA, Fort J and Dias MP (2023) Conservation status and overview of threats to seabirds. In Young LC, and VanderWerf EA (eds), *Conservation of Marine Birds*. Cambridge: Academic Press, Chapter 2, 33–56. doi:10.1016/B978-0-323-88539-3.00015-7
- Phillips RA, Silk JRD, Croxall JP and Afanasyev V (2006) Year-round distribution of white-chinned petrels from South Georgia: Relationships with oceanography and fisheries. *Biological Conservation* 129, 336–347. doi:10.1016/j.biocon.2005.10.046
- Poncet S, Wolfaardt AC, Barbraud C, Reyes-Arriagada R, Black A, Powell RB and Phillips RA (2020) The distribution, abundance, status and global importance of giant petrels (*Macronectes giganteus* and *M. halli*) breeding at South Georgia. *Polar Biology* 43(1), 17–34. doi:10.1007/s00300-019-02608-y
- Poncet S, Wolfaardt AC, Black A, Browning S, Lawton K, Lee J, Passfield K, Strange G and Phillips RA (2017) Recent trends in numbers of wandering (*Diomedea exulans*), black-browed (*Thalassarche melanophris*) and grey-headed (*T. chrysostoma*) albatrosses breeding at South Georgia. *Polar Biology* 40(7), 1347–1358. doi:10.1007/s00300-016-2057-0
- Puryear WB and Runstadler JA (2024) High-pathogenicity avian influenza in wildlife: A changing disease dynamic that is expanding in wild birds and having an increasing impact on a growing number of mammals. *Journal of the American Veterinary Medical Association* 1(aop), 1–9. doi:10.2460/javma.24.01.0053
- Quillfeldt P, Arriero E, Martínez J, Masello JF and Merino S (2011) Prevalence of blood parasites in seabirds—a review. *Frontiers in Zoology* 8, 26. doi:10.1186/1742-9994-8-26
- Quillfeldt P, Martínez J, Hennicke J, Ludynia K, Gladbach A, Masello JF, Riou S and Merino S (2010) Haemosporidian blood parasites in seabirds—a comparative genetic study of species from Antarctic to tropical habitats. *Naturwissenschaften* 97, 809–817. doi:10.1007/s00114-010-0698-3
- Rar V and Golovljova I (2011) *Anaplasma*, *Ehrlichia*, and “candidate *Neoehrlichia*” bacteria: Pathogenicity, biodiversity, and molecular genetic

- characteristics, a review. *Infection Genetics & Evolution* **11**(8), 1842–1861. doi:10.1016/j.meegid.2011.09.019
- Roberts LC, Abolnik C, Waller LJ, Shaw K, Ludynia K, Roberts DG, Kock AA, Makhado AB, Snyman A and Abernethy D** (2023) Descriptive epidemiology of and response to the high pathogenicity avian influenza (H5N8) epidemic in South African coastal seabirds, 2018. *Transboundary and Emerging Diseases* **1**, 2708458. doi:10.1155/2023/2708458
- Sallaberry-Pincheira N, Gonzalez-Acuña D, Herrera-Tello Y, Dantas GP, Luna-Jorquera G, Frere E, Valdés-Velasquez A, Simeone A and Vianna JA** (2015) Molecular epidemiology of avian malaria in wild breeding colonies of Humboldt and Magellanic penguins in South America. *EcoHealth* **12**, 267–277. doi:10.1007/s10393-014-0995-y
- Sambrook J, Fritsch ER and Maniatis T** (1989) *Molecular Cloning: A Laboratory Manual*. New York: Cold Spring Harbor.
- Schramm F, Gauthier-Clerc M, Fournier JC, McCoy KD, Barthel C, Postic D, Handrich Y, Maho YL and Jaulhac B** (2014) First detection of *Borrelia burgdorferi* sensu lato DNA in king penguins (*Aptenodytes patagonicus halli*). *Ticks and Tick-borne Diseases* **5**(6), 939–942. doi:10.1016/j.tbd.2014.07.013
- Sgarioni AZ, Serafini P, Pereira A, Emmerich T, Pontes TP, Machado DC, Ribeiro PR, Amorim DB, Klafke G and Reck J** (2024) Molecular survey of haemosporidian parasites in procellariiformes sampled in Southern Brazil, 2013–22. *The Journal of Wildlife Diseases* **60**(2), 413–420. doi:10.7589/JWD-D-23-00087
- Sijbranda DC, Hunter S, Howe L, Lenting B, Argilla L and Gartrell BD** (2017) Cases of mortality in little penguins (*Eudyptula minor*) in New Zealand associated with avian malaria. *New Zealand Veterinary Journal* **65**(6), 332–337. doi:10.1080/00480169.2017.1359124
- Steig EJ, Schneider DP, Rutherford SD, Mann ME, Comiso JC and Shindell DT** (2009) Warming of the Antarctic ice-sheet surface since the 1957 International Geophysical Year. *Nature* **457**(7228), 459–462. doi:10.1038/nature07669
- Thomas NJ, Bunikis J, Barbour AG and Wolcott MJ** (2002) Fatal spirochetosis due to a relapsing fever-like *Borrelia* sp. in a northern spotted owl. *Journal of Wildlife Diseases* **38**(1), 187–193. doi:10.7589/0090-3558-38.1.187
- Uhart MM, Gallo L and Quintana F** (2018) Review of diseases (pathogen isolation, direct recovery and antibodies) in albatrosses and large petrels worldwide. *Bird Conservation International* **28**(2), 169–196. doi:10.1017/S0959270916000629
- Valkiunas G** (2005) *Avian Malaria Parasites and Other Haemosporidia*, 1st edn. Boca Raton, FL, USA: CRC Press.
- Vanstreels RET, Palma RL and Mironov SV** (2020) Arthropod parasites of Antarctic and Subantarctic birds and pinnipeds: A review of host-parasite associations. *International Journal for Parasitology: Parasites and Wildlife* **12**, 275–290. doi:10.1016/j.ijppaw.2020.03.007
- Vanstreels RET, Parsons NJ, Pistorius PA and Schaefer AM** (2019) Prognostic indicators of immature rehabilitated African penguins (*Spheniscus demersus*) in South Africa. *The Journal of Wildlife Diseases* **55**, 758–769. doi:10.7589/2018-05-134
- Vanstreels RET, Uhart MM and Work TM** (2023) Health and diseases, (Young LC, and VanderWerf EA). (eds.) *Conservation of Marine Birds*. Cambridge: Academic Press, 131–176. doi:10.1016/B978-0-323-88539-3.0003-0.
- Vanstreels RET, Yabsley MJ, Parsons NJ, Swanepoel L and Pistorius PA** (2018) A novel candidate species of *Anaplasma* that infects avian erythrocytes. *Parasites and Vectors* **11**(1), 1–7. doi:10.1186/s13071-018-3089-9
- Warner RE** (1968) The role of introduced diseases in the extinction of the endemic Hawaiian Avifauna. *The Condor* **70**, 101–120. doi:10.2307/1365954
- Wolcott KA, Margos G, Fingerle V and Becker NS** (2021) Host association of *Borrelia burgdorferi* sensu lato: A review. *Ticks and Tick-borne Diseases* **12**(5), 101766. doi:10.1016/j.tbd.2021.101766
- Wyatt KB, Campos PF, Gilbert MTP, Kolokotronis SO, Hynes WH, DeSalle R and Greenwood AD** (2008) Historical mammal extinction on Christmas Island (Indian Ocean) correlates with introduced infectious disease. *PLoS ONE* **3**(11), e3602. doi:10.1371/annotation/9f9e687-682e-4010-97e4-139b33343d34
- Yabsley MJ, Parsons NJ, Horne EC, Shock BC and Purdee M** (2012) Novel relapsing fever *Borrelia* detected in African penguins (*Spheniscus demersus*) admitted to two rehabilitation centres in South Africa. *Parasitology Research* **110**(3), 1125–1130. doi:10.1007/s00436-011-2602-2
- Zagalska-Neubauer M and Bensch S** (2016) High prevalence of *Leucocytozoon* parasites in fresh water breeding gulls. *Journal of Ornithology* **157**, 525–532. doi:10.1007/s10336-015-1291-5