



Triassic outliers in the Peak District, central England, and a stratigraphical revision of the Miocene Brassington Formation

James B. Riding^{1*}, Vanessa J. Banks^{1,2}, Peter F. Jones² and Matthew J. Pound³

¹ British Geological Survey, Keyworth, Nottingham NG12 5GG, UK

² School of Sciences, University of Derby, Derby DE22 1GB, UK

³ School of Geography and Natural Sciences, Northumbria University, Newcastle upon Tyne NE1 8ST, UK

JBR, 0000-0002-5529-8989; VJB, 0000-0001-6335-7080; PFJ, 0009-0005-9504-8596; MJP, 0000-0001-8029-9548

* Correspondence: jbri@bgs.ac.uk

Abstract: The Miocene Brassington Formation of Derbyshire and Staffordshire in central England, as originally conceived, is preserved in sinkholes in the Lower Carboniferous/Mississippian Peak Limestone Group. However, contrary to the original Neogene age interpretation, most of the Brassington Formation is demonstrated herein to be of Triassic age, based upon gross lithofacies, material properties, palaeontology, palaeoclimatic considerations and palaeomagnetism. Specifically, the supposedly Miocene Kirkham and Bees Nest members are reinterpreted as the Triassic Sherwood Sandstone and Mercia Mudstone groups respectively. The Brassington Formation is therefore emended to exclude the Kirkham and Bees Nest members, and to include the original Kenslow Member and the new Friden Member. The latter unit is a clay-dominated succession exposed at Kenslow Top Pit, near Friden, Derbyshire. Both the Kenslow and Friden members have yielded Middle–Late Miocene pollen and spores. Evidentially, both the Lower–Middle Triassic Sherwood Sandstone and Mercia Mudstone groups were originally deposited in the White Peak area of Derbyshire and Staffordshire before subsiding into karstic voids. Geological implications of this revised interpretation include that the Sherwood Sandstone Group formerly extended further northwards in the Peak District than was previously supposed, and was laid down in a more complex and extensive depositional setting. The newly recognized Sherwood Sandstone Group and Mercia Mudstone Group successions in the Peak District have implications for the interpretation of hydrological evolution and karstification in the subjacent Peak Limestone Group. Furthermore, the palynology of the emended Brassington Formation may help to determine altitude of deposition, and hence provide improved understanding of Cenozoic uplift rates and interpretations of the drainage and landscape evolution of upland Britain.

Supplementary material: Background material to this article comprising three appendices, 14 figures and four tables is available at <https://doi.org/10.6084/m9.figshare.c.8179347>

Received 10 May 2025; **revised** 4 November 2025; **accepted** 7 November 2025

Karst landscapes can play an important role in preserving parts of sedimentary successions that might otherwise have been lost due to erosion (Coxon 2005). Sinkholes allow sediments to subside below erosion surfaces, thereby enhancing the survival of foundered strata. Important examples of this within upland Britain are the ‘pocket deposits’ found in ~60 karst cavities in the Peak Limestone Group (Lower Carboniferous/Mississippian) of Derbyshire and Staffordshire. These preserved sediments were formalized by Boulter *et al.* (1971) as the Brassington Formation (Figs 1, 2) and regarded as a stratigraphically coherent, fining-upwards succession entirely of Neogene age. As originally defined, this unit largely comprises the sand-dominated Kirkham Member, locally overlain by the thinner, mud-dominated, Bees Nest and Kenslow members (Figs 2, 3; Table 1). A Neogene age was proposed for the entire formation based on the palynology of the uppermost bed, the Kenslow Member (Boulter 1970, 1971a, b; Boulter and Chaloner 1970). The relatively sharp boundaries between the three members were interpreted by Boulter *et al.* (1971) and Walsh *et al.* (1972, 1980) as representing conformable transitions. Consequently, a single depositional sequence was envisaged with consecutive fluvial, lacustrine and paludal palaeoenvironments.

Recent palaeontological studies (e.g. Pound *et al.* 2012a) have confirmed a Miocene age for the Kenslow Member. The underlying Kirkham and Bees Nest members are palaeontologically barren (Boulter *et al.* 1971). Earlier workers, for example Howe (1897) and Kent (1957), suggested a Triassic age for these lowermost units because of their resemblance to *in situ* Triassic strata within the wider region. Based on an assessment of the available evidence described herein, the present authors support and amplify these

observations. We explain why the concept of the Brassington Formation as a genetically related Neogene succession is flawed, and demonstrate that the Kirkham and Bees Nest members correlate with local Lower–Middle Triassic lithostratigraphical units; these are the Sherwood Sandstone and Mercia Mudstone groups respectively. This contribution provides a thorough stratigraphical revision of the Brassington Formation and discusses the associated geological implications.

Geological background

Carboniferous geological setting, burial history and uplift

The area of the ‘pocket deposits’ is situated to the NE of the Midlands Microcraton on the Woo Dale Tilt Block (Smith *et al.* 2005). The Peak Limestone Group accumulated on this basement as extensive platform carbonate sediments deposited in a shallow shelf sea. They exhibit strong cyclicity that has been related to glacioeustasy and tectonic adjustments (Ford 1968; Manifold *et al.* 2020; Cózar *et al.* 2022). Platform growth ceased as it was drowned by the muds of the Middle Carboniferous Bowland Shale Formation, still evident around the platform margin (Gutteridge 2024) and by implication capping it (Ford 1999; Walsh *et al.* 2018). Platform-marginal, fault-guided subsidence formed troughs such as the Widmerpool Gulf to the south (Hennissen *et al.* 2017, fig. 1). During the Late Carboniferous the Bonsall Fault Terrace (Smith *et al.* 2005) became a focus for Variscan inversion, rifting and relative uplift on the SW side of the platform. Rifting was a focus for




Fig. 1. The geology of the central and southern Peak District, central England, illustrating major bedrock outcrops and the distribution of the 'pocket deposits'. Individual members of the former Brassington Formation are too small in area to be discernible at this scale. Note the concentration of the 'pockets' in the dolomitized limestone, and the NW–SE orientation of the two major clusters in Derbyshire. The western cluster is located south of Waterhouses, Staffordshire. Localities where pebbles characteristic of the Sherwood Sandstone Group occur in the topsoil are also indicated. Note the northwards overstep of the Triassic Sherwood Sandstone Group onto the Carboniferous strata immediately north of ~53° N. Source: British Geological Survey. DiGMapGB-50 (SHAPE geospatial data), scale 1:50 000, Tiles: ew111, ew112, ew124, ew125, updated: 30 November 2016, BGS, using: EDINA Geology Digimap Service, <https://digimap.edina.ac.uk>, downloaded: 26 April 2024 16:00:21.688. OS Open Roads (SHAPE geospatial data), scale 1:25 000, Tiles: sk, updated: 9 October 2023, Ordnance Survey (GB), using: EDINA Digimap Ordnance Survey Service, <https://digimap.edina.ac.uk>, downloaded: 26 April 2024 16:15:07.883. Contains OS data © Crown copyright and database right (2025) and British Geological Survey materials © UKRI (2025).




Fig. 2. A graphic log of the type section of the Brassington Formation of Boulter *et al.* (1971) at Bees Nest Pit, near Brassington with a key ((a) and (b) respectively), and a geological map of Bees Nest Pit (c). (a) A composite type section compiled from four successions measured and logged on the east side of the pit during the early 1970s by Ijtaba (1973, p. 17, fig. 3). The location of this composite section is indicated in (c) by four solid gold/yellow lines interpreted from information in Ijtaba (1973). Twenty-four numbered beds were recognized (Supplementary Material table 4). The ventifact-bearing interval (beds 12–16) is indicated. (b) Lithological key. (c) Sketch map of Bees Nest Pit during the early 1970s when quarrying operations were ongoing, and there was virtually full exposure of the different lithological units. Abbreviations: BNM, Bees Nest Member; BSF, Bowland Shale Formation; B. Fm., Brassington Formation; KM, Kenslow Member; MMG, Mercia Mudstone Group; P, plant fossils. Springer Nature are thanked for granting permission under licence number 6153741144237 to reuse graphics from figure 1 of Boulter *et al.* (1971) ‘Brassington Formation: a newly recognized Tertiary Formation in the Southern Pennines’. *Nature Physical Science*, 231, 134–136 (<https://www.nature.com/articles/physci231134a0>). Source: (a) and (b) are substantially modified from Boulter *et al.* (1971, fig. 1) and Ijtaba (1973, fig. 3), and (c) is considerably amended after Walsh *et al.* (1972, fig. 2A).

early dolomitization of the platform carbonates (Breislin *et al.* 2020, 2023) and probably provided the structural guidance for the ‘pocket deposits’ (Ford and King 1968; Walsh *et al.* 2018). The UK-wide uplift at the Carboniferous–Permian transition (~305–290 Ma) exposed strata of this age to the erosion event that resulted in a significant sub-Triassic unconformity (Fig. 4).

Despite scant physical evidence, there appears to have been substantial Mesozoic cover over northern England; Huddart (2002) estimated this to be 1–3 km thick. Studies of the burial history and thermal maturation of the Bowland Shale Formation in the East Midlands (e.g. Andrews 2013; Lodhia *et al.* 2023) include one at Carsington, ~4 km SE of Brassington. This suggested 2200 m of post-Permian strata (Hao *et al.* 2021).

Triassic stratigraphy and sedimentation adjacent to the Peak District

Lithostratigraphy of the Sherwood Sandstone and Mercia Mudstone groups in the vicinity of the ‘pocket deposits’

In the areas surrounding the Peak District, the Triassic is represented by the Sherwood Sandstone and Mercia Mudstone groups (Fig. 1; Wills 1970; Warrington *et al.* 1980). The former is an extensive

varicoloured sandstone unit that is locally pebbly and fine-grained; the lowermost part is conglomeratic (Ambrose *et al.* 2014). This rock is friable because of weak cementation due to weathering (Steel and Thompson 1983). The pebbles are dominated by smooth, well-rounded quartz and quartzite clasts derived from the Armorican Massif in NW France and the UK (Smith and Edwards 1991; Radley and Coram 2016; Burgess *et al.* 2024, fig. 7). Some of the pebbles exhibit faceting and/or pressure-pitting (Supplementary Material figs 1, 2; Warrington *et al.* 1980).

Within the Sherwood Sandstone Group, the Chester and Helsby Sandstone formations are most relevant to this contribution. The Chester Formation (formerly the Nottingham Castle Sandstone Formation) is Early Triassic (Olenekian) in age, and generally fines upwards. Castle Rock, a reference section in central Nottingham (UK National Grid Reference (NGR) SK 56927 39397), exposes up to 35 m of red–brown or grey, medium- to coarse-grained, pebbly, cross-bedded, fluvial/alluvial sandstone of this unit (Howard *et al.* 2009). The Helsby Sandstone Formation is of Middle Triassic (Anisian) age, and was previously mapped as the Hollington Formation in the Ashbourne area (Chisholm *et al.* 1988). This unit is lithologically and genetically comparable with the Chester Formation. The Helsby Sandstone Formation comprises beds of red, brown and grey pebbly sandstones with conglomeratic bases,




Fig. 3. The main part of the type section of the Brassington Formation of Boulter *et al.* (1971) at Bees Nest Pit, near Brassington. This is the northernmost solid yellow line in Figure 2c. The central northern face is pictured, looking northeastwards. The majority of the exposure is the Kirkham Member, now the Sherwood Sandstone Group, marked SSG. This is overlain by the Bees Nest Member, now the Mercia Mudstone Group (marked MMG), which dips steeply towards the left (i.e. northwestwards). The uppermost unit is the relatively unconsolidated Kenslow Member, marked KM. This unit tends to be washed downslope as is evident above the person on the left. Source: photograph by Peter T. Walsh taken during the mid-1970s, and reproduced with permission.

interbedded with red and brown siltstones, and mudstones (Newell 2018). This unit is reported to thin over areas of contemporary high ground including Charnwood Forest, north Leicestershire (Scotney *et al.* 2012) and the White Peak (Newell 2018, fig. 15).

The Middle to Late Triassic (Anisian–Rhaetian) Mercia Mudstone Group is a thick, geographically extensive succession that dominantly comprises brown, red and green mudstones and siltstones. Of particular relevance here is the Tarporley Siltstone Formation (formerly the Sneinton Formation) of Anisian age, which comprises interbedded laminated and varicoloured micaceous mudstones, siltstones and fine-grained sandstones. It outcrops to the south of Brassington around Burton, Derby and Nottingham (Wills 1970; Carney *et al.* 2001). The Sherwood Sandstone Group–Tarporley Siltstone Formation transition is diachronous, and is demarcated where mudstone/siltstone is dominant over sandstone (Warrington *et al.* 1980; British Geological Survey 2025). Swinnerton (1935) observed that this transition is characterized by ventifacts.

Outcrop distribution

Permian strata thin westwards from the East Midlands Shelf, and consequently the Sherwood Sandstone Group rests on a buried ridge of Lower Carboniferous strata near Ashbourne and immediately SW

of Derby (Newell 2018, fig. 15). The distribution of Triassic outliers reflects both the sub-Triassic topography and subsequent erosion histories. Outliers of the Chester Formation occur SE of Brassington (Fig. 1; British Geological Survey 1983). Relatively complete Triassic successions are present in the Cheshire Basin to the west, the Needwood Basin to the south and the East Midlands Shelf to the east (Warrington *et al.* 1980). The sub-Triassic unconformity is recognized at Nottingham where Lower Triassic rocks overlie Upper Carboniferous (upper Bashkirian to lower Moscovian) strata. Westwards, the Sherwood Sandstone Group overlies older (Bashkirian/Langsettian) Carboniferous rocks. This westerly trend indicates that Late Carboniferous uplift and Permian–Triassic denudation were greater in the west (Fig. 4).

Triassic palaeogeography and depositional environments

Sherwood Sandstone Group facies represent continental aeolian and fluvial sedimentation (Medici *et al.* 2015). The regional palaeo-current was towards the NW (Ambrose *et al.* 2014), although a mean direction of NW–SE was reported for Styrrup Quarry, Nottinghamshire (Wakefield *et al.* 2015). The Mercia Mudstone Group largely represents arid continental conditions, with intermittent marine-influenced playa lacustrine settings (Warrington *et al.* 1980; Wilson 1993; Howard *et al.* 2008; Jones *et al.* 2025). The top of the Sherwood Sandstone Group was envisaged as a wind-deflated desert pavement that was subsequently buried by the Mercia Mudstone Group. Swinnerton (1935) interpreted the Sherwood Sandstone Group–Mercia Mudstone Group transition as a sustained hiatus; this has been construed as a manifestation of the Hardegsen unconformity of the central European and North Sea basins (Smith and Warrington 1971). At Budleigh Salterton in Devon, this hiatus is associated with a clay-rich horizon, interpreted as a ‘reg’ palaeosol formed by prolonged pedogenesis under arid conditions (Wright *et al.* 1991).

Prior to the Hardegsen unconformity, deposition of the Chester Formation is attributed to the Budleighensis river system, which drained periodically towards the NW into the Staffordshire, Cheshire and Irish Sea basins, or NE towards the East Midlands Shelf of the Southern North Sea (endorheic) Basin (Aitkenhead *et al.* 2002). This bifurcation resulted from asynchronous avulsion events. NE drainage lay between the southern flank of the Pennines and the northern spur of the Charnwood Massif, and thence onto the East Midlands Shelf and into the Southern North Sea Basin (Warrington and Ivey-Cook 1992; Newell 2018). Although overlapped by the Mercia Mudstone Group during the Middle to Late Triassic, the Budleighensis river never overflowed the Charnwood Massif (Carney *et al.* 2001). Evidence from the outliers of the Chester Formation SE of Brassington (Fig. 1; Chisholm *et al.* 1988) indicates that the Sherwood Sandstone Group spread north onto the southern Pennines NW of Derby. To the south, post-Hardegsen unconformity deposition of the Helsby Formation was partly fault-controlled (Carney *et al.* 2001). Deposition was consistently north and NW through the Needwood, Stafford, Cheshire and Irish Sea basins, spilling onto the Pennines around Ashbourne (Chisholm *et al.* 1988), where fluvial sediments intercalated with coeval aeolian sands (Newell 2018). Contemporaneously, the East Midlands Shelf experienced a prolonged period of aeolian deflation

Table 1. The stratigraphy of the Brassington Formation as defined by Boulter *et al.* (1971) in the left-hand column, directly compared with the revised stratigraphical interpretations herein in the centre and right-hand columns

Stratigraphy of Boulter <i>et al.</i> (1971)	Reinterpreted lithostratigraphy and chronostratigraphy herein	
Not recognized	Frیدن Member (Brassington Formation)	Late Miocene
Kenslow Member (Brassington Formation – Neogene)	Kenslow Member (Brassington Formation)	Middle–Late Miocene
Bees Nest Member (Brassington Formation – Neogene)	Mercia Mudstone Group (Tarporley Mudstone Formation)	Middle Triassic
Kirkham Member (Brassington Formation – Neogene)	Sherwood Sandstone Group (Chester and Helsby Sandstone formations)	Early–Middle Triassic




Fig. 4. A cross-section illustrating the Carboniferous Peak Limestone, Craven and Millstone Grit groups, and the Triassic Sherwood Sandstone Group (stippled) from Turnditch in the SE, to west of Wirksworth in the NW, a distance of ~10 km (see Fig. 1), to illustrate the sub-Triassic unconformity. The projection of this unconformity to the Peak District in the north clearly demonstrates that the occurrence of the Sherwood Sandstone and Mercia Mudstone groups in the Peak District is highly likely in gross geological terms. Specifically, assuming no major fault movement, if the gradient of the base of the Triassic is projected northwards to Wirksworth from the *in situ* outcrop at Turnditch, the Sherwood Sandstone Group can be placed directly on top of the Lower Carboniferous Peak Limestone Group at, or immediately above, the contemporary land surface. This places the Sherwood Sandstone Group onto the Peak District upland surface at about 300 m above ordnance datum. The projected sub-Triassic unconformity in the NW indicates the probable presence of the Bowland Shale Formation overlying the Peak Limestone Group to source the foundered shale blocks in some of the larger ‘pockets’. For orientation, the village of Turnditch is ~12 km east of Ashbourne at NGR SK 293 463, and Yokecliffe Rake is a mineralized Mississippi Valley Type mineralized fault trending east–west to the west of Wirksworth centred on NGR SK 2679 5400. Abbreviation: MMG, Millstone Grit Group. Source: considerably modified after Frost and Smart (1979, fig. 56).

and ventifact concentration (Warrington and Ivimey-Cook 1992) and the Chester Formation is overlain by the westwards-onlapping Tarporley Siltstone Formation with a deflation unconformity at the boundary (Howard *et al.* 2009).

Post-Triassic structural evolution

Jurassic extensional tectonics episodically extended across the East Midlands (Smith *et al.* 2005), potentially contributing to the evolution of the sinkholes hosting the ‘pocket deposits’. Regional Mesozoic subsidence accommodated any post-Permian strata. Late Cretaceous epicontinental seas were relatively shallow, starved of terrigenous sediment and associated with the onset of a prolonged period of uplift that continued to the Paleocene (King 2006; Gale and Lovell 2018; Püttmann and Mutterlose 2021). This regionally discrete uplift reflected differences in rheological crustal properties (Green *et al.* 2001), possibly resulting from the development of the Iceland Plume and the onset of seafloor spreading between Europe and Greenland, coeval with the Alpine orogeny. Although Green *et al.* (2001) noted that this pre-dated their apatite fission track uplift dates, Smith *et al.* (2005) proposed regional basin inversion. Westaway (2009, 2020) postulated a lower crustal flow model supported by detailed reviews of the geomorphological indicators of lithospheric buoyancy in central and northern England. This accounts for proposed uplift of ~300–400 m since the Miocene, and by >100 m since the Middle Pleistocene.

The Cenozoic

Superimposed on the uplift history, the warm and wet climatic conditions of the Paleocene and Eocene (e.g. Kender *et al.* 2012) probably facilitated ongoing weathering of exposed rock. There is no evidence of strata of this age in the Pennines. Isolated, terrestrially derived Miocene sediments recorded in the Brassington–Friden area, Anglesey and SW England were described as palaeogeographical outliers by King *et al.* (2016), in stark contrast to the extensive coeval successions offshore (Cope *et al.* 1992). Post-Miocene erosion precludes determination of the full thickness of the Miocene in the White Peak. Accommodation space may have been limited to paludal depocentres within subsidence hollows.

Glacial deposits are scarce in the White Peak where the bedrock comprises limestones. The area was probably subject to direct action of southerly moving glaciers during the main expansion of Marine Isotope Stage 12 (MIS12) ice and earlier (Lee *et al.* 2011). By contrast, during MIS2–4, ice was limited to the perimeter of the Peak District and periglacial conditions prevailed. Surviving glacial

deposits, including till, and rafted sediments are preserved in the ‘pockets’ (Jones *et al.* 2016).

The Brassington Formation

As proposed by Boulter *et al.* (1971), the Brassington Formation was a sand- and clay-dominated succession, entirely of Neogene age. Its type section is at Bees Nest Pit, ~1 km east of Brassington village (Figs 2, 3; Supplementary Material fig. 3). The Brassington Formation was subdivided into the Kirkham, Bees Nest and Kenslow members by Boulter *et al.* (1971) (Table 1). Only the Kenslow Member has been biostratigraphically dated (Boulter 1970, 1971a, b; Boulter and Chaloner 1970; Pound *et al.* 2012a; Pound and Riding 2016). Formalization of the Brassington Formation followed on from the work of Yorke (1954, 1960, 1961) and Ford and King (1968, 1969), and was referenced in the maps and sheet memoirs produced by the British Geological Survey (Frost and Smart 1979; Aitkenhead *et al.* 1985; Chisholm *et al.* 1988). Further contributions on this unit include Kent (1957), Ford (1972), Wilson (1979) and Walsh *et al.* (1972, 1980, 2018).

The sinkholes preserving these sediments are steep-sided, of differing shapes and volumes, and largely in the dolomitized Bee Low Limestone Formation (Asbian). They occur in three clusters, mainly at elevations of between 320 and 275 m. These are centred around Friden and Brassington in Derbyshire, an area ~24 km long and ~6 km wide trending WNW–ESE, and a smaller area in the Weaver Hills, south of Waterhouses, Staffordshire (Fig. 1; Walsh *et al.* 2018, figs 1, 2). The Friden and Brassington ‘pockets’ appear to be related to the alignment of fault-controlled dolomitization (Fig. 1; Breislin *et al.* 2023). Many of the ‘pockets’ have not been mapped in detail, and many others undoubtedly remain undiscovered (Jones and Banks 2014; Riding 2021, fig. 13) as indicated by local occurrences of pebbly soils, topographical hollows, and areas of broom and gorse bushes (Supplementary Material fig. 1).

Ford and King (1968, 1969) and Walsh *et al.* (1972, 1980, fig. 15) postulated that the sand-dominated Kirkham Member was derived from the Neogene erosion of southward-retreating scarps of the Sherwood Sandstone Group. These sediments supposedly formed a decametre-scale alluvial/fluviial sand sheet (Supplementary Material fig. 4). The overlying Bees Nest and Kenslow members are, by contrast, metre-scale clay-dominated units, interpreted respectively as an alluvial siliciclastic sheet and lacustrine/paludal clays deposited locally in small hollows. Commonly, sediments in the ‘pockets’ are underlain by an insoluble chert–clay residue derived from the host limestone. Blocks of dolomitized limestone derived from roof-collapse and subsided

Table 2. A selective summary of the age interpretations of the fills of the ‘pocket deposits’, and the evidential basis for these

Author(s) and year	Age of the ‘pocket deposit’ fills	Evidence
Brown (1867)	Carboniferous or Triassic	Lithological similarity
Brodie (1886)	Triassic or Paleogene	Lithological similarity
Howe (1897)	Triassic	Lithological similarity
Thomas <i>et al.</i> (1920)	post-Triassic to Pleistocene	Overall lithofacies
Scott (1927)	Carboniferous	Lithological similarity
Kent (1957)	Triassic	Lithological similarity
Chaloner (1961)	Neogene	Palaeobotany and palynology*
This paper	Triassic and Miocene	See main text

The early works of Brown (1867), Brodie (1886) and Thomas *et al.* (1920) gave somewhat equivocal age assessments. By contrast, Howe (1897), Scott (1927) and Chaloner (1961) (in bold font) postulated Triassic, Carboniferous and Neogene ages respectively. The latter three studies were the first to indicate those ages, and therefore represent the principal breakthroughs on this topic. The title of Kent (1957) perceptively referred to the ‘pocket deposits’ as ‘Triassic relics’. *Note that the work of Chaloner (1961) is based on the Kenslow Member only.

autochthonous masses of the Bowland Shale Formation are also present (Ford and King 1969; Walsh *et al.* 2018). Furthermore, many of the ‘pockets’ are capped by Pleistocene till and/or head (Supplementary Material table 1). Brecciated mineral accumulations, typically baryte slab breccias, sporadically occur in some of the ‘pockets’ (Supplementary Material fig. 5). Reported examples include Golconda Mine ~2 km NE of Brassington (NGR SK 2488 5517, Fig. 1) (Ford and Jones 2007, pl. 9). Fragments of baryte and galena are also present in some underground alluvial red sands (Worley 1978; Ford and Jones 2007, pl. 8; Ford and Worley 2016). This material represents glacial meltwater flows and is not included in Supplementary Material table 1.

Typically, the Brassington Formation forms sag-synclinal structures with localized small- and moderate-scale faulting and steep dips at the margins, especially of the larger ‘pockets’ (Fig. 3; Walsh *et al.* 2018, figs 9, 12). A large mass of vertically dipping Bees Nest Member at Bees Nest Pit is depicted in Supplementary Material fig. 6; this faulted block indicates significant movement along a subvertical linear fault. Minor (metre-scale) faulting locally affects both the Kirkham and Bees Nest members at Bees Nest Pit (e.g. Ijtaba 1973, pl. 2) with local fracturing of cemented parts of the Kirkham Member. These structural features may represent the reactivation of pre-Triassic faults in addition to the subsidence/collapse process.

The age controversy

The chronostratigraphy of the Brassington Formation has proved contentious, with proposed ages ranging from Carboniferous to Pleistocene (Table 2). Early workers such as Brown (1867) invoked hybrid ‘middle’ Carboniferous and Triassic ages, while Brodie (1886) suggested a possible Paleogene age. Later researchers including Howe (1897), Arnold-Bemrose (1910), Barke *et al.* (1920) and Fearnside (1932) favoured a Triassic age based on lithological similarities. Yorke (1954, 1960, 1961), Edwards and Trotter (1954) and other geological surveyors, and Kent (1957) also supported a Triassic age. A more detailed commentary on this topic is provided as Supplementary Material appendix 1.

Chaloner (1961) reported Neogene plant macrofossils and pollen from the uppermost clays of the Brassington Formation. Ford and King (1968, 1969) concluded that the Kirkham Member comprised eroded Triassic sand deposited during the Neogene. Consequently, Boulter *et al.* (1971) established the Brassington Formation as entirely of Neogene age based on palynological data from the Kenslow Member. Recent research has refined the age of the Kenslow Member as Middle–Late Miocene (Serravalian–Tortonian) (Pound *et al.* 2012a, 2019; Pound and Riding 2016). The identification of a major unconformity below the Kenslow Member has contributed to a more objective chronostratigraphical interpretation (Supplementary Material fig. 7).

Evidence for the Triassic age of the Kirkham and Bees Nest members

This section documents nine lines of geological evidence pertaining to the Triassic age of the Kirkham and Bees Nest members of the Brassington Formation of Boulter *et al.* (1971); these are listed in Supplementary Material table 2.

The sub-Triassic unconformity

A major stratigraphical break immediately south of the Peak District brings Triassic strata into contact with a range of Carboniferous lithostratigraphical units (Fig. 4). NE of Derby, the Triassic Sherwood Sandstone Group overlies the Millstone Grit Group and the Pennine Lower Coal Measures Formation (uppermost Mississippian and Pennsylvanian). Westwards, between Derby and Ashbourne, Early Triassic strata rest on the Mississippian Widmerpool and Bowland Shale formations (British Geological Survey 2014). Further west, on the eastern edge of the Weaver Hills, the Sherwood Sandstone Group is in direct contact with the Mississippian Peak Limestone Group (British Geological Survey 1983). A ‘pocket deposit’ at Sallymoor (SK 083 464) includes sediment closely resembling local Triassic bedrock and blocks of red and yellow pebbly sand/sandstone of Triassic aspect that are typical of several ‘pockets’ within the western cluster (Chisholm *et al.* 1988) (Fig. 1; Supplementary Material table 3).

If projected northwestwards from Derby, the base of the Sherwood Sandstone Group would lie immediately above the eroded surface of the Peak Limestone Group near Wirksworth and Brassington (Fig. 4). This suggests that the Peak Limestone Group was formerly overlain by Triassic strata within what is now the southern cluster of ‘pocket deposits’ (Figs 1, 4). Detached masses of the Bowland Shale Formation in the ‘pockets’ imply that a thin layer of this unit was present above the Peak Limestone Group between the Late Carboniferous and the earliest Triassic (Walsh *et al.* 1972, 2018). Secondary staining (‘reddening’) of Carboniferous strata is known at numerous localities south of the Peak District (Fig. 4; Frost and Smart 1979; Chisholm *et al.* 1988). The staining affects mudstones and sandstones immediately below the sub-Triassic unconformity, including the shale masses within the ‘pockets’. Such secondary effects are attributed to weathering at the sub-Triassic surface, or to downward percolation of iron-bearing solutions from a former Triassic cover (Stevenson and Gaunt 1971). The juxtaposition of Triassic strata above Lower Carboniferous limestones and mudstones accords with the lithostratigraphical sequences within the larger ‘pockets’ (Supplementary Material table 1).

Lithological characteristics of the Kirkham Member

The Kirkham Member of Boulter *et al.* (1971) has been recorded in all the known ‘pockets’, and is mostly an unlithified cream/white,

medium- to coarse-grained sand (Fig. 5). It is frequently pebbly and contains occasional green and red clay/silt intraclasts and stringers. Locally, the Kirkham Member is weakly to relatively well-lithified (Supplementary Material fig. 8). The vertical thickness is variable because this parameter is controlled primarily by the depth of each 'pocket'. For example, Boulter *et al.* (1971) and Ijtaba (1973) recorded 35.31 m at the type section at Bees Nest Pit (Supplementary Material table 4). Walsh *et al.* (2018, table 1) mentioned thicknesses of ~40–50 m elsewhere.

Masses of pebbly medium- to coarse-grained coherent sandstone are generally yellow, red–brown and grey in colour, whereas weakly lithified and unconsolidated sand with well-rounded quartzite and quartz pebbles is largely cream/white in colour but locally is red to red/brown (Supplementary Material fig. 9; Aitkenhead *et al.* 1985). The cream/white deposits appear to have been bleached, perhaps by hypogene fluids and/or exposure to strong sunlight. Typically, the grains are coated by thin pellicles of kaolinite which are indicative of intense alteration, presumably of feldspar (Walsh *et al.* 1972, 1980, 2018). Together with sustained weathering, this may also account for the weak or absent cementation. The silica and kaolinite content, and largely uncemented characteristics were the principal reasons the Kirkham Member was quarried for refractory brick manufacture (Supplementary Material appendix 2; Boswell 1918). The Kirkham Member displays many features that typify the

Sherwood Sandstone Group of the English Midlands. These include the absence of fossils, colour, weak cementation, particle morphology, sedimentary structures, grain and pebble composition, lithology, overall palaeocurrent flow direction, overall sedimentology and texture. Pebbles from the Kirkham Member and the Sherwood Sandstone Group are closely comparable (Supplementary Material fig. 2).

A cemented sandstone in the Kirkham Member is exposed alongside the old access track at the west side of Bees Nest Pit, near Brassington (NGR SK 24108 54574; Supplementary Material fig. 8). This comprises 5–6 m of well- to moderately well-consolidated yellow–brown sandstone. Here, the lowermost part, a partially lithified conglomerate referred to as Bed 7 of the Kirkham Member (Ijtaba 1973, fig. 3; Walsh *et al.* 1980, fig. 2) closely resembles the pebble beds at the base of the Chester Formation.

Because the cemented sandstone masses in the Kirkham Member (Supplementary Material fig. 8) resemble the Chester and Helsby Sandstone formations, these parts of the Kirkham Member of Boulter *et al.* (1971) are referred to the Sherwood Sandstone Group. Likewise, the weakly lithified sands within the Kirkham Member are laterally contiguous with the cemented masses, and are also lithologically similar. Hence, the latter facies is also assigned to the Sherwood Sandstone Group. The contention of Ford and King (1969, p. 60) and Walsh *et al.* (1972, p. 522) that the cemented



Fig. 5. A montage of four photographs of former working quarries illustrating the unlithified facies of the Kirkham Member, now the Sherwood Sandstone Group, around Brassington and Friden taken in the early 1970s (see also Walsh *et al.* 2018, figs 5, 6, 11). These are now largely obscured, and Kirkham's Pit (c) has been infilled. (a) The NW corner of Bees Nest Pit, near Brassington, taken from the SE corner (NGR SK 24159 54491). (b) The north side of Green Clay Pit, near Brassington, taken from the southern side (NGR SK 23974 54703). (c) The SE part of Kirkham's Pit, west of Brassington, taken from the NW side (NGR SK 21729 54138). (d) The NE part of Kenslow Top Pit, near Friden, taken from near the access road looking towards the SW (NGR SK 18223 61507). Source: photographs are by Peter T. Walsh and are used with permission.

masses represent localized secondary re-cementation of Triassic erosion debris during the Miocene is considered to be highly unlikely. We propose that these are relicts of *in situ*, incompletely weathered, Sherwood Sandstone Group that have subsided into the ‘pockets’.

Lithological characteristics of the Bees Nest Member

The Bees Nest Member of Boulter *et al.* (1971) is represented by 6–9 m of lithified varicoloured, largely red and green mudstones and siltstones, and is only present in the centres of the larger ‘pockets’ (Walsh *et al.* 2018). Thicknesses of ~30 m and up to 40 m at Kenslow Top Pit reported by Ijtaba (1973, p. 25) and Aitkenhead *et al.* (1985, p. 107) respectively are considered herein to be apparent thicknesses and substantial overestimates of the true stratigraphical thickness of the steeply inclined foundered strata. Currently, the only known exposures of the Bees Nest Member are in Bees Nest Pit (Fig. 6; NGR SK 24100 54580 and NGR SK 24170 54537) and at Spencer’s Pit (NGR SK 21508 54147) ~1 km ENE and ~1.75 km west of Brassington respectively. This unit was previously exposed in other silica sand pits including Kenslow Top and Kirkham’s pits, but these sections are no longer visible. At the type section in Bees Nest Pit, the exposed Bees Nest Member



Fig. 6. The uppermost part of the Bees Nest Member, now the Mercia Mudstone Group, at Bees Nest Pit. This is the type section of the former Bees Nest Member, east of the main access ramp on the north side of the quarry. The outcrop on the right has been cleaned to show the beds of reddish-brown mudstones interbedded with green siltstones and pale-brown, fine-grained sandstones. This unit overlies the Kirkham Member/Sherwood Sandstone Group, and is overlain, with a sharp boundary, by the Kenslow Member. Source: photograph taken in September 2019 by Peter F. Jones; geological hammer for scale.

comprises 6.17 m of interbedded unfossiliferous red, brown, grey and green mudstones and siltstones, with occasional fine-grained sandstones (Figs 3, 6; Supplementary Material fig. 10). Some of the sandy beds are pebbly, and the pebbles are well-rounded quartz and quartzite, reminiscent of those in the Sherwood Sandstone Group.

The Bees Nest Member is lithologically very similar to the Middle to Late Triassic Mercia Mudstone Group. Specifically, this unit preserved at Bees Nest and Spencer’s pits closely resembles the Tarporley Siltstone Formation. Due to the occurrence of pebbles in the Bees Nest Member, this unit is reminiscent of the lowermost Tarporley Siltstone Formation of the Nottingham area, specifically the Woodthorpe Member (Howard *et al.* 2008). In summary, because the Bees Nest Member resembles the Tarporley Siltstone Formation east and south of Brassington, these two units are interpreted herein to be equivalent. This means that the representative units of the Sherwood Sandstone and Mercia Mudstone groups (i.e. the Kirkham and Bees Nest members respectively) in the ‘pocket deposits’ are thus preserved in their expected stratigraphical order.

The transition between the Kirkham and Bees Nest members at the type section

Significant parts of the Kirkham, Bees Nest and Kenslow members of Boulter *et al.* (1971) are still visible at the type section (Figs 3 and 5–7), although the exposures have deteriorated since quarrying ceased. Records of the geology at Bees Nest Pit were made when the succession was better exposed than at present (e.g. Yorke 1954, 1960, 1961; Walsh *et al.* 1972, 1980; Ijtaba 1973), and collectively comprise a valuable geological archive. A log of the type section presented by Boulter *et al.* (1971, fig. 1) is a composite of several exposures on the southern, eastern and northern sides of the pit. Each lithological unit is numbered consecutively upwards from the base of the 42.55 m thick succession (Fig. 2; Supplementary Material table 4).

Between the Kirkham and Bees Nest members, there is a transition zone of at least 3 m within which this boundary could have been placed. Whereas the Bees Nest Member largely comprises thin units of red or green silty clay, the lowermost 0.53 m is pebbly. Conversely, below Bed 12 (pebbly sand) of the underlying Kirkham Member, four thin clay/silt units (beds 8–11) are present within a predominantly sandy succession. This situation is analogous to the boundary between the Sherwood Sandstone and Mercia Mudstone groups in the Triassic successions of the East Midlands.

Wind-etched pebbles at the Kirkham Member/Bees Nest Member transition

At the Kirkham Member/Bees Nest Member transition, matrix-supported pebbles exhibit markedly etched surface textures with mesoscale flutes, grooves and pits (Fig. 8; Supplementary Material fig. 11). These are interpreted as ventifacts from wind action in unvegetated environments (Viles and Bourke 2007; Durand and Bourquin 2013). This contrasts with smooth-surfaced pebbles elsewhere in the Kirkham Member (Supplementary Material fig. 2).

Three potential explanations for the occurrence of this zone of concentrated ventifacts at the Kirkham Member/Bees Nest Member transition are:

- (i) *in situ* erosion by wind action during the Neogene;
- (ii) derivation from ventifact-bearing Triassic source rocks with redeposition during the Neogene; and
- (iii) ventifacts formed at the Triassic land surface, and subsequently buried by younger Triassic sediments.



Fig. 7. The Kenslow Member of the Brassington Formation at the type section, the central–northern area of Bees Nest Pit, near Brassington. Abundant plant fossils are present, dominated by blackened and dark brown fragments of mummified wood (Mustoe 2018). Source: photograph taken during the late 1970s by Peter F. Jones; geological hammer for scale.

Given the evidence for a forest-dominated landscape and a humid climate during the Miocene (Pound and Riding 2016; McCoy *et al.* 2022), the first possibility can be discounted. The second is also highly unlikely owing to the well-preserved wind-etched features. Because of the hot, arid climate during the Triassic, the third possibility appears to be by far the most plausible.

With the exception of Hughes (1952) and Yorke (1961), there are few recorded details of wind-worn pebbles from the Kirkham Member. Unpublished research by D. B. Thompson suggests that ventifacts are rare in the Kirkham Member. By contrast, these sculpted pebbles are common in the Triassic of the English Midlands. Examples have been reported from the Triassic close to the southern Pennines (e.g. Lamplugh *et al.* 1908; Swinnerton 1914; Thompson and Worsley 1967; Frost and Smart 1979; Howard *et al.* 2009). The distinctly wind-worn pebbles occur in a thin interval (beds 12–16 of Ijtaba 1973), straddling the contact (i.e. the base of Bed 13) between the Kirkham and Bees Nest members (Fig. 2). This zone of exclusively wind-worn pebbles indicates a relict desert pavement, perhaps reminiscent of the upper surface of the Lower Triassic Budleigh Salterton Pebble Bed at Budleigh

Salterton, Devon (Wright *et al.* 1991). Similarly, the concentration of ventifacts reported to overlie the Sherwood Sandstone Group at Sneinton, Nottingham (Supplementary material fig. 12; Swinnerton 1914; Howard *et al.* 2009), is probably an additional example of such a pavement, and is significant for being located much closer to the Brassington area. It seems probable that the relatively sharp lithostratigraphical change from the Kirkham Member to the Bees Nest Member represents the Triassic Hardeggen unconformity. Furthermore, for Triassic-derived ventifacts to be concentrated in this interval during the Neogene requires highly unusual circumstances: a source of Triassic wind-worn pebbles in the local area; erosion from the host rock; water transport with limited damage to the surface ornamentation (delicate micro-pitting in some cases); and the exclusion of Sherwood Sandstone Group pebbles without the surface textures. A control sample taken from Bed 7 of Boulter *et al.* (1971) stratigraphically lower in the Kirkham Member did not reveal sculpted pebbles. The pebbles in Bed 7 are typically smooth and well-rounded (Supplementary material fig. 2).

Material properties and engineering geology

Advances in the analysis of material properties since Walsh *et al.* (1972) and Ijtaba (1973) underpin the evidence-based comparison with more recent Triassic literature. The Kirkham Member is kaolinitic, thus matching the clay assemblage of the Sherwood Sandstone Group (Table 3; Ijtaba 1973; Bath *et al.* 1987). Similarly, the illite-rich Bees Nest Member is entirely consistent with the Mercia Mudstone Group (Table 3; Hobbs *et al.* 2002; Jones *et al.* 2025). Heavy mineral analysis shows both the Kirkham Member and the Sherwood Sandstone Group in Derbyshire are sparse in staurolite, and rich in tourmaline and zircon (Table 3; Ijtaba 1973; Jeans *et al.* 1993). Subsidence and possible contact with hypogene fluids (Banks *et al.* 2015) may have further reduced the mechanical strength of these units, as could exposure to hydrological weathering during the Quaternary, for example, resulting in the dissolution of any carbonate cement. The grading analyses of Ijtaba (1973) were also considered in the discussion of the lithostratigraphy herein. Pre-consolidation pressure tests of clays show significant differences between the Kirkham (888 kN m⁻²) and Kenslow (257 kN m⁻²) members, far exceeding stress from their current separation, suggesting considerably different burial histories and geological ages (Table 4; Ijtaba 1973).

Wilson (1979) studied sand grain surface textures from the Kirkham Member. These match the equivalent features in the



Fig. 8. A wind-etched pebble with flutes and grooves from the transition zone between the Kirkham and Bees Nest members (now the Sherwood Sandstone and Mercia Mudstone groups respectively) from the north side of Bees Nest Pit, near Brassington, and interpreted as a ventifact. Length is 70 mm, and breadth is 43 mm. This surface texture contrasts markedly with the pebbles from Bed 7, which are smooth and well-rounded (Supplementary Material fig. 2). Source: photograph taken by Peter F. Jones.

Table 3. A comparison of relevant studies on the clay mineralogy and the heavy minerals of the Brassington Formation of [Boulter *et al.* \(1971\)](#), and the Triassic Sherwood Sandstone and Mercia Mudstone groups

Mineral species	Mineralogical analyses of the Brassington Formation	Mineralogical analyses of Triassic strata	Stratigraphical interpretation
Clay minerals (analysed by X-ray diffraction)			
Gibbsite	Ijtaba (1973) studied the Kenslow Member (Bed 24), and found abundant gibbsite; this suggests humid, warm weathering of Miocene lacustrine clays during the Pliocene (Gasparini <i>et al.</i> 2022)	N/A	Miocene (Kenslow Member)
Illite	Ijtaba (1973) studied the Bees Nest Member, beds 19–21, and found this succession to be rich in illite	Hobbs <i>et al.</i> (2002) reported that the clays of Mercia Mudstone Group units A and B of NE England are illite-rich (see also Cripps and Taylor 1987 and Jones <i>et al.</i> 2025)	Triassic (Bees Nest Member)
Kaolinite	Ijtaba (1973) studied beds 2, 6 and 8 of the Kirkham Member and found them to be highly kaolinitic	Bath <i>et al.</i> (1987) investigated the diagenesis of the Sherwood Sandstone Group, and found K-feldspar and authigenic kaolinite to be present	Triassic (Kirkham Member)
Heavy mineral analyses			
Anatase	Ijtaba (1973) from the Kirkham Member	Jeans <i>et al.</i> (1993) – specific gravity >2.92; 64–250 µm	
Brookite	Scarce or absent; euhedral bluish, dusky, yellowish tabular grains	Staurolite is common in the Sherwood Sandstone Group of Nottinghamshire, but it is sparse in Derbyshire. Jeans <i>et al.</i> (1993) suggested that heavy mineral analyses of the overlying Mercia Mudstone Group would be informative; their work indicated a lower tourmaline content with locally rapid changes in garnet and sphene concentrations	
Garnet	Scarce or absent; colourless, fractured and isotropic grains		
Monazite	Scarce or absent; biaxial, light yellow rounded grains		
Rutile	Minor; red (ubiquitous) and yellow; rounded to subrounded; transverse striations on some grains		
Staurolite	Minor; straw yellow, marked pleochroism and with conchoidal surfaces on some grains		Triassic (Kirkham Member)
Titanite	Two grains observed from Bees Nest Pit		
Tourmaline	Abundant; blue, brown, green and yellow; angular to rounded; roundness increases with size; variable intensity of pleochroism		Triassic (Kirkham Member)
Zircon	Dominant; grain roundness increases with size; several grains with faint dusty zoning parallel to the crystal faces		Triassic (Kirkham Member)
Opaques	Hematite, ilmenite, leucoxene coating and magnetite		

This indicates that the Kirkham and Bees Nest Members of the Brassington Formation are likely to be of Triassic age, and that the Kenslow Member is of Miocene age.

Table 4. A summary of sand grain surface texture analyses, pebble solution marks and pre-consolidation pressure tests on material from the Brassington Formation of Boulter *et al.* (1971)

Summary details		Stratigraphical interpretation
1. Sand grain surface texture analyses		
Wilson (1979) examined nine sand samples from the Kirkham Member using a scanning electron microscope, and compared these with a sample of the Chester Formation of the Sherwood Sandstone Group from Hulland Quarry (NGR SK 280 455) 10 km south of Brassington	Wilson (1979) stated that the surface features of the grains from the 'pocket deposits' lack high-energy indicators of secondary subaqueous abrasion and high-energy chemical weathering; however, they do exhibit evidence of chemical abrasion and mechanical fracturing	Triassic (Kirkham Member)
2. Pebbles with pressure solution marks		
Some pebbles from Bed 7 of the Kirkham Member exhibit pressure pitting, which strongly suggests significant overburden pressure		Triassic (Kirkham Member)
3. Pre-consolidation pressure tests		
Ijtaba (1973) found that the pre-consolidation pressure of Bed 10 of the Kirkham Member is 829 kN m^{-2}	Ijtaba (1973) concluded that beds 10 and 24 are both normally consolidated clays, but there is a significant difference in the pre-consolidation pressures; Davis and Chandler (1973) also discussed the pre-consolidation pressures of weathered Mercia Mudstone Group material	Bed 10 (Kirkham Member) is Triassic, and Bed 24 (Kenslow Member) is Miocene
Ijtaba (1973) found that the pre-consolidation pressure of the Kenslow Member (Bed 24) is 257 kN m^{-2}		
Beds 10 and 24 are separated by 5–6 m of sediment (around 120 kN m^{-2}); the coefficients of volume compressibility (Mv) and coefficient of consolidation (Cv) values were not given by Ijtaba (1973)		

The top two rows (Part 1) summarize the work of Wilson (1979) on the surface textures of sand grains from the Kirkham Member. This indicates that these are closely comparable to the Triassic Chester Formation of the Sherwood Sandstone Group. Below this (Part 2) is the observation that the surfaces of some quartz and quartzite/metaquartzite pebbles from Bed 7 of the Kirkham Member exhibit pressure solution marks indicative of considerable post-depositional sediment loading during the Mesozoic and Paleogene. This means that the Kirkham Member is far more likely to be of Triassic, rather than Neogene, age. Part 3 documents data on pre-consolidation pressure tests of beds 10 and 24 of Boulter *et al.* (1971) and Ijtaba (1973). The differences noted are consistent with the Triassic Kirkham Member being deeply buried and consolidated below a younger Mesozoic cover, and then unroofed by erosion prior to deposition of the Miocene Kenslow Member.

Chester Formation, with no unequivocal evidence for subsequent fluvial action. This is contrary to the hypothesis of the Kirkham Member being the result of a second phase of riverine reworking of the Sherwood Sandstone Group of Walsh *et al.* (1972) (Table 4). Some of the pebbles exhibit small white pits, which appear to be pressure solution features (Supplementary material fig. 2). These pits are interpreted as being developed during post-depositional sediment loading (Table 4), suggesting that there was a significant thickness of post-deposition strata that has subsequently been removed by erosion. This supports the view that Bed 7 of the Kirkham Member represents *in situ* Triassic Sherwood Sandstone Group that subsequently subsided into dissolution cavities without further overland movement. This is because, were the pebbles derived from reworking of Triassic sandstones, the pressure solution marks would probably have been destroyed.

Palaeontology

In common with the Sherwood Sandstone Group (Holloway *et al.* 1989; Ambrose *et al.* 2014; Warrington and Pollard 2021) the Kirkham Member is apparently devoid of fossils including palynomorphs (Boulter *et al.* 1971, JBR personal observations). Similarly, the Bees Nest Member is macropalaeontologically barren, comparable with the Mercia Mudstone Group, which is extremely sparse in macrofossils (e.g. Howard *et al.* 2008; Gallois 2019). Six palynological samples analysed from the Bees Nest Member of Bees Nest Pit (Fig. 6) proved entirely devoid of palynomorphs, and contained only dark fragments of woody tissue. This depauperate nature is consistent with the Tarporley Siltstone Formation of the East Midlands (Howard *et al.* 2009, p. 113 and references therein; Hodgskiss *et al.* 2024).

Conversely, the Kenslow Member is extremely rich in macrobotanical, microbotanical and fungal fossils indicative of a Miocene age (Boulter 1971a; Pound and Riding 2016; Pound *et al.* 2019, 2022; O'Keefe *et al.* 2020; McCoy *et al.* 2022). The marked difference in the fossil contents of the Bees Nest and Kenslow members represents a depositional hiatus and disparate palaeoenvironmental conditions. If these two clay-rich units were both

Miocene in age, as hypothesized by Ford and King (1968, 1969), Boulter *et al.* (1971) and Walsh *et al.* (1972), it is unlikely that they would exhibit such a marked difference in palaeontological content. For example, if the Kirkham Member represents reworking of the Triassic Sherwood Sandstone Group during the Miocene, and redeposited in a well-vegetated alluvial/fluvial setting with a pervasive, temperate climate, then it should contain abundant plant fossils, like the Kenslow Member.

Palaeoclimate

Clay mineralogy, colour, relative lack of fossils, mud flake intraclasts and sedimentology all indicate that the Triassic Sherwood Sandstone and Mercia Mudstone groups were deposited in a predominantly arid, monsoonal palaeoclimatic regime with intense seasonality (e.g. Parrish 1993; Sellwood and Valdes 2006; Preto *et al.* 2010). Furthermore, ventifacts are present, and bioturbation and organic palaeosols are absent in the Sherwood Sandstone Group/Kirkham Member of the Brassington Formation of Boulter *et al.* (1971) (Fig. 2). These Triassic strata are interpreted as being deposited in an extreme greenhouse palaeoclimate (Radley and Coram 2016). The predominantly red colour with green/grey mottling and layering of the Bees Nest Member is consistent with a highly arid depositional setting, comparable to the Mercia Mudstone Group (Howard *et al.* 2008; Milroy *et al.* 2019).

In stark contrast, the European Miocene was relatively warm, wet and temperate, and sedimentary successions with proxies for hot and dry conditions are regionally unknown (Bruch *et al.* 2007; Pound *et al.* 2011, 2012b; Steinhilber *et al.* 2021). The Kenslow Member was deposited in a low-relief lacustrine/paludal environment under warmer and wetter conditions than present-day UK (Pound *et al.* 2012a, 2022; Pound and Riding 2016; Gibson *et al.* 2022; McCoy *et al.* 2022).

Palaeomagnetism

The palaeomagnetism of the cemented sandstone in the Kirkham Member and the Bees Nest Member at Bees Nest Pit was analysed

by Conall Mac Niocaill (University of Oxford). Several core samples were taken in 2018 from a metre-scale succession of lithified former Kirkham Member on the west side of the access track to Bees Nest Pit (Supplementary Material fig. 8; NGR SK 24108 54574), and from the type section of the Bees Nest Member at Bees Nest Pit (Figs 3, 6; NGR SK 24100 54580). The shallow palaeomagnetic inclinations recorded from these members are consistent with a Triassic age, when what is now the UK lay at tropical latitudes (Warrington and Ivimey-Cook 1992). Steeper inclinations would be expected in Miocene deposits, when Europe was close to its present temperate latitude (Tauxe 2010).

Revised lithostratigraphy and regional implications

The evidence presented above requires that the Brassington Formation of Boulter *et al.* (1971) is redefined herein. Consequently, the Kirkham and Bees Nest members are discarded as redundant lithostratigraphical units, and formally referred to the Sherwood Sandstone Group (Chester and Helsby Sandstone formations) and the Mercia Mudstone Group (Tarporley Siltstone Formation) respectively (Table 1). The Brassington Formation is emended such that the Kenslow Member of Boulter *et al.* (1971) is retained, and a new Friden Member is proposed (Appendix A and Supplementary Material appendix 3). This revision denotes a radical reduction of Peak District sites containing sediments attributable to the Brassington Formation. These are: Bees Nest Pit (Kenslow Member), Kirkham's Pit (Kenslow Member) and Kenslow Top Pit (Kenslow and Friden members). Other sites where organic deposits have been reported include Green Clay, Minninglow, Blake Moor, Heathcote and Sallymoor pits, and Hindlow Quarry. Although these are potential correlatives of the Kenslow Member, they lack detailed analysis (Supplementary Material table 3).

The presence in the 'pockets' of subsided masses of Triassic sediment has significant implications. For example, it indicates that whilst there is no representation of Upper Carboniferous, Permian and lowermost Triassic strata, the Sherwood Sandstone Group formerly extended northwards at least as far as the Brassington–Friden area. This implies the existence of a more complex and laterally extensive fluvial palaeoenvironment during the Early–Middle Triassic. It also questions traditional views that the northward-flowing 'Budleighensis' river system was deflected northwestwards and northeastwards by higher ground on reaching the southern Pennines (e.g. Radley and Coram 2016). Although limited in number, the Mercia Mudstone Group occurrences in the 'pockets' are among the first to be recorded in the southern Pennines. Triassic successions within the Peak District are less complete than in the adjacent basins. However, they offer the potential to add to existing information on palaeoenvironments and to knowledge of post-Carboniferous history, with associated implications for the interpretation of karst processes, and the exhumation history of the carbonate platform and its hydrological evolution.

Miocene non-marine sediments are rare in Britain (King 2006). This has been attributed to a hiatus in deposition during uplift associated with the Alpine orogeny (Jackson 2008). Although the depositional context requires further analysis, the freshwater Brassington Formation, as revised herein, possibly accumulated in subsidence-related depressions reactivated by Alpine uplift and associated faulting (Andrews 2013). It was suggested by Walsh *et al.* (1972, 2018) that these sediments formed close to sea-level. If future palynological studies can establish the precise altitude of deposition, the potential exists to substantially refine our understanding of Neogene uplift rates for the Peak District. This would have additional applications for interpretations of the landscape and drainage evolution of upland Britain. Therefore, despite its limited

occurrence, the emended Brassington Formation is a particularly significant onshore Miocene deposit in the UK, reinforcing the views expressed by Pound and Riding (2015).

Conclusions

This contribution fundamentally revises the stratigraphy of the 'pocket deposits' in the southern Pennines. All three members of the Brassington Formation of Boulter *et al.* (1971) were originally interpreted as being of Neogene age. The revision assigns the lower two units to the Triassic. The Kirkham Member comprises unfossiliferous, weathered pebbly sandstone. Based on exposure analogues, absence of fossils, geotechnical properties, lithofacies, mineralogy, palaeomagnetism and sedimentology, this succession is referred to the Early–Middle Triassic Sherwood Sandstone Group, as several authors previously postulated. The overlying Bees Nest Member comprises unfossiliferous, varicoloured mudstones and siltstones. Evidence from clay mineralogy, lithology, palaeomagnetism and the transitional boundary indicates correlation with the Middle–Late Triassic Mercia Mudstone Group. This stratigraphical juxtaposition is entirely consistent with Triassic successions throughout southern Britain (Howard *et al.* 2008; Ambrose *et al.* 2014; Newell 2018). Only the uppermost Kenslow Member and newly defined Friden Member (see Appendix A) yield Miocene index fossils, and now constitute the revised Brassington Formation. Therefore, we contend that the concept of an entirely Neogene succession of Boulter *et al.* (1971) lacks empirical proof. Specifically, there is no *prima facie* evidence for a Neogene age for the Kirkham and Bees Nest members. Their gross lithofacial nature clearly indicates the preservation of a classic Triassic stratal succession.

This occurrence of the Sherwood Sandstone Group at the southern end of the Pennines adds to our understanding of Triassic palaeogeography and the fluctuating nature of the 'Budleighensis' fluvial system. Together with the first reported occurrence of the Mercia Mudstone Group in the White Peak, the stratigraphical revisions herein highlight that a more extensive Triassic cover was formerly present at the southern end of the Pennines, in addition to reaffirming the negligible presence of Miocene rocks throughout onshore UK. Nevertheless, more detailed analyses of the revised Miocene Brassington Formation potentially offer significant insights with respect to the complex uplift history of the Pennines.

In summary, the evidence presented herein overwhelmingly favours a Triassic, rather than Miocene, age for the Kirkham and Bees Nest members of the Brassington Formation of Boulter *et al.* (1971), indicating relatively minor northerly palaeogeographical extensions of two of the most distinctive and extensive lithostratigraphical units in northern Europe.

Scientific editing by Neil Davies

Acknowledgements The constructive comments of two anonymous reviewers and Andy Farrant (BGS) have substantially improved this contribution. We thank the late Peter T. Walsh for his generous sharing of knowledge and data. For the purposes of open access, the authors have applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission. VJB and JBR publish with the approval of the Director, British Geological Survey (NERC).

Author contributions JBR: investigation (lead), project administration (lead), writing – original draft (lead), writing – review & editing (lead); VJB: conceptualization (equal), formal analysis (equal), investigation (equal), writing – original draft (supporting), writing – review & editing (equal); PFJ: conceptualization (equal), formal analysis (equal), investigation (equal), writing – original draft (supporting), writing – review & editing (supporting); MJP: conceptualization (supporting), investigation (equal), writing – original draft (supporting), writing – review & editing (equal)

Funding This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Competing interests The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper

Data availability All data generated or analysed during this study are included in this published article (and if present, its [supplementary information files](#)).

Appendix A: Revised lithostratigraphy of the Brassington Formation

As outlined in the main text, this contribution fundamentally revises the Miocene Brassington Formation of [Boulter *et al.* \(1971\)](#) (Table 1). Specifically, the Kirkham and Bees Nest members of these authors are discarded as redundant lithostratigraphical units, which are referred to the Triassic Sherwood Sandstone Group and the Mercia Mudstone Group respectively. The Brassington Formation is emended herein such that the Kenslow Member of [Boulter *et al.* \(1971\)](#) is retained, and the overlying Friden Member is newly proposed. The full lithostratigraphical revision is presented in appendix 3 of the [Supplementary Material](#). However, because the Friden Member is a lithostratigraphical novelty, this unit is formally outlined below.

The Friden Member of the emended Brassington Formation (new unit)

Overview: The Friden Member is formally established herein. It is a clay- and sand-dominated unit found only at the southern side of Kenslow Top Pit, and comprises a large glaciotectionic raft ~3 m thick within subglacial sediments (Fig. 9). It was originally described by [Jones *et al.* \(2016\)](#), and is their units B and C ([Supplementary Material](#) figs 13, 14 respectively). [Pound *et al.* \(2012a\)](#) recovered a Late Miocene (Tortonian) pollen flora from the Friden Member, which they misidentified as the Kenslow Member. The Friden Member is younger than the Kenslow Member based upon mapping at Kenslow Top Pit, the lithological dissimilarity to the Middle–Late Miocene Kenslow Member at the type section and pollen biostratigraphy ([Pound *et al.* 2012a](#)). It is therefore younger than the Kenslow Member, which is Middle–Late Miocene in age ([Pound and Riding 2016](#)), and is distinguished from the latter unit on clear lithological grounds.

Type section: The southern side of Kenslow Top Pit, near Friden, Derbyshire, UK (Fig. 9; NGR SK 18289 61420). There are no other exposures of the Friden Member, and hence a separate reference section cannot be designated.

Lithology: Subglacially sheared, interbedded brown and grey clay, yellow–brown sand, and brown gravels. The beds are streaky and locally deformed due to cryoturbation/glaciotectionics, and there is local hydrofracturing ([Jones *et al.* 2016](#), figs 3–6).

Upper boundary: The top of the Friden Member is defined by its upper boundary with Quaternary glaciogenic sediments, interpreted as subglacial fluvial deposits (unit D of [Jones *et al.* 2016](#)) at Kenslow Top Pit.

Lower boundary: The base of the Friden Member is defined by the boundary between units B and A of [Jones *et al.* \(2016\)](#); unit A was interpreted as locally derived subglacial debris. The transition with the underlying Kenslow Member has therefore not been observed.

Distribution and thickness: The Friden Member has only been recorded from the type section at the southern end of Kenslow Top Pit; it is up to ~3 m in thickness.



Fig. 9. The type section and only known outcrop of the Friden Member of the Brassington Formation at the south side of Kenslow Top Pit, near Friden (Appendix A and [Supplementary Material appendix 3](#)). This was originally described by [Jones *et al.* \(2016\)](#) as units B and C. The base of the Friden Member is the base of the prominent grey clay (unit B) marked by the base of the cleaned section at the break of slope immediately above the field notebook. The top of the Friden Member is the upper boundary of Unit C, the streaky succession of interbedded brown clays and yellow sands; the geological hammer is in the centre of this unit. Unit B is underlain by locally derived subglacial debris, and Unit C is overlain by lighter brown clays with gravels (head). Photograph taken in June 2015 by Peter F. Jones; geological hammer and spade for scale.

Genetic interpretation: The brown and grey clays are interpreted as lacustrine deposits. Periodically, substantial influxes of sand and gravel were transported into the lake, perhaps due to storm activity.

Age: Late Miocene (Tortonian) ([Pound *et al.* 2012a](#)).

References

- Aitkenhead, N., Chisholm, J.I. and Stevenson, I.P. 1985. *Geology of the Country Around Buxton, Leek and Bakewell. Memoir of the British Geological Survey*. Her Majesty's Stationery Office, London.
- Aitkenhead, N., Barclay, W.J., Brandon, A., Chadwick, R.A., Chisholm, J.I., Cooper, A.H. and Johnson, E.W. 2002. *British Regional Geology: The Pennines and Adjacent Areas*. 4th edn. British Geological Survey, Nottingham.
- Ambrose, K., Hough, E., Smith, N.J.P. and Warrington, G. 2014. *Lithostratigraphy of the Sherwood Sandstone Group of England, Wales and South-West Scotland*. British Geological Survey Research Report, **RR/14/01**. British Geological Survey, Nottingham.
- Andrews, I.J. 2013. *The Carboniferous Bowland Shale Gas Study: Geology and Resource Estimation*. British Geological Survey for Department of Energy and Climate Change, London, UK.
- Arnold-Bemrose, H.H. 1910. Chapter XXIII. The Lower Carboniferous rocks of Derbyshire. *Proceedings of the Geologists' Association*, **1–20**, 540–563, [https://doi.org/10.1016/S0016-7878\(10\)80005-7](https://doi.org/10.1016/S0016-7878(10)80005-7)
- Banks, V.J., Jones, P.F. and Raines, M.G. 2015. Karst geohazards associated with a carbonate platform edge. XVI European Conference on Soil Mechanics and Geotechnical Engineering, September 2015, Edinburgh, *Proceedings Volume*, 1–6.
- Barke, F., Hind, W. and Scott, A. 1920. Quartzose conglomerate at Caldon Low, Staffordshire. *Geological Magazine*, **57**, 76–82, <https://doi.org/10.1017/S0016756800100986>
- Bath, A.H., Milodowski, A.E. and Strong, G.E. 1987. Fluid flow and diagenesis in the East Midlands Triassic sandstone aquifer. *Geological Society, London, Special Publications*, **34**, 127–140, <https://doi.org/10.1144/GSL.SP.1987.034.01.09>
- Boswell, P.H.G. 1918. *A Memoir of the British Resources of Refractory Sands, Part 1*. Taylor and Francis, London.
- Boulter, M.C. 1970. *An Upper Tertiary Flora from Derbyshire*. Unpublished PhD thesis, University College London.
- Boulter, M.C. 1971a. A palynological study of the Neogene plant beds in Derbyshire. *Bulletin of the British Museum (Natural History) Geology*, **19**, 359–410.

- Boulter, M.C. 1971b. A survey of the Neogene flora from two Derbyshire pocket deposits. *Mercian Geologist*, **4**, 45–62.
- Boulter, M.C. and Chaloner, W.G. 1970. Neogene fossil plants from Derbyshire (England). *Review of Palaeobotany and Palynology*, **10**, 61–78, [https://doi.org/10.1016/0034-6667\(70\)90022-9](https://doi.org/10.1016/0034-6667(70)90022-9)
- Boulter, M.C., Ford, T.D., Ijtaba, M. and Walsh, P.T. 1971. Brassington Formation: a newly recognised Tertiary Formation in the Southern Pennines. *Nature Physical Science*, **231**, 134–136, <https://doi.org/10.1038/physci231134a0>
- Breislin, C., Crowley, S., Banks, V.J., Marshall, J.D., Millar, I.L., Riding, J.B. and Hollis, C. 2020. Controls on dolomitization in extensional basins: an example from the Derbyshire Platform, UK. *Journal of Sedimentary Research*, **90**, 1156–1174, <https://doi.org/10.2110/jsr.2020.58>
- Breislin, C.J., Banks, V.J., Crowley, S., Millar, I.L., Marshall, J.D., Riding, J.B. and Hollis, C. 2023. Mechanisms controlling the localisation of fault-controlled hydrothermal dolomitization on platform margins, Derbyshire Platform, UK. *The Depositional Record*, **9**, 734–758, <https://doi.org/10.1002/dep2.214>
- British Geological Survey 1983. *Ashbourne. England and Wales Sheet 124. Solid with Drift Edition. 1:50 000*. British Geological Survey, Keyworth, Nottingham.
- British Geological Survey 2014. *Derby (125) Sheet 1:50 000 Geological Map of England and Wales*. Natural Environment Research Council.
- British Geological Survey 2025. *The BGS Lexicon of Named Rock Units*, <https://webapps.bgs.ac.uk/lexicon/lexicon.cfm?pub=SSG> [last accessed January 2025].
- Brodie, P.B. 1886. On a remarkable section in Derbyshire. *Geological Magazine*, **3**, 432, <https://doi.org/10.1017/S0016756800467270>
- Brown, E. 1867. The Weaver clays. *Geological Magazine*, **4**, 381–382, <https://doi.org/10.1017/S0016756800206894>
- Bruch, A.A., Uhl, D. and Mosbrugger, V. 2007. Miocene climate in Europe – patterns and evolution: a first synthesis of NECLIME. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **253**, 1–7, <https://doi.org/10.1016/j.palaeo.2007.03.030>
- Burgess, P.M., Dobromylskyj, E. and Lovell–Kennedy, J. 2024. Probably proximal pebbles? An outcrop-constrained quantitative analysis of clast transport distances in the lower Triassic Sherwood Sandstone Group, UK. *Journal of the Geological Society, London*, **182**, jgs2024-155, <https://doi.org/10.1144/jgs2024-15>
- Carney, J.N., Ambrose, K., Brandon, A., Royles, C.P., Cornwell, J.D. and Lewis, M.A. 2001. *Geology of the Country Between Loughborough, Burton and Derby. Sheet Description of the British Geological Survey, 1:50 000 Series Sheet 141 Loughborough (England and Wales)*. British Geological Survey.
- Chaloner, W.G. 1961. Tertiary. In: *Summary of Progress of the Geological Survey of Great Britain and the Museum of Practical Geology for the Year 1960*. Her Majesty's Stationery Office, London, 52.
- Chisholm, J.L., Charsley, T.J. and Aitkenhead, N. 1988. *Geology of the country around Ashbourne and Cheadle. Memoir for 1:50 000 geological sheet 124 (England and Wales)*. Her Majesty's Stationery Office, London.
- Cope, J.C.W., Ingham, J.K. and Rawson, P.F. (eds) 1992. *Atlas of Palaeogeography and Lithofacies*. Geological Society, London, *Memoirs*, **13**.
- Coxon, P. 2005. The late Tertiary landscapes of western Ireland. *Irish Geography*, **38**, 111–127, <https://doi.org/10.1080/00750770509555854>
- Cózar, P., Somerville, I.D., Hounslow, M.W. and Coronado, I. 2022. Far-field correlation of palaeokarstic surfaces in Mississippian successions using high-frequency foraminiferal diversity trends. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **601**, 111088, <https://doi.org/10.1016/j.palaeo.2022.111088>
- Cripps, J.C. and Taylor, R.K. 1987. Engineering characteristics of British over-consolidated clays and mudrocks. II. Mesozoic deposits. *Engineering Geology*, **23**, 213–253, [https://doi.org/10.1016/0013-7952\(87\)90091-3](https://doi.org/10.1016/0013-7952(87)90091-3)
- Davis, A.G. and Chandler, R.J. 1973. *Further Work on the Engineering Properties of Keuper Marl*. Construction Industry Research and Information Association Report, 47.
- Durand, M. and Bourquin, S. 2013. Criteria for the identification of ventifacts in the geological record: a review and new insights. *Comptes Rendus Geoscience*, **345**, 111–125, <https://doi.org/10.1016/j.crte.2013.02.004>
- Edwards, W.N. and Trotter, F.M. 1954. *The Pennines and Adjacent Areas*. 3rd edn. British Regional Geology. Her Majesty's Stationery Office, London.
- Fearnside, W.G. 1932. The geology of the eastern part of the Peak District. *Proceedings of the Geologists' Association*, **43**, 152–191, [https://doi.org/10.1016/S0016-7878\(32\)80026-X](https://doi.org/10.1016/S0016-7878(32)80026-X)
- Ford, T.D. 1968. The Carboniferous Limestone. In: Sylvester-Bradley, P.C. and Ford, T.D. (eds) *The Geology of the East Midlands*. Leicester University Press, 59–82.
- Ford, T.D. 1972. Field meeting in the Peak District. Report by the Director: Trevor D. Ford. *Proceedings of the Geologists' Association*, **83**, 231–236, [https://doi.org/10.1016/S0016-7878\(72\)80009-9](https://doi.org/10.1016/S0016-7878(72)80009-9)
- Ford, T.D. 1999. The growth of geological knowledge in the Peak District. *Mercian Geologist*, **14**, 161–190.
- Ford, T.D. and Jones, J.A. 2007. The geological setting of the mineral deposits at Brassington and Carsington, Derbyshire. *Mining History*, **16**, 1–23.
- Ford, T.D. and King, R.J. 1968. Outliers of possible Tertiary age. In: Sylvester-Bradley, P.C. and Ford, T.D. (eds) *The Geology of the East Midlands*. Leicester University Press, 324–331.
- Ford, T.D. and King, R.J. 1969. The origin of the silica sand pockets in Derbyshire Limestone. *Mercian Geologist*, **3**, 51–69.
- Ford, T.D. and Worley, N.E. 2016. Mineralization of the South Pennine orefield, UK – a review. *Proceedings of the Yorkshire Geological Society*, **61**, 55–86, <https://doi.org/10.1144/pygs2015-364>
- Frost, D.V. and Smart, J.G.O. 1979. *Geology of the Country North of Derby. Memoir for 1:50 000 Geological Sheet 125*. Her Majesty's Stationery Office, London.
- Gale, A.S. and Lovell, B. 2018. The Cretaceous–Paleogene unconformity in England: uplift and erosion related to the Iceland mantle plume. *Proceedings of the Geologists' Association*, **129**, 421–435, <https://doi.org/10.1016/j.pgeola.2017.04.002>
- Gallois, R. 2019. The stratigraphy of the Permo-Triassic rocks of the Dorset and East Devon Coast World Heritage Site, U.K. *Proceedings of the Geologists' Association*, **130**, 274–293, <https://doi.org/10.1016/j.pgeola.2018.01.006>
- Gasparini, A.S., Fontes, M.P.F., Pacheco, A.A. and Ker, J.C. 2022. Gibbsite crystallinity and morphology in ferralsols and bauxites. *Minerals*, **12**, 1441, <https://doi.org/10.3390/min12111441>
- Gibson, M.E., McCoy, J., O'Keefe, J.M.K., Nuñez Otaño, N.B., Warny, S. and Pound, M.J. 2022. Reconstructing terrestrial paleoclimates: a comparison of the co-existence approach, Bayesian and probability reconstruction techniques using the UK Neogene. *Paleoceanography and Paleoclimatology*, **37**, e2021PA004358, <https://doi.org/10.1029/2021PA004358>
- Green, P.F., Thompson, K. and Hudson, J.D. 2001. Recognition of tectonic events in undeformed regions: contrasting results from the Midland Platform and East Midlands Shelf, Central England. *Journal of the Geological Society, London*, **158**, 59–73, <https://doi.org/10.1144/jgs.158.1.59>
- Gutteridge, P. 2024. Lacustrine and palustrine carbonates in a Brigantian (late Dinantian) intrashelf basin in the Derbyshire carbonate platform. *Geological Journal*, **59**, 951–964, <https://doi.org/10.1002/gj.4901>
- Hao, J., Taylor, K.G. and Hollis, C. 2021. Mineral diagenesis and inferred fluids in basinal mudstones: the Carboniferous Morridge Formation, Widmerpool Gulf, UK. *Marine and Petroleum Geology*, **134**, 105373.
- Hennissen, J.A.I., Hough, E., Vane, C.H., Leng, M.J., Kemp, S.J. and Stephenson, M.H. 2017. The prospectivity of a potential shale gas play: an example from the southern Pennine Basin (central England, UK). *Marine and Petroleum Geology*, **86**, 1047–1066, <https://doi.org/10.1016/j.marpetgeo.2017.06.033>
- Hobbs, P.R.N., Hallam, J.R. *et al.* 2002. *Engineering Geology of British Rocks and Soils – Mudstones of the Mercia Mudstone Group*. British Geological Survey Research Report, **RR/01/02**.
- Hodgskiss, M.S.W., Roberts, N.M.W., Paiste, P., Rameil, N., Hammer, E., Brunstad, H. and Lepland, A. 2024. Direct dating of deposition and rift-related alteration of fossil-barren red bed units in the North Sea. *Journal of the Geological Society, London*, **181**, jgs2023-052, <https://doi.org/10.1144/jgs2023-052>
- Holloway, S., Milodowski, A.E., Strong, G.E. and Warrington, G. 1989. The Sherwood Sandstone Group (Triassic) of the Wessex Basin, southern England. *Proceedings of the Geologists' Association*, **100**, 383–394, [https://doi.org/10.1016/S0016-7878\(89\)80056-2](https://doi.org/10.1016/S0016-7878(89)80056-2)
- Howard, A.S., Warrington, G., Ambrose, K. and Rees, J.G. 2008. *A Formational Framework for the Mercia Mudstone Group (Triassic) of England and Wales*. British Geological Survey Research Report, **RR/08/04**.
- Howard, A.S., Warrington, G., Carney, J.N., Ambrose, K., Young, S.R., Pharaoh, T.C. and Cheney, C.S. 2009. *Geology of the Nottingham District. Memoir of the British Geological Survey, Sheet 126 (England and Wales)*. Her Majesty's Stationery Office, London.
- Howe, J.A. 1897. Notes on the pockets of sands and clay in the limestone of Derbyshire and Staffordshire. *Transactions of the North Staffordshire Field Club*, **31**, 143–149.
- Huddart, D. 2002. Pre-Quaternary landscape development. In: Huddart, D. and Glasser, N.F. (eds) *Quaternary of Northern England. Geological Conservation Review Series*, **25**. Joint Nature Conservation Committee, Peterborough, 20–29.
- Hughes, E.M. 1952. *The Geology of an Area North of Brassington*. Unpublished thesis, University of Nottingham.
- Ijtaba, M. 1973. *The Stratigraphy and Sedimentology of the 'Pocket-Deposits' in the Bees Nest and Kirkham Pits, Near Brassington, Derbyshire*. Unpublished Master of Philosophy thesis, Chelsea College of Science and Technology, University of London, repository.royalholloway.ac.uk/file/783f9326-5819-4ade-9852-459fc35e98c6/1/Ijtaba.pdf
- Jackson, A.A. 2008. *Bedrock Geology UK South. An Explanation of the Bedrock Geology Map of England and Wales - 1:625 000*. 5th edn. British Geological Survey, Nottingham.
- Jones, C.V., Reed, S.J.B. and Xing, M. 1993. Heavy mineral stratigraphy in the UK Trias: western Approaches, onshore England and the Central North Sea. *Geological Society, London, Petroleum Geology Conference Series*, **4**, 609–624, <https://doi.org/10.1144/0040609>
- Jones, P.F. and Banks, V.J. 2014. Generating new geo-data. *The Geoscientist*, **24**, 12–17.
- Jones, P.F., Banks, V.J., Pound, M.J. and Riding, J.B. 2016. Glacitectonic structures in the White Peak, Derbyshire. *Quaternary Newsletter*, **140**, 5–19.
- Jones, S.T.J., Worden, R.H., Faulkner, D.R., Fisher, Q.J., McEvoy, J.A. and Utley, J.E.P. 2025. Deposition, diagenesis, and porosity of a siliciclastic caprock: the Late Triassic Mercia Mudstone Group. *Journal of the Geological Society, London*, **182**, jgs2025-086, <https://doi.org/10.1144/jgs2025-086>

- Kender, S., Stephenson, M.H. *et al.* 2012. Marine and terrestrial environmental changes in NW Europe preceding carbon release at the Paleocene–Eocene transition. *Earth and Planetary Science Letters*, **353–354**, 108–120, <https://doi.org/10.1016/j.epsl.2012.08.011>
- Kent, P. 1957. Triassic relics and the 1,000 foot surface in the Southern Pennines. *East Midland Geographer*, **1**, 3–10.
- King, C. 2006. Palaeogene and Neogene: uplift and a cooling climate. In: Brenchley, P.J. and Rawson, P.F. (eds) *The Geology of England and Wales*. 2nd edn. The Geological Society, London, 395–428.
- King, C., Gale, A.S. and Barry, T.L. (eds) 2016. *A Revised Correlation of Tertiary Rocks in the British Isles and Adjacent Areas of NW Europe*. Geological Society Special Report, 27.
- Lamplugh, G.W., Gibson, W., Sherlock, R.L. and Wright, W.B. 1908. *The Geology of the Country Between Newark and Nottingham. Memoir of the Geological Survey of Great Britain, Sheet 126 (England and Wales)*. Her Majesty's Stationery Office, London.
- Lee, J.R., Rose, J. *et al.* 2011. The glacial history of the British Isles during the Early and Middle Pleistocene: implications for the long-term development of the British Ice Sheet. In: Ehlers, J., Gibbard, P.L. and Hughes, P.D. (eds) *Quaternary Glaciations—Extent and Chronology, A Closer Look. Developments in Quaternary Science*, **15**. Elsevier, Amsterdam, The Netherlands, 59–74.
- Lodhia, B.H., Parent, A., Fraser, A.J., Nuemaier, M. and Hennissen, J.A.I. 2023. Thermal evolution and resources of the Bowland Basin (NW England) from apatite fission-track analyses and multidimensional basin modelling. *Geological Society, London, Special Publications*, **534**, 39–60, <https://doi.org/10.1144/SP534-2022-15>
- Manifold, L., Hollis, C. and Burgess, P. 2020. The anatomy of a Mississippian (Viséan) carbonate platform interior, UK: depositional cycles, glacioeustasy and facies mosaics. *Sedimentary Geology*, **401**, 105633, <https://doi.org/10.1016/j.sedgeo.2020.105633>
- McCoy, J.L., Barrass-Barker, T., Hocking, E.P., O'Keefe, J.M.K., Riding, J.B. and Pound, M.J. 2022. Middle Miocene (Serravallian) wetland development on the northwest edge of Europe based on palynological analysis of the uppermost Brassington Formation of Derbyshire, United Kingdom. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **603**, 111180, <https://doi.org/10.1016/j.palaeo.2022.111180>
- Medici, G., Boulesteix, K., Mountney, N.P., West, L.J. and Odling, N.E. 2015. Palaeoenvironment of braided fluvial systems in different tectonic realms of the Triassic Sherwood Sandstone Group, UK. *Sedimentary Geology*, **329**, 188–210, <https://doi.org/10.1016/j.sedgeo.2015.09.012>
- Milroy, P., Wright, V.P. and Simms, M.J. 2019. Dryland continental mudstones: deciphering environmental changes in problematic mudstones from the Upper Triassic (Carnian to Norian) Mercia Mudstone Group, south-west Britain. *Sedimentology*, **66**, 2557–2589, <https://doi.org/10.1111/sed.12626>
- Mustoe, G.E. 2018. Non-mineralized fossil wood. *Geosciences*, **8**, 223, <https://doi.org/10.3390/geosciences8060223>
- Newell, A.J. 2018. Rifts, rivers and climate recovery: a new model for the Triassic of England. *Proceedings of the Geologists' Association*, **129**, 352–371, <https://doi.org/10.1016/j.pgeola.2017.04.001>
- O'Keefe, J.M.K., Pound, M.J., Riding, J.B. and Vane, C.H. 2020. Cellular preservation and maceral development in lignite and wood from the Brassington Formation (Miocene), Derbyshire, UK. *International Journal of Coal Geology*, **222**, 103452, <https://doi.org/10.1016/j.coal.2020.103452>
- Parrish, J.T. 1993. Climate of the supercontinent Pangea. *Journal of Geology*, **101**, 215–233, <https://doi.org/10.1086/648217>
- Pound, M.J. and Riding, J.B. 2015. Miocene in the UK!. *The Geoscientist*, **25**, 20–22.
- Pound, M.J. and Riding, J.B. 2016. Palaeoenvironment, palaeoclimate and age of the Brassington Formation (Miocene) of Derbyshire, UK. *Journal of the Geological Society, London*, **173**, 306–319, <https://doi.org/10.1144/jgs2015-050>
- Pound, M.J., Haywood, A.M., Salzmann, U., Riding, J.B., Lunt, D.J. and Hunter, S.J. 2011. A Tortonian (Late Miocene, 11.61–7.25 Ma) global vegetation reconstruction. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **300**, 29–45, <https://doi.org/10.1016/j.palaeo.2010.11.029>
- Pound, M.J., Riding, J.B., Donders, T.H. and Daskova, J. 2012a. The palynostratigraphy of the Brassington Formation (Upper Miocene) of the southern Pennines, central England. *Palynology*, **36**, 26–37, <https://doi.org/10.1080/01916122.2011.643066>
- Pound, M.J., Haywood, A.M., Salzmann, U. and Riding, J.B. 2012b. Global vegetation dynamics and latitudinal temperature gradients during the Mid to Late Miocene (15.97–5.33 Ma). *Earth-Science Reviews*, **112**, 1–22, <https://doi.org/10.1016/j.earscirev.2012.02.005>
- Pound, M.J., O'Keefe, J.M.K., Nuñez Otaño, N.B. and Riding, J.B. 2019. Three new Miocene fungal palynomorphs from the Brassington Formation, Derbyshire, UK. *Palynology*, **43**, 596–607, <https://doi.org/10.1080/01916122.2018.1473300>
- Pound, M.J., Nuñez Otaño, N.B., Romero, I.C., Lim, M., Riding, J.B. and O'Keefe, J.M.K. 2022. The fungal ecology of the Brassington Formation (Middle Miocene) of Derbyshire, UK, and a new method for palaeoclimate reconstruction. *Frontiers in Ecology and Evolution*, **10**, 947623, <https://doi.org/10.3389/fevo.2022.947623>
- Preto, N., Kustatscher, E. and Wignall, P.B. 2010. Triassic climates – state of the art and perspectives. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **290**, 1–10, <https://doi.org/10.1016/j.palaeo.2010.03.015>
- Püttmann, T. and Mutterlose, J. 2021. Palaeoecology of Late Cretaceous coccolithophores: insights from the shallow-marine record. *Palaeogeography and Palaeoclimatology*, **36**, e2020PA004161, <https://doi.org/10.1029/2020PA004161>
- Radley, J.D. and Coram, R.A. 2016. The Chester Formation (Early Triassic, southern Britain): sedimentary response to extreme greenhouse climate? *Proceedings of the Geologists' Association*, **127**, 552–557, <https://doi.org/10.1016/j.pgeola.2016.08.002>
- Riding, J.B. 2021. A guide to preparation protocols in palynology. *Palynology*, **45**(Supplement 1), 1–157, <https://doi.org/10.1080/01916122.2021.1878305>
- Scotney, P.M., Carney, J.N. and Harwood, M. 2012. New information on Neoproterozoic–Cambrian geology and the Triassic unconformity around Groby, southern Charnwood Forest, UK. *Proceedings of the Yorkshire Geological Society*, **59**, 37–51, <https://doi.org/10.1144/pygs.59.1.302>
- Scott, A. 1927. The origin of the High Peak sand and clay deposits. *Transactions of the British Ceramic Society*, **26**, 255–260.
- Sellwood, B.W. and Valdes, P.J. 2006. Mesozoic climates: general circulation models and the rock record. *Sedimentary Geology*, **190**, 269–287, <https://doi.org/10.1016/j.sedgeo.2006.05.013>
- Smith, S.A. and Edwards, R.A. 1991. Regional sedimentological variations in Lower Triassic fluvial conglomerates (Budleigh Salterton Pebble Beds), southwest England: some implications for palaeogeography and basin evolution. *Geological Journal*, **26**, 65–83, <https://doi.org/10.1002/gj.3350260105>
- Smith, E.G. and Warrington, G. 1971. The age and relationships of the Triassic rocks assigned to the lower part of the Keuper in North Nottinghamshire, north-west Lincolnshire and south Yorkshire. *Proceedings of the Yorkshire Geological Society*, **38**, 201–227, <https://doi.org/10.1144/pygs.38.2.201>
- Smith, N.J.P., Kirby, G.A. and Pharaoh, T.C. 2005. *Structure and Evolution of the South-West Pennine Basin and Adjacent Area*. Subsurface Memoir of the British Geological Survey.
- Steel, R.J. and Thompson, D.B. 1983. Structures and textures in Triassic braided stream conglomerates ('Bunter' Pebble Beds) in the Sherwood Sandstone Group, North Staffordshire, England. *Sedimentology*, **30**, 341–367, <https://doi.org/10.1111/j.1365-3091.1983.tb00677.x>
- Steinthorsdottir, M., Coxall, H.K. *et al.* 2021. The Miocene: the future of the past. *Palaeogeography and Palaeoclimatology*, **36**, e2020PA004037, <https://doi.org/10.1029/2020PA004037>
- Stevenson, I.P. and Gaunt, G.D. 1971. *Geology of the Country Around Chapel-en-le-Frith. Memoirs of the Geological Survey of Great Britain, sheet 99*.
- Swinerton, H.H. 1914. III.—Periods of dreikanter formation in south Notts. *Geological Magazine*, **1**, 208–211, <https://doi.org/10.1017/S0016756800196761>
- Swinerton, H.H. 1935. *The Denudation of the East Midlands*. Report of the British Association for the Advancement of Science, 375.
- Tauxe, L. 2010. *Essentials of Palaeomagnetism*. University of California Press, Berkeley and Los Angeles, USA.
- Thomas, H.H., Hallimond, A.S. and Radley, E.G. 1920. *Refractory Materials: Gannister, Silica Rock, etc. Special Reports on the Mineral Resources of Great Britain*, **16**. Geological Survey, London.
- Thompson, D.B. and Worsley, P. 1967. Periods of ventifact formation in the Permo-Triassic and Quaternary of the north east Cheshire Basin. *Mercian Geologist*, **2**, 279–298.
- Viles, H.A. and Bourke, M.C. 2007. Aeolian features. In: Bourke, M.C. and Viles, H.A. (eds) *A Photographic Atlas of Rock Breakdown Features in Geomorphic Environments*. Planetary Science Institute, Tucson, Arizona, 6–22.
- Wakefield, O.J.W., Hough, E. and Peatfield, A.W. 2015. Architectural analysis of a Triassic fluvial system: the Sherwood Sandstone of the East Midlands Shelf, UK. *Sedimentary Geology*, **327**, 1–13, <https://doi.org/10.1016/j.sedgeo.2015.07.006>
- Walsh, P.T., Boulter, M.C., Ijtaba, M. and Urbani, D.M. 1972. The preservation of the Neogene Brassington Formation of the southern Pennines and its bearing on the evolution of upland Britain. *Journal of the Geological Society, London*, **128**, 519–559, <https://doi.org/10.1144/gsjgs.128.6.0519>
- Walsh, P.T., Collins, P., Ijtaba, M., Newton, J.P., Scott, N.H. and Turner, P.R. 1980. Palaeocurrent directions and their bearing on the origin of the Brassington Formation (Miocene–Pliocene) of the Southern Pennines, Derbyshire, England. *Mercian Geologist*, **8**, 47–62.
- Walsh, P.T., Banks, V.J., Jones, P.F., Pound, M.J. and Riding, J.B. 2018. A reassessment of the Brassington Formation (Miocene) of Derbyshire, UK and a review of related hypogene karst suffosion processes. *Journal of the Geological Society, London*, **175**, 443–463, <https://doi.org/10.1144/jgs2017-029>
- Warrington, G. and Ivey-Cook, H.C. 1992. Triassic. In: Cope, J.C.W., Ingham, J.K. and Rawson, P.F. (eds) *Atlas of Palaeogeography and Lithofacies*. Geological Society, London, Memoirs, **13**, 97–106.
- Warrington, G. and Pollard, J.E. 2021. On the records of the brachiopod 'Lingula' and associated fossils in Mid-Triassic deposits in England. *Proceedings of the*

- Yorkshire Geological Society*, **63**, 319–327, <https://doi.org/10.1144/pygs2020-015>
- Warrington, G., Audley-Charles, M.G. *et al.* 1980. *A Correlation of the Triassic Rocks in the British Isles. Special Report of the Geological Society of London*, **13**.
- Westaway, R. 2009. Quaternary uplift of northern England. *Global and Planetary Change*, **68**, 357–382, <https://doi.org/10.1016/j.gloplacha.2009.03.005>
- Westaway, R. 2020. Late Cenozoic uplift history of the Peak District, central England, inferred from dated cave deposits and integrated with regional drainage development: a review and synthesis. *Quaternary International*, **546**, 20–41, <https://doi.org/10.1016/j.quaint.2019.10.002>
- Wills, L.J. 1970. The Triassic succession in the central Midlands in its regional setting. *Quarterly Journal of the Geological Society of London*, **126**, 225–285, <https://doi.org/10.1144/gsjgs.126.1.0225>
- Wilson, P. 1979. Surface features of quartz sand grains from the Brassington Formation. *Mercian Geologist*, **7**, 19–30.
- Wilson, A.A. 1993. The Mercia Mudstone Group (Trias) of the Cheshire Basin. *Proceedings of the Yorkshire Geological Society*, **49**, 171–188, <https://doi.org/10.1144/pygs.49.3.171>
- Worley, N.E. 1978. *Stratigraphical Control of Mineralisation in the Peak District of Derbyshire*. PhD thesis, University of Leicester, UK, <https://hdl.handle.net/2381/8668>
- Wright, V.P., Marriott, S.B. and Vanstone, S.D. 1991. A ‘reg’ palaeosol from the Lower Triassic of south Devon: stratigraphic and palaeoclimatic implications. *Geological Magazine*, **128**, 517–523, <https://doi.org/10.1017/S0016756800018653>
- Yorke, C. 1954. *The Pocket Deposits of Derbyshire*, **3**. Privately published, Birkenhead.
- Yorke, C. 1960. *The Pocket Deposits of Derbyshire. A General Survey*. Privately published, Birkenhead.
- Yorke, C. 1961. *The Pocket Deposits of Derbyshire. A General Survey*. Privately published, Birkenhead.