

# From Carbon to Plastics: Partitioning of Biogeochemical and Anthropogenic Particles in Penguin Guano

Erica Sparaventi,\* Emily Rowlands, Federico Giglio, Araceli Rodríguez-Romero, Antonio Tovar-Sánchez, and Clara Manno\*



Cite This: <https://doi.org/10.1021/acs.est.5c17586>



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

**ABSTRACT:** Penguins are emblematic inhabitants of the Antarctic continent and play a significant role in the biogeochemical cycles of the Southern Ocean by transferring essential nutrients to marine ecosystems via guano production. Despite their ecological importance, the contribution of penguin guano to carbon cycling remains largely unexplored. Microplastics (MPs) add complexity to these dynamics. MPs in fecal material can limit carbon export by reducing sinking rates and increasing remineralization. We examined guano from two penguin species, the Chinstrap (*P. antarcticus*) ( $n = 25$ ) and Gentoo (*P. papua*) ( $n = 7$ ), from the South Shetland Islands. We quantified biogeochemical particulate components (carbon, nitrogen, and biogenic silica) and characterized MP polymeric composition. Both species showed similar values for natural particulate fractions. Microplastics were detected in 91% of samples, dominated by small particles (25–50  $\mu\text{m}$ , 46%). Chinstrap guano contained the highest amount of MPs. Polypropylene was the predominant polymer (34% in Chinstrap, 75% in Gentoo), followed by polyethylene (37% in Chinstrap, not found in Gentoo). This study provides the first survey of the smallest MP fraction (down to 25  $\mu\text{m}$ ) in penguin guano, offering new insight into how a shift in the partitioning from natural to MP particles, previously overlooked, may influence guano-mediated carbon pathways.

**KEYWORDS:** carbon, microplastics, penguins, guano, Antarctic, biogeochemical cycle



## 1. INTRODUCTION

Penguins represent 90% of seabirds' biomass in the Southern Ocean (SO).<sup>1</sup> Of the 18 existing species, five inhabit the Antarctic continent, including three from the genus *Pygoscelis*, the Adélie (*P. adeliae*), Chinstrap (*P. antarcticus*), and Gentoo (*P. papua*). These are considered the sentinel indicators of marine ecosystem health due to their large populations (estimated at 13.5 million individuals<sup>2</sup>) and their circumpolar distributions. Spending about 75% of their life at sea and occupying a middle position in the Antarctic food web, they predominantly feed on Antarctic krill (*Euphausia superba*, hereafter krill) and are effective indicators of ocean health.<sup>3,4</sup> Penguin guano is the main source of macronutrients (e.g., nitrogen and phosphorus) and micronutrients (e.g., trace metals) on the Antarctic land.<sup>5</sup> Furthermore, recent studies highlighted the important role of penguin guano in SO biogeochemical cycles. The release of dissolved trace metals (e.g., iron, zinc, and copper) mediated by guano enriches the upper layers of the iron-limited SO.<sup>6,7</sup> This input enhances phytoplankton communities' growth, stimulating primary production.<sup>7–9</sup>

The SO plays a key role in global climate system regulation, by coupling physical (e.g., circulation dynamics, upwelling of nutrient-rich waters, and the formation of deep and


intermediated water masses that transport carbon to the ocean floor) and biological processes that drive both local carbon fixation and long-term carbon sequestration.<sup>10–12</sup> In particular, previous studies showed that fecal material produced by organisms at a low trophic level (i.e., zooplankton) represents a fundamental route in the SO for carbon to reach the deep ocean, supporting oceanic carbon sequestration.<sup>13–16</sup> However, the potential contribution of feces produced by upper pelagic predators in the Antarctic ecosystem, including penguin guano, in mediating the downward flux of carbon to the oceanic depths remains almost unexplored. Understanding this contribution could provide valuable insights into the complex mechanisms that regulate carbon export dynamics in the SO.

Microplastic (MP, size between 0.1  $\mu\text{m}$  and 5 mm) pollution further complicates these dynamics. The presence of MPs within fecal pellets can attenuate sinking rates, thereby

**Received:** December 5, 2025

**Revised:** May 28, 2026

**Accepted:** May 29, 2026



**Figure 1.** Maps showing sample collection areas. (a) The Antarctic Peninsula with the South Shetland Islands, the yellow square indicates the two areas considered for the sampling; (b) Sentinel 2A scene of Deception Island on March 17, 2023, the red squares indicate the colonies where Chinstrap penguin guano samples were collected during the austral summers 2020/2021 and 2021/2022; and (c) Sentinel 2A scene of Livingston Island on March 17, 2023, the red squares indicate the colonies where Gentoo penguin guano samples were collected during the austral summer 2021/2022. The size of each square is proportional to the population of penguins in each colony.

enhancing remineralization and limiting vertical carbon export.<sup>17–19</sup> Recent evidence reveals that exposure to polystyrene nanoparticles (NPs, size  $<1\ \mu\text{m}$ ) accelerates krill fecal pellet degradation, potentially reducing carbon sequestration by up to 27% (5.5 Mt. C) per productive season.<sup>20,21</sup> Microplastics have been detected in penguin guano from different species (Gentoo, Adélie, Chinstrap, and King penguins) and locations (South Georgia, South Orkney Islands, and across the Antarctic Peninsula).<sup>22–24</sup> Dietary intake is identified as the primary pathway for MPs, through accidental direct ingestion or indirectly via krill consumption.<sup>24,25</sup> MPs in the SO may originate from local sources, including ship traffic and Antarctic research facilities,<sup>26</sup> as well as from distant lower-latitude seas,<sup>27</sup> with meridional transport across the Antarctic Circumpolar current. While MP concentrations may be minimal when averaged across the SO,<sup>28</sup> they can be relatively high in close proximity to local emission sources, particularly in coastal areas nearby research facilities and shipping routes, where they may pose a localized ecological risk.<sup>26,29–31</sup>


Here, we investigated the particulate composition of guano from two key Antarctic penguin species, Chinstrap and

Gentoo, at Deception and Livingston Islands (South Shetland Islands, off the Antarctic Peninsula). We provide the first integrated quantification of both the biogeochemical (organic carbon (POC), inorganic carbon (PIC), total nitrogen (TN), and biogenic silica (BSi)) and anthropogenic (MP) particulate composition. We identified the MP polymer type down to a size of  $25\ \mu\text{m}$ , an unprecedented resolution for such samples in penguin guano, despite most of the environmental available MPs in Antarctica having a size smaller than  $50\ \mu\text{m}$ .<sup>32,33</sup> Finally, we examined the guano carbon–MPs fraction interaction and the potential implications for the carbon cycle.

## 2. MATERIALS AND METHODS

### 2.1. Penguin Guano Samples Collection

A total of 32 fresh penguin guano samples were collected during two Antarctic campaigns in the South Shetland Islands (SSI). In the austral summers of 2020/2021 and 2021/2022, samples were collected from Chinstrap penguin breeding colonies at Vapour Col ( $62^{\circ}59'S$ ,  $60^{\circ}44'W$ ) (20,000 breeding pairs,<sup>34</sup>) and Baily Head ( $62^{\circ}58'S$ ,  $60^{\circ}30'W$ ) (50,000 breeding pairs,<sup>35</sup>) on Deception Island ( $n = 25$ ) (Figure 1). In the austral summer of 2021/2022, samples were collected from Gentoo penguin breeding colonies at Hannah Point ( $62^{\circ}39'S$ ,  $60^{\circ}36'W$ ), Sally Rock ( $62^{\circ}42'S$ ,  $60^{\circ}25'W$ ), and



**Figure 2.** Schematic illustrations showing the steps involved in the methodology employed for the extraction, characterization (with a  $\mu$ -FTIR spectrometer), and identifications of MPs in penguin guano.

Argentina Cove (62°40'S 60°24'W) on Livingston Island ( $n = 7$ ) (Figure 1). Single colony census is not available for Gentoo penguins since they are distributed across multiple smaller colonies, but total abundance is estimated to be on the order of several thousand breeding pairs ( $10^3$ – $10^4$ ), based on site-specific monitoring data on Livingston Island. Wet guano was carefully collected by hand or with a plastic (polystyrene, PS) spoon from snow, ice, or rocks, avoiding collecting ground remains, placed in a conical centrifuge tube or in a zip-lock bag, and frozen at  $-20$  °C prior to processing in the laboratory. Permissions to work and collect guano samples in the study area were granted by the Spanish Polar Committee.

## 2.2. Quality Assurance/Quality Control (QA/QC)

To avoid potential contamination, all handling steps were performed under controlled clean conditions. Guano samples were collected with PS spoons and stored using falcon tubes and zip-lock bags; however, all storage materials were new and the spoon was cleaned after each use. Samples were handled to minimize external contamination prior to frozen condition storage, with exposure kept to a minimum. Once in the lab, for the microplastic particles extraction procedures, all reagents were first filtered through a  $2$   $\mu$ m polycarbonate filter. All glass and beakers were cleaned with 95% ethanol and ultrapure water before use. Furthermore, all the steps included in the protocol were performed in a laminar flow hood in a restricted access laboratory, and only cotton lab coats and nitrile gloves were used. To check for contamination, three types of blanks were performed: (i) procedural blanks (3 replicates); (ii) lab contamination air blanks (2 replicates); and (iii) storage blanks, for zip-lock bags (2 replicates) and falcon tubes (2 replicates). In particular, a total of 3 procedural blanks were performed for the entire analysis and processed with the same protocol as that used for the samples but using only filtered Milli-Q. In addition, lab contamination air blanks used polycarbonate filters (47 mm diameter and  $0.2$   $\mu$ m pore size) placed in glass Petri dishes on the laboratory bench and simulated all the steps and procedures that have been carried out. Furthermore, storage blanks, consisting of 2 empty zip-lock bags and 2 falcon tubes stored for over a year, were analyzed to check potential contamination during storage. No field blanks were taken to evaluate background contamination since guano sampling was not initially designed for investigating plastic pollution.<sup>23</sup>

## 2.3. Biogeochemical Analysis

Once in the lab, the 32 guano samples were freeze-dried (LABCONCO, model 798,030). Subsamples of dried guano (3.5 mg) were used for analyses of particulate biogeochemical matter. Total particulate carbon (TC), particulate organic carbon (POC), and total nitrogen (TN) were measured in triplicate by combustion in an elemental analyzer (CHN, Exeter Analytical Inc. CE440 elemental analyzer, accuracy  $\pm 0.15\%$ ).<sup>36</sup> For POC determination, samples were pretreated with 1 N hydrochloric acid (HCl, analytical grade,  $\geq 37\%$ , then diluted, Sigma-Aldrich). Particulate inorganic carbon (PIC) was obtained by determining the difference between TC and POC. Three blank filters (GF/F filters, 25 mm,  $0.45$   $\mu$ m pore size) were

interspersed regularly between samples (run every 25 samples) to measure and correct for drift. The amount of particulate biogenic silica (BSi) in guano was quantified by colorimetric analysis (Jasco UVIVDEC accuracy  $\pm 1\%$ ). For the silicate extraction, approximately 10–20 mg of the guano sample was suspended in 50 mL of 0.5 M sodium hydroxide (NaOH, pellets, Carlo Erba, Milan, Italy) at 85 °C for 5 h. Following a progressive dissolution method, an aliquot of each sample was taken for analysis every hour, and the relative silica data were extrapolated back to time zero to correct for the silica originating from coexisting clay minerals.<sup>37</sup> The dissolved silicate was then reacted with ammonium molybdate tetrahydrate (99+%, Thermo Scientific). To shift the solution toward blue (measured at 810 nm), a reducing solution was added, composed of metol sulfite (sodium sulfite anhydrous, Thermo Scientific), 4-methylaminophenol sulfate (99% extra pure, Thermo Scientific), oxalic acid dihydrate (VWR), and sulfuric acid ( $H_2SO_4$ , diluted 50%, 96% purity, Panreac AppliChem).

## 2.4. Microplastics Extraction

We adapted and further improved a methodology used for the extraction and identification of MPs in penguin guano<sup>24</sup> to enable the analyses of smaller MPs ( $>25$   $\mu$ m) (Figure 2). This utilized alkaline digestion, which has been proposed as an effective and cheaper option to enzymatic digestion as it is also less damaging for plastic particles or fibers.<sup>38,39</sup> An initial optimization step was performed to determine the appropriate sample mass required to ensure complete digestion of the guano matrix. Different sample weights were tested, after which a standardized mass of approximately 1.5 g, was selected and used for subsequent analyses. The 32 freeze-dried samples were weighed (1–3 g), transferred into clean glass beakers with a 10% potassium hydroxide (KOH) solution, and placed on a shaker for 72 h at room temperature. The volume of KOH added was 3 times the volume of the biological material. After 72 h, the floating phase was separated from the settled part, by filtration onto separate 10  $\mu$ m metal meshes using a glass filtration setup (45 mm) with a vacuum pump. Due to the large amount of organic matter (mainly due to the presence of a large volume of krill cuticles), the filters from both the floating and settled parts were placed in clean beakers, and a further 10% hydrogen peroxide ( $H_2O_2$ ) solution was added for 24 h. The volume of the solution depended on the amount of the sample, up to a maximum of 100 mL. The digested floating phase filters were then vacuum filtered on Anodiscs (Anopore Inorganic Membrane,  $0.2$   $\mu$ m, 25 mm, Whatman) to allow a full scan of the filter surface for the analysis of the smaller plastics (detection limit down to 11  $\mu$ m) using a Focal Plane Array (FPA)-based imaging. The settled parts were vacuum filtered onto another 10  $\mu$ m metal mesh. The meshes from the settled parts were systematically examined using a stereomicroscope LEICA M80 (Leica Microsystems GmbH) to identify potential MPs particles and/or fibers candidates, which were then hand-picked using forceps and placed onto the Anodiscs for the further analysis. For the selection of possible MPs, the criteria proposed by Hidalgo-Ruz et

al.<sup>40</sup> have been followed as guidelines (i.e., choosing fibers with equal thickness and homogeneous colors).

### 2.5. Microplastics Characterization

A total of 11 digested floating phase Anodiscs were randomly selected from the 32 digested samples, five of Chinstrap penguin guano and six of Gentoo penguin samples, and analyzed for the identification of MP particles using FPA-based imaging. To identify the chemical composition of the particles/fibers, the Anodisc filters were analyzed using the Agilent 670 micro-Fourier Transform Infrared Spectroscopy system ( $\mu$ -FTIR, Agilent Technologies, Santa Clara, CA, USA). The spectrometer was coupled to an Agilent 620 microscope with an automated XYZ-stage and a 128 × 128 focal plane array (FPA) detector, cooled with liquid nitrogen, in the transmission mode. FTIR spectral analysis was conducted in the spectral range of 3600–1250  $\text{cm}^{-1}$  (spectral resolution: 8  $\text{cm}^{-1}$ ). Each filter was fully analyzed by sequentially scanning one-half at a time to ensure accurate FTIR measurement. Before each sample scan, a background scan was collected on a clean Anodisc in the same spectral range. Spectra were further analyzed via Purity Microplastics Finder (MPF) (Purity GmbH, Austria) running a machine learning model (by Hufnagl et al.),<sup>41</sup> which supports the identification of 21 polymers (Supporting Information, Table S1). This post hoc spectral analysis in Purity used a relevance tolerance limit of 0.6, and spectral matches were filtered to match the lower size threshold of the analytical technique of 11  $\mu\text{m}$ .

### 2.6. Microplastic Mass Calculations

The mass of MP particles was determined following the method presented in Rowlands et al.,<sup>42</sup> where the calculation referred to fragments. For fragments, volume was calculated as a cross-sectional area multiplied by thickness. Area measurements derived from the Purity software, which account for concave regions of the 2D shape, were multiplied by fragment thickness ( $T$ ) after adjusting thickness to account for potential (unobserved) concavity across the third dimension. The thickness was estimated from length and width measurements by assuming equality of the width/length and thickness/width ratios. The volume calculation was therefore  $V = P \times T \times \text{area}$ . Fragment mass was then calculated assuming the rough volumes of particles and by multiplying volume by the polymer density (Supporting Information, Table S2).

### 2.7. Statistical Analysis

Statistical differences in the biogeochemical fraction (POC, PIC, TN, BSi, POC/N, and POC/PIC) and the MP polymers (type, size, and the percentage of MPs relative to the total particulate MPs plus POC) between the two studied species have been analyzed using R studio software version 4.4.3 (2025-02-28 ucrt). Normality was tested using Shapiro–Wilk tests; when data met normality assumption, significant differences among species were assessed by one-way ANOVA; otherwise, the nonparametric analysis of variance Kruskal–Wallis was applied. The significance level was set at  $p < 0.05$ .

## 3. RESULTS


### 3.1. QA/QC Results

The three types of blanks performed showed negligible microplastic presence. The procedural blanks (performed along the entire analysis) contained an average of 2 items per filter, with an average size of  $61 \pm 45 \mu\text{m}$ . The lab contamination air blanks contained an average of 2.2 particles per filter (with 94% PA) with an average size of  $41 \pm 15 \mu\text{m}$  (Supporting Information, Table S3). Only two polyamide (PA) particles were identified in the storage blanks, with sizes 39.9 and 62.6  $\mu\text{m}$ . No microplastic items were found in falcon tubes blanks. Therefore, the potential contamination during storage and laboratory processing was minimal. The three types of blanks were used for normalizing the number of MPs extracted and identified from guano samples. For the

procedural and storage blanks data correction, we subtracted the total number of MPs observed from the total number of MPs in samples due to matching in polymer composition and size. For the air contamination blank data correction, we subtracted the mean MP value reported in blanks from the mean MP value of samples.<sup>43</sup>

### 3.2. Guano Biogeochemical Composition

The relative abundance of the biogeochemical particulate components was similar for both penguin species (Figure 3a).




**Figure 3.** (a) Chemical particulate characterization expressed as % of their dry mass) of POC, PIC, TN, and BSi and (b) expressed as POC:N and POC:PIC ratios of Chinstrap and Gentoo penguin guano collected off the Antarctic Peninsula during the austral summers 2020/2021 and 2021/2022. Different letters indicate significant differences ( $p < 0.05$ ) between the two species.

Overall POC was the dominant component of the total guano biomass, followed by TN and PIC. BSi values were under the detection limits for all the samples. The POC concentration, analyzed for each sample in triplicate, from Chinstrap ( $n = 75$ ) and Gentoo ( $n = 21$ ) guano showed no significant difference ( $F_{1,94} = 0.25$ ,  $p > 0.05$ ), with the respective averages of  $25.4 \pm 4.1\%$  and  $26.0 \pm 4.9\%$  of their dry mass (dw). Significant differences were found in PIC ( $H(1) = 6.05$ ,  $p < 0.05$ ) and TN ( $H(1) = 7.43$ ,  $p < 0.05$ ) levels between guano of the two species, being higher in Chinstrap than in Gentoo. Chinstrap PIC average was  $5.9 \pm 3.4\%$  and that of Gentoo was  $3.9 \pm 2.8\%$ ; TN levels were  $10.6 \pm 3.3\%$  for Chinstrap and  $7.9 \pm 1.5\%$  for Gentoo. POC:N and POC:PIC ratios were also significantly different ( $H(1) = 11.79$ ,  $p < 0.001$  and  $H(1) = 4.55$ ,  $p < 0.05$ , respectively) between guano from the two species (Figure 3b). The average value of POC:N in Chinstrap was  $2.6 \pm 0.8$  and that for Gentoo was  $3.4 \pm 0.7$ ; POC:PIC in Chinstrap was  $6.6 \pm 7.0$  and that in Gentoo was  $7.9 \pm 5.0$ .

### 3.3. Guano Microplastic Composition

Microplastic particles were found in 91% of guano samples (10 out of 11); the only MP-free sample was from Gentoo. In total, 101 MP particles were recovered, normalized on guano digested volume ( $\text{MP}/\text{cm}^3$ ). Chinstrap penguin showed a higher presence of MPs, with an average of  $15 \pm 14 \text{MP}/\text{cm}^3$  being found compared to an average of  $4 \pm 2 \text{MP}/\text{cm}^3$  across Gentoo samples (Supporting Information, Table S3), with significant differences found regarding the number of MP particles in guano between the two penguin species ( $H(1) = 11.851$ ,  $p < 0.05$ ). Seven different types of polymers were identified, with the highest frequency polymer being polypropylene (PP) found in 34% of Chinstrap and 75% of Gentoo samples. The second most frequent polymer was polyethylene (PE) with a frequency of 37% in Chinstrap, but it was not found in Gentoo. Polyamide (PA) frequency of occurrence was 18% in Chinstrap samples. Polystyrene (PS) was present in 8% in Chinstrap and 9% in Gentoo samples,


while polyethylene terephthalate (PET) was found in 1% and 15%, respectively. Cellulose acetate (CA) with 1% and polyacrylonitrile (PAN) with 1% were the less common type of polymers, found only in Chinstrap guano samples. A total of 6 fibers were manually isolated. Chinstrap penguin samples showed a higher abundance, with 4 fibers (two PAN, one PA, and one PET), whereas in Gentoo samples were found 2 items (both PET fibers) (Figure 4).



**Figure 4.** Total number and frequency of occurrence (%) of particles and fibers identified via  $\mu$ -FTIR, normalized on guano digested volume ( $\text{cm}^3$ ) across the two penguin species, Chinstrap ( $n = 5$ ) and Gentoo ( $n = 6$ ), categorized by polymer type: polypropylene (PP), polyethylene (PE), polystyrene (PS), polyamide (PA), cellulose acetate (CA), polyethylene terephthalate (PET), and polyacrylonitrile (PAN).

The MPs class size range was from 25  $\mu\text{m}$  to  $>5$  mm, with the smallest (within 25–50  $\mu\text{m}$ ) being the most representative size class, with 68 MPs found (46% of frequency of occurrence), and they were more abundant in Chinstrap samples (41%) than in Gentoo (5%) (Figure 5). No significant differences were found between species and sizes in the Kruskal–Wallis test ( $H(1) = 0.019$ ,  $p > 0.05$ ). The identified fibers ranged in size from 1.3 to  $>5$  mm (Figure 5).

The mass of small MP particles calculated in each sample ranged from 0.05 to 11.6  $\mu\text{g}$ , with an average of  $4.0 \pm 4.6$   $\mu\text{g}$  for Chinstrap samples and  $1.5 \pm 2.3$   $\mu\text{g}$  in Gentoo penguins




**Figure 5.** Frequency of occurrence (%) of MPs divided by size class ( $\mu\text{m}$ ), found in penguin guano samples from Chinstrap and Gentoo penguins.

(Figure 6). MP concentrations, expressed as  $\mu\text{g mg}^{-1}$  of guano dry weight (dw), varied between 0.0001 and 0.004  $\mu\text{g mg}^{-1}$ , with an average of  $0.002 \pm 0.002$   $\mu\text{g mg}^{-1}$  of guano (dw) in Chinstrap and  $0.001 \pm 0.001$   $\mu\text{g mg}^{-1}$  of guano (dw) in Gentoo (Supporting Information, Table S4). The percentage of MPs relative to the total particulate MPs plus POC was less than 2%, Chinstrap:  $0.5 \pm 0.6\%$  and Gentoo:  $0.1 \pm 0.2\%$ , with no significant difference between the two species ( $H(1) = 1.63$ ,  $p > 0.05$ ) and an average value of  $0.3 \pm 0.5\%$  among the two species (Supporting Information, Table S4). This proportion increased to  $3 \pm 8\%$  when plastic fibers  $>5$  mm were also included.

## 4. DISCUSSION

### 4.1. Guano Biogeochemical Particulate Composition

To assess the potential input of carbon by penguin guano in the SO ecosystem, we analyzed the biogeochemical particulate composition of guano from two species of Antarctic penguin, the Chinstrap and Gentoo, from Deception and Livingston Islands. We found that guano from the two penguin species exhibited broadly similar levels of POC and PIC, TN, and BSi. Existing information relating to POC in penguin guano is relatively scarce, and no data regarding on PIC concentrations are currently available in penguin guano samples. Previous studies only reported total organic carbon (indicated as “TC” or “TOC”) values of an average of  $21 \pm 7\%$  (dw) in Chinstrap samples from South Shetland Islands,<sup>44</sup> of up to 30% (dw) in Adélie penguin guano in the Ross Sea region,<sup>45</sup> and of 12% (dw) in emperor penguin guano in eastern Antarctica,<sup>46</sup> which are in accordance with our findings (16.1–37.2% dw). The presence of PIC in both species may derive from minor prey items, such as invertebrate with a carbonate shell, including mollusk gastropod,<sup>47–49</sup> and from lithogenic fragments (i.e., rocks, stones, or pebbles). The stones found in penguin guano are primarily associated with their digestive processes. Penguins ingest stone, known as gastroliths, which serve various functions, including aiding in digestion and potentially assisting in deep diving, or either ingest accidentally through dietary intake.<sup>47,50</sup> The negligible levels of BSi are related to the fact that penguins do not directly consume diatoms, despite a minor amount likely being unintentionally ingested while feeding in seawater, as well as through consumption of krill, which in turn ingest diatoms.<sup>51</sup> The observed higher POC:N ratio in Gentoo guano (2.6–5.0) reveals a relatively more carbon-rich material compared to Chinstrap guano (1.2–4.2), suggesting a different biochemical composition of the prey.<sup>52,53</sup> Different dietary strategies (i.e., whether they feed more on krill or fish/squid) can influence their POC:N ratio.<sup>54,55</sup> Furthermore, Gentoo penguins forage in coastal waters ( $<200$  m water depth), performing both pelagic and especially benthic dives in response to local oceanographic and seafloor features.<sup>56,57</sup> However, Chinstrap forage near the shelf break ( $>200$  m water depth) in the epipelagic/mid layer, likely due to the high availability of krill, where oceanographic conditions promote the formation of dense aggregations of prey and continual replenishment.<sup>56,58</sup> Guano biogeochemical composition can provide information on the distinctive trophic patterns of the two species, despite occupying a shared ecological niche and their similar feeding habits.<sup>49</sup> In this study, POC content in penguin guano contributed on average to 26% of the total guano dry mass, which is very similar to the maximum value observed in krill feces (29%). In the SO, krill



**Figure 6.** (a) Total weight ( $\mu\text{g}$ ) of microplastic particles and fibers and (b) only small MP weight in guano digested samples across the two penguin species, Chinstrap ( $n = 5$ ) and Gentoo ( $n = 6$ ), categorized by polymer type: polypropylene (PP), polyethylene (PE), polystyrene (PS), polyamide (PA), cellulose acetate (CA), polyethylene terephthalate (PET), and polyacrylonitrile (PAN).

feces play a crucial role in the carbon cycle, contributing over 70% of the POC vertical flux.<sup>16,51,58–60</sup> Consequently, penguin guano, similar to krill feces, could be considered an important source of POC in the SO, potentially entering and sustaining the benthic domain. Given the large Pygoscelid penguin population and the huge quantities of guano produced, penguins may represent key elements to conduit carbon export in SO. This is in accordance with the study of Gambi et al.<sup>61</sup> which found an accumulation of particulate organic matter up to 70 m depth, in correspondence of one of the largest Adélie penguin colony in the Ross Sea region.

#### 4.2. Guano Microplastic Composition

We found MPs in 91% of our guano samples (10 out of 11), which is in agreement with a previous study that found MPs in 77% of King penguin (*Aptenodytes patagonicus*) guano samples at South Georgia Island.<sup>23</sup> In contrast, previous reports found lower MPs in penguin guano, ranging between 20% and 29% in Adélie, Chinstrap, and Gentoo penguins across the Antarctic Peninsula.<sup>22,24</sup> However, all these studies focus on MPs larger than  $63 \mu\text{m}$ . Here, we report for the first time the identification of smaller MP particles (down to  $25 \mu\text{m}$ ), a size fraction unexplored in penguin guano. We found 69 MPs, representing 46% of the total MPs, ranging between 25 and  $50 \mu\text{m}$ . This small size range is of particular interest since MPs  $<50 \mu\text{m}$  represents the most abundant size fraction observed in the Southern Ocean. In sea surface water samples from the Weddell Sea, Leistenschneider et al.<sup>33</sup> found the smallest MPs size representing 42% ( $11 \mu\text{m}$ ) and 25% ( $25–50 \mu\text{m}$ ) of the total MPs abundance. Jones-Williams et al.<sup>32</sup> also reported that particles and fibers smaller than  $50 \mu\text{m}$  accounted for 95% of the total MP abundance in Antarctic snow. Our results suggest that previous investigations on MPs in penguin guano may have underestimated the number of MP particles since those measuring less than  $63 \mu\text{m}$  were not considered in previous studies due to procedural limitations. Furthermore, particular attention should be given to the small size range of MPs within fecal matter as larger particles are likely to be retained in the digestive system and/or fragmented prior to excretion.<sup>62</sup> MPs observed here may derive from direct ingestion or may have originated from the breakdown of larger plastic fragments, which represent the most commonly ingested type of MPs by seabirds.<sup>25,63,64</sup> MPs detected in krill span from 2 to  $193 \mu\text{m}$ .<sup>65,66</sup> Therefore, penguins may be exposed to MPs indirectly through trophic transfer via krill consumption.

Although guano sampling was not initially designed for microplastic analysis and minor background contamination cannot be completely ruled out, the consistency of our results with previous studies supports their overall reliability. Here, we found that MP particles were more abundant in Chinstrap (75 MPs) than in Gentoo (27 MPs) penguin guano samples, in accordance with the results of Fragão et al.<sup>24</sup> Different foraging depths between the two species may also partly drive the difference in ingestion, accumulation, and excretion of MPs (in addition to the difference in diet discussed above). This has been observed for other organisms such as two sympatric species of coastal dolphins with different feeding habits in depths, studied in the Atlantic Ocean.<sup>67</sup> Chinstrap penguins, which are epipelagic feeders, may have a greater encounter rate of plastic than Gentoo penguins, which feed predominantly in the benthic domain,<sup>25,55</sup> because low-density MPs are typically buoyant in the water column. This could explain why we only found PET polymers in Gentoo penguin guano, given the PET polymer density ( $1.38 \text{ g cm}^{-3}$ ) is higher than that of the seawater and therefore it is more readily available in benthos. We found PP to be the polymer with the highest frequency of occurrence (34% in Chinstrap and 75% in Gentoo), followed by PE and PA (37% and 18% in Chinstrap penguin). This is in agreement with previous revision studies, which found PP to be the most common polymer detected in Antarctic seabirds' matrices (such as pellets and stomach samples) and in *Pygoscelis* penguin species, among others.<sup>68</sup> More broadly, PP has been noted as the most ingested polymer by seabirds worldwide.<sup>25</sup> Typically, PP is used for packaging, laboratory equipment, bottle caps, and the fishery industry.<sup>69,70</sup> PA, the second most common polymer, is linked to textiles derived from technical clothing and equipment like ropes or route-marking flags.<sup>32</sup> Its ubiquity in Antarctica is well-documented, for example, it is found in feces from three sympatric seal species of the Western Antarctic Peninsula,<sup>71</sup> in penguins' gastrointestinal tracks,<sup>63</sup> and in snow from sites near the Antarctic research camp,<sup>32</sup> indicating a local pollution source (such as the wastewater treatment plants of scientific stations).<sup>72</sup> PE was found the most representative for Spheniscidae,<sup>68</sup> in *Pygoscelis* guano samples and in the digestive tracks of Gentoo penguins' chicks.<sup>24,64</sup> It could be associated with ropes and gear of the fishery industry,<sup>69</sup> which is among the main sources of plastic pollution in the Antarctic region. While the presence of MPs detected here was considerably higher than previously reported (such as number of items), their overall contribution in terms of mass can be similar since

the dimensions cover a size range notably lower than those found in penguin guano elsewhere. For this reason, reporting both the total particles number and their corresponding mass is essential to enable proper comparison among studies as previous research on penguin guano reported MP abundance only, not providing the corresponding MP mass.

### 4.3. Biogeochemical Cycles Implication

Although the biogeochemical and MP fractions of guano were analyzed independently, their interaction has important implications for carbon cycling. The high particulate organic carbon (POC) and carbonate (PIC) contents indicate that penguin guano represents a dense, carbon-rich material with the potential to contribute to vertical carbon export. However, the coexistence of abundant small, low-density MP particles (e.g., PP and PE) within this organic matrix may alter the physicochemical properties of guano, affecting its compactness and sinking behavior (e.g., by increasing buoyancy). In this sense, MPs do not modify the chemical composition of guano directly but may influence its ballasting capacity and remineralization rate, ultimately modulating the efficiency of fecal matter-mediated carbon fluxes in the Southern Ocean.

Here, we show that penguin guano can represent a major source of carbon in coastal areas near the colonies. Therefore, a deeper understanding of the fate and vertical dynamics in the water column (sinking versus floating) of this rich-carbon fecal material is essential to assess the role of penguin guano in mediating carbon export from the ocean surface to the seafloor. The pulpy consistency of penguin guano is more similar to liquid whale feces than zooplankton fecal pellets, which often contain a peritrophic membrane, preventing the release of nutrients and ensuring a more efficient sinking rate.<sup>73</sup> However, despite guano consistency (which will promote the recycling of organic matter in the surface ocean), part of guano will eventually start to sink (e.g., due to ballasting mechanisms for the presence of stones and fragments of organic matter such as krill exoskeletons), acting as a vector of carbon export to the deep ocean. Qualitative observation of guano content confirms the high presence of krill exoskeletons (Supporting Information, Figure S1), which are already considered an important vector for carbon fluxes in the Southern Ocean.<sup>60</sup> Similarly, small guano particles can repack and aggregate with other particles, generating new formations, similar to marine snow, which consists of organic material and phytodetritus that incorporates with inorganic matter, resulting in higher sinking velocity.<sup>74,75</sup> On the other hand, the high presence of small-size MPs found in this study potentially contributes to altering the stability, integrity, and compactness of penguin guano. Laboratory experiments have shown that the incorporation of low-density plastic fragments can alter the structure and density of organic matter, resulting in changes to sinking and remineralization rates, altering POC aggregation,<sup>76,77</sup> thereby decreasing the carbon export to the deep sea. Given the amorphous nature of guano compared with structured fecal pellets, empirical sinking experiments will be required to confirm these physical interactions.

Based on an average guano production rate of 84.5 g of dry guano per individual per day<sup>78</sup> and a POC content of 26% (this study), each penguin contributes approximately 21.97 g of POC daily through guano deposition. Over a 120 day breeding season, this corresponds to an individual export of ~2.6 kg of POC, highlighting the potential role of penguin colonies in carbon fluxes. As for our results, MP particles

represent an average of 0.3% of the total weight of POC plus MPs in the penguins' guano. Thus, we can estimate daily export of MPs per individual to be ~0.07 g. Over a 120 day breeding period, this corresponds to ~8 g per penguin. However, upon scaling up to guano production across Chinstrap and Gentoo penguin populations in the South Shetland Islands (SSI) (about 986,440 Chinstrap individuals;<sup>79</sup> 108,788 Gentoo individuals estimated from Herman et al.<sup>80</sup>), this small amount could be relatively important. Considering that approximately 54% of the total amount of guano produced may contain detectable MPs (data derived from reports on MPs and penguin guano, including the present study), the SSI population could contribute an estimate of 4.7 tons of MPs discharged for the breeding season. Once released into the seawater, MPs in the guano can re-enter biogeochemical cycles and become bioavailable again to zooplankton or undergo flocculation processes. In this context, penguins may act as a vector or transport mechanism for MPs injection in the water column. Moreover, plastic production is increasing globally;<sup>81</sup> therefore, MPs in the SO are expected to rise, in particular in the northern region of the Antarctic peninsula, which is considered as a hotspot of ecological impact for MP accumulation.<sup>26</sup> Consequently, as middle/top epipelagic predators, Antarctic penguins will likely ingest and excrete higher MPs projecting into the future. Furthermore, it is possible to calculate the MPs:POC ratio to evaluate the "natural" versus "anthropogenic" carbon (Table 1). Using an

**Table 1. Estimation of the "Natural" versus "Anthropogenic" Carbon<sup>ab</sup>**

Penguin Species	Total MP in Guano (mg)	Carbon Content of Plastic	Plastic Carbon (mg)	POC Guano (mg)	MPs:POC Ratio
Chinstrap	0.004	0.8268	0.003	0.96	0.003
Gentoo	0.0015	0.8268	0.001	0.99	0.001

<sup>a</sup>Calculation for the MPs:POC ratio in guano from Chinstrap and Gentoo penguins from Deception and Livingston Islands. <sup>b</sup>Note: the carbon content of plastic was estimated in the study by Zhu et al.;<sup>82</sup> the total MP in guano (mg) considered only the smallest fraction of MPs found in this study (not fibers).

average carbon content of plastic of 82.68% (plastic carbon = total plastic  $\times$  0.8268),<sup>82</sup> we calculated the mass ratio between plastic-derived carbon and natural POC (ratio = plastic carbon:POC). The MPs:POC ratio is 0.003 for Chinstrap and 0.001 for Gentoo, indicating that, although present, plastic carbon currently represents less than 0.4% of the total organic carbon pool in the guano. Therefore, while the 'anthropogenic' inclusion of carbon is detectable, it does not yet significantly alter the bulk density or the primary stoichiometry of the carbon export mediated by individuals.

Overall, this study helps boost the information on Chinstrap and Gentoo penguin guano composition in terms of biogeochemical and anthropogenic particular matter, providing new insight into the presence of the smallest class sizes of MP polymers in penguin guano, which had not been detected previously. Our findings suggest that penguin colonies may act as localized hotspots for organic carbon transfer at the terrestrial-maritime interface. Future research should quantify the proportion of guano that remains afloat versus the fraction that sinks, in order to distinguish between material recycled in surface waters and material contributing to carbon export to the deep ocean. On the other hand, our results highlight that

penguins' guano is acting as a vector of MPs in the SO ecosystem. The distinct trophic patterns of the two penguin species define the difference in the biochemical composition of their guano, for both carbon and MPs, thereby influencing its fate and the potential contribution to carbon export to the deep ocean. Identifying the main mechanisms that drive these pathways is essential to estimate the amount of carbon export mediated by guano and to understand how MPs pollution can interfere with the carbon sinking pathway.

## ■ ASSOCIATED CONTENT

### Data Availability Statement

The Supporting Information set is provided in the [Supporting Information](#) section and all data is available at NERC EDS UK Polar Data Centre (10.5285/88c79a01-d3c0-4fca-835c-481930948a5e).

### SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.est.5c17586>.

The supporting data set includes polymers identifiable by the Purity Microplastics Finder (MPF), polymer densities, and detailed information on microplastic polymers and particle masses ([PDF](#))

## ■ AUTHOR INFORMATION

### Corresponding Authors

**Erica Sparaventi** – British Antarctic Survey, Cambridge CB3 0ET, U.K.; Department of Genetic Toxicology and Cancer Biology, National Institute of Biology, Ljubljana 1000, Slovenia; Department of Ecology and Coastal Management, Institute of Marine Sciences of Andalusia, ICMAN (CSIC), Cádiz 11510, Spain; [orcid.org/0000-0002-7578-8918](https://orcid.org/0000-0002-7578-8918); Email: [erica.sparaventi@nib.si](mailto:erica.sparaventi@nib.si)

**Clara Manno** – British Antarctic Survey, Cambridge CB3 0ET, U.K.; Email: [clanno@bas.ac.uk](mailto:clanno@bas.ac.uk)

### Authors

**Emily Rowlands** – British Antarctic Survey, Cambridge CB3 0ET, U.K.

**Federico Giglio** – CNR-ISP—National Research Council of Italy—Institute of Polar Sciences, Bologna 40129, Italy

**Araceli Rodríguez-Romero** – Department of Ecology and Coastal Management, Institute of Marine Sciences of Andalusia, ICMAN (CSIC), Cádiz 11510, Spain

**Antonio Tovar-Sánchez** – Department of Ecology and Coastal Management, Institute of Marine Sciences of Andalusia, ICMAN (CSIC), Cádiz 11510, Spain;

[orcid.org/0000-0003-4375-1982](https://orcid.org/0000-0003-4375-1982)

Complete contact information is available at: <https://pubs.acs.org/10.1021/acs.est.5c17586>

### Author Contributions

E.S.: conceptualization, methodology, fieldwork, data analysis, and original drafting of the manuscript. E.R.: conceptualization, general overview, methodology, data interpretation, writing—reviewing and editing. F.G.: methodology, data interpretation, writing—reviewing and editing. A.R.R.: general overview, conceptualization, writing—reviewing and editing. A.T.S.: funding acquisition, conceptualization, fieldwork, writing—reviewing and editing. C.M.: funding acquisition,

conceptualization, methodology, data analysis, and original drafting of the manuscript.

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

This research falls under the framework of the Spanish Government project PIMETAN (RTI2018-098048-B-I00) and the UKRI-FLF project CUPIDO (MR/T020962/1). E.S. was supported by the Spanish predoctoral FPI grant (PRE2019-089679) and a bursary from Antarctic Science Ltd. We would extend our gratitude to D. Roque for the support in the field, P. Geissler for carried out the CHN analysis, A. Sakovich for her support in the lab, and Dr. Aidan Hunter for assistance with mass calculations. We also thank V. A. Morilla for the scientific illustrations of the Antarctic penguins. This research is part of the POLARCSIC research initiative. We thank the Spanish Antarctic Stations “Gabriel de Castilla” and “Juan Carlos I” and the transportation by the RV Hesperides, as the logistic support by the Marine Technological Unit of CSIC.

## ■ REFERENCES

- (1) Williams, T. Annual Variation in Breeding Biology of Gentoo Penguins, *Pygoscelis Papua*, at Bird Island, South Georgia. *J. Zool.* **1990**, *222* (2), 247–258.
- (2) BirdLife International. *State of the World 'S Birds 2004*, 2004.
- (3) Boersma, P. D. Penguins as Marine Sentinels. *BioScience* **2008**, *58* (7), 597–607.
- (4) Cherel, Y. Isotopic Niches of Emperor and Adélie Penguins in Adélie Land, Antarctica. *Mar. Biol.* **2008**, *154* (5), 813–821.
- (5) Zdanowski, M. K.; Zmuda, M. J.; Zwolska, I. Bacterial Role in the Decomposition of Marine-Derived Material (Penguin Guano) in the Terrestrial Maritime Antarctic. *Soil Biol. Biochem.* **2005**, *37* (3), 581–595.
- (6) Sparaventi, E.; Rodríguez-Romero, A.; Barbosa, A.; Ramajo, L.; Tovar-Sánchez, A. Trace Elements in Antarctic Penguins and the Potential Role of Guano as Source of Recycled Metals in the Southern Ocean. *Chemosphere* **2021**, *285*, 131423.
- (7) Belyaev, O.; Sparaventi, E.; Navarro, G.; Rodríguez-Romero, A.; Tovar-Sánchez, A. The Contribution of Penguin Guano to the Southern Ocean Iron Pool. *Nat. Commun.* **2023**, *14* (1), 1781.
- (8) Shatova, O.; Wing, S. R.; Gault-Ringold, M.; Wing, L.; Hoffmann, L. J. Seabird Guano Enhances Phytoplankton Production in the Southern Ocean. *J. Exp. Mar. Biol. Ecol.* **2016**, *483*, 74–87.
- (9) Shatova, O. A.; Wing, S. R.; Hoffmann, L. J.; Wing, L. C.; Gault-Ringold, M. Phytoplankton Community Structure Is Influenced by Seabird Guano Enrichment in the Southern Ocean. *Estuar. Coast. Shelf Sci.* **2017**, *191*, 125–135.
- (10) Khatiwala, S.; Primeau, F.; Hall, T. Reconstruction of the History of Anthropogenic CO<sub>2</sub> Concentrations in the Ocean. *Nature* **2009**, *462* (7271), 346–349.
- (11) Frölicher, T. L.; Sarmiento, J. L.; Paynter, D. J.; Dunne, J. P.; Krasting, J. P.; Winton, M. Dominance of the Southern Ocean in Anthropogenic Carbon and Heat Uptake in CMIP5 Models. *J. Clim.* **2015**, *28* (2), 862–886.
- (12) Gray, A. R. The Four-Dimensional Carbon Cycle of the Southern Ocean. *Annu. Rev. Mar. Sci.* **2024**, *16* (1), 163–190.
- (13) Manno, C.; Stowasser, G.; Enderlein, P.; Fielding, S.; Tarling, G. A. The Contribution of Zooplankton Faecal Pellets to Deep-Carbon Transport in the Scotia Sea (Southern Ocean). *Biogeosciences* **2015**, *12* (6), 1955–1965.
- (14) Manno, C.; Giglio, F.; Stowasser, G.; Fielding, S.; Enderlein, P.; Tarling, G. A. Threatened Species Drive the Strength of the Carbonate Pump in the Northern Scotia Sea. *Nat. Commun.* **2018**, *9* (1), 4592.

- (15) Iversen, M. H.; Pakhomov, E. A.; Hunt, B. P. V.; Van Der Jagt, H.; Wolf-Gladrow, D.; Klaas, C. Sinkers or Floaters? Contribution from Salp Pellets to the Export Flux during a Large Bloom Event in the Southern Ocean. *Deep Sea Res. Part II Top. Stud. Oceanogr.* **2017**, *138*, 116–125.
- (16) Belcher, A.; Henson, S. A.; Manno, C.; Hill, S. L.; Atkinson, A.; Thorpe, S. E.; Fretwell, P.; Ireland, L.; Tarling, G. A. Krill Faecal Pellets Drive Hidden Pulses of Particulate Organic Carbon in the Marginal Ice Zone. *Nat. Commun.* **2019**, *10* (1), 1–8.
- (17) Cole, M.; Lindeque, P. K.; Fileman, E.; Clark, J.; Lewis, C.; Halsband, C.; Galloway, T. S. Microplastics Alter the Properties and Sinking Rates of Zooplankton Faecal Pellets. *Environ. Sci. Technol.* **2016**, *50*, 3239–3246.
- (18) Wieczorek, A. M.; Croot, P. L.; Lombard, F.; Sheahan, J. N.; Doyle, T. K. Microplastic Ingestion by Gelatinous Zooplankton May Lower Efficiency of the Biological Pump. *Environ. Sci. Technol.* **2019**, *53* (9), 5387–5395.
- (19) Bergami, E.; Manno, C.; Cappello, S.; Vannuccini, M. L.; Corsi, I. Nanoplastics Affect Moulting and Faecal Pellet Sinking in Antarctic Krill (*Euphausia Superba*) Juveniles. *Environ. Int.* **2020**, *143* (February), 105999.
- (20) Manno, C.; Corsi, I.; Rowlands, E.; Bergami, E. Plastics Counteract the Ability of Antarctic Krill to Promote the Blue Carbon Pathway in the Deep Ocean. *Mar. Pollut. Bull.* **2024**, *209*, 117238.
- (21) Barnes, D. K. A.; Galgani, F.; Thompson, R. C.; Barlaz, M. Accumulation and Fragmentation of Plastic Debris in Global Environments. *Philos. Trans. R. Soc., B* **2009**, *364* (1526), 1985–1998.
- (22) Bessa, F.; Ratcliffe, N.; Otero, V.; Sobral, P.; Marques, J. C.; Waluda, C. M.; Trathan, P. N.; Xavier, J. C. Microplastics in Gentoo Penguins from the Antarctic Region. *Sci. Rep.* **2019**, *9* (1), 14191.
- (23) Le Guen, C.; Suaria, G.; Sherley, R. B.; Ryan, P. G.; Aliani, S.; Boehme, L.; Brierley, A. S. Microplastic Study Reveals the Presence of Natural and Synthetic Fibres in the Diet of King Penguins (*Aptenodytes Patagonicus*) Foraging from South Georgia. *Environ. Int.* **2020**, *134*, 105303.
- (24) Fragão, J.; Bessa, F.; Otero, V.; Barbosa, A.; Sobral, P.; Waluda, C. M.; Guimarães, H. R.; Xavier, J. C. Microplastics and Other Anthropogenic Particles in Antarctica: Using Penguins as Biological Samplers. *Sci. Total Environ.* **2021**, *788*, 147698.
- (25) Kibria, G. Contamination of Coastal and Marine Bird Species with Plastics: Global Analysis and Synthesis. *Mar. Pollut. Bull.* **2024**, *206*, 116687.
- (26) Hunter, A.; Thorpe, S. E.; McCarthy, A. H.; Manno, C. Microplastic Hotspots Mapped across the Southern Ocean Reveal Areas of Potential Ecological Impact. *Sci. Rep.* **2024**, *14* (1), 31599.
- (27) Isobe, A.; Uchiyama-Matsumoto, K.; Uchida, K.; Tokai, T. Microplastics in the Southern Ocean. *Mar. Pollut. Bull.* **2017**, *114* (1), 623–626.
- (28) Suaria, G.; Perold, V.; Lee, J. R.; Lebouard, F.; Aliani, S.; Ryan, P. G. Floating Macro- and Microplastics around the Southern Ocean: Results from the Antarctic Circumnavigation Expedition. *Environ. Int.* **2020**, *136*, 105494.
- (29) Gurumoorthi, K.; Luis, A. J. Recent Trends on Microplastics Abundance and Risk Assessment in Coastal Antarctica: Regional Meta-Analysis. *Environ. Pollut.* **2023**, *324*, 121385.
- (30) Gonzalez-Pineda, M.; Salvadó, H.; Avila, C. Do Antarctic Bivalves Present Microdebris? The Case of Livingston Island. *Environ. Pollut.* **2024**, *351*, 124086.
- (31) Jain, A.; Hughes, K. A.; Manno, C. Low-Cost Preventative Measures Can Effectively Mitigate Microplastic Release from Scientific Research Facilities in Antarctica. *Antarct. Sci.* **2025**, *37*, 372–383.
- (32) Jones-Williams, K.; Rowlands, E.; Primpke, S.; Galloway, T.; Cole, M.; Waluda, C.; Manno, C. Microplastics in Antarctica—A Plastic Legacy in the Antarctic Snow? *Sci. Total Environ.* **2025**, *966*, 178543.
- (33) Leistenschneider, C.; Wu, F.; Primpke, S.; Gerdt, G.; Burkhardt-Holm, P. Unveiling High Concentrations of Small Microplastics (11–500 Mm) in Surface Water Samples from the Southern Weddell Sea off Antarctica. *Sci. Total Environ.* **2024**, *927*, 172124.
- (34) Belyaev, O.; Román, A.; Belliure, J.; Navarro, G.; Barbero, L.; Tovar-Sánchez, A. Assessing Topographic Features and Population Abundance in an Antarctic Penguin Colony through UAV-Based Deep-Learning Models. *Int. J. Appl. Earth Obs. Geoinformation* **2024**, *134*, 104124.
- (35) Naveen, R.; Lynch, H. J.; Forrest, S.; Mueller, T.; Polito, M. First Direct, Site-Wide Penguin Survey at Deception Island, Antarctica, Suggests Significant Declines in Breeding Chinstrap Penguins. *Polar Biol.* **2012**, *35* (12), 1879–1888.
- (36) Nieuwenhuize, J.; Maas, Y. E. M.; Middelburg, J. J. Rapid Analysis of Organic Carbon and Nitrogen in Particulate Materials. *Mar. Chem.* **1994**, *45* (3), 217–224.
- (37) Demaster, D. J. Measuring Biogenic Silica in Marine Sediments and Suspended Matter. In *Geophysical Monograph Series*; Hurd, D. C., Spencer, D. W., Eds.; American Geophysical Union: Washington, D. C., 1991; pp 363–367.
- (38) Jones-Williams, K.; Galloway, T.; Cole, M.; Stowasser, G.; Waluda, C.; Manno, C. Close Encounters - Microplastic Availability to Pelagic Amphipods in Sub-Antarctic and Antarctic Surface Waters. *Environ. Int.* **2020**, *140*, 105792.
- (39) Kühn, S.; Van Werven, B.; Van Oyen, A.; Meijboom, A.; Bravo Rebolledo, E. L.; Van Franeker, J. A. The Use of Potassium Hydroxide (KOH) Solution as a Suitable Approach to Isolate Plastics Ingested by Marine Organisms. *Mar. Pollut. Bull.* **2017**, *115* (1–2), 86–90.
- (40) Hidalgo-Ruz, V.; Gutow, L.; Thompson, R. C.; Thiel, M. Microplastics in the Marine Environment: A Review of the Methods Used for Identification and Quantification. *Environ. Sci. Technol.* **2012**, *46* (6), 3060–3075.
- (41) Hufnagl, B.; Stibi, M.; Martirosyan, H.; Wilczek, U.; Möller, J. N.; Löder, M. G. J.; Laforsch, C.; Löhniger, H. Computer-Assisted Analysis of Microplastics in Environmental Samples Based on  $\mu$ FTIR Imaging in Combination with Machine Learning. *Environ. Sci. Technol. Lett.* **2022**, *9* (1), 90–95.
- (42) Rowlands, E.; Galloway, T.; Cole, M.; Peck, V. L.; Posacka, A.; Thorpe, S.; Manno, C. Vertical Flux of Microplastic, a Case Study in the Southern Ocean, South Georgia. *Mar. Pollut. Bull.* **2023**, *193*, 115117.
- (43) Dawson, A. L.; Santana, M. F. M.; Nelis, J. L. D.; Motti, C. A. Taking Control of Microplastics Data: A Comparison of Control and Blank Data Correction Methods. *J. Hazard. Mater.* **2023**, *443*, 130218.
- (44) Pérez-Fernández, B.; Caza, J.; De La Cerda, M. C.; Pérez-Alberti, A.; Viñas, L.; Oliva, M.; Ruiz-Fernández, J.; García-Hernández, C.; Otero, X. L. Occurrence and Sources of Pollutants in the Soils of Penguin Colonies in the South Shetland Islands, North-West Coast of the Antarctic Peninsula. *Geoderma* **2025**, *461*, 117482.
- (45) Liu, X.; Nie, Y.; Sun, L.; Emslie, S. D. Eco-Environmental Implications of Elemental and Carbon Isotope Distributions in Ornithogenic Sediments from the Ross Sea Region, Antarctica. *Geochim. Cosmochim. Acta* **2013**, *117*, 99–114.
- (46) Zhu, R.; Liu, Y.; Ma, E.; Sun, J.; Xu, H.; Sun, L. Greenhouse Gas Emissions from Penguin Guanos and Ornithogenic Soils in Coastal Antarctica: Effects of Freezing-Thawing Cycles. *Atmos. Environ.* **2009**, *43* (14), 2336–2347.
- (47) Clausen, A.; Pütz, K. Winter Diet and Foraging Range of Gentoo Penguins (*Pygoscelis Papua*) from Kidney Cove, Falkland Islands. *Polar Biol.* **2003**, *26* (1), 32–40.
- (48) Tabassum, N.; Lee, J.-H.; Lee, S.-R.; Kim, J.-U.; Park, H.; Kim, H.-W.; Kim, J.-H. Molecular Diet Analysis of Adélie Penguins (*Pygoscelis Adeliae*) in the Ross Sea Using Fecal DNA. *Biology* **2022**, *11* (2), 182.
- (49) Wawrzyniek-Borejko, J.; Panasiuk, A.; Hinke, J. T.; Korczak-Abshire, M. Are the Diets of Sympatric Pygoscelid Penguins More Similar than Previously Thought? *Polar Biol.* **2022**, *45* (10), 1559–1569.
- (50) Beaune, D.; Le Bohec, C.; Lucas, F.; Gauthier-Clerc, M.; Le Maho, Y. Stomach Stones in King Penguin Chicks. *Polar Biol.* **2009**, *32* (4), 593–597.

- (51) Haberman, K. L.; Ross, R. M.; Quetin, L. B. Diet of the Antarctic Krill (*Euphausia Superba* Dana): II. Selective Grazing in Mixed Phytoplankton Assemblages. *J. Exp. Mar. Biol. Ecol.* **2003**, *283* (1–2), 97–113.
- (52) Gleiber, M.; Steinberg, D.; Ducklow, H. Time Series of Vertical Flux of Zooplankton Fecal Pellets on the Continental Shelf of the Western Antarctic Peninsula. *Mar. Ecol.: Prog. Ser.* **2012**, *471*, 23–36.
- (53) Morales, C. Carbon and Nitrogen Content of Copepod Faecal Pellets: Effect of Food Concentration and Feeding Behavior. *Mar. Ecol.: Prog. Ser.* **1987**, *36*, 107–114.
- (54) Atkinson, A.; Schmidt, K.; Fielding, S.; Kawaguchi, S.; Geissler, P. A. Variable Food Absorption by Antarctic Krill: Relationships between Diet, Egestion Rate and the Composition and Sinking Rates of Their Fecal Pellets. *Deep Sea Res. Part II Top. Stud. Oceanogr.* **2012**, *59*–60, 147–158.
- (55) Saba, G. K.; Burd, A. B.; Dunne, J. P.; Hernández-León, S.; Martin, A. H.; Rose, K. A.; Salisbury, J.; Steinberg, D. K.; Trueman, C. N.; Wilson, R. W.; Wilson, S. E. Toward a Better Understanding of Fish-based Contribution to Ocean Carbon Flux. *Limnol. Oceanogr.* **2021**, *66* (5), 1639–1664.
- (56) Kokubun, N.; Takahashi, A.; Mori, Y.; Watanabe, S.; Shin, H.-C. Comparison of Diving Behavior and Foraging Habitat Use between Chinstrap and Gentoo Penguins Breeding in the South Shetland Islands, Antarctica. *Mar. Biol.* **2010**, *157* (4), 811–825.
- (57) Takahashi, A.; Kokubun, N.; Mori, Y.; Shin, H.-C. Krill-Feeding Behaviour of Gentoo Penguins as Shown by Animal-Borne Camera Loggers. *Polar Biol.* **2008**, *31* (10), 1291–1294.
- (58) Warwick-Evans, V.; Kelly, N.; Dalla Rosa, L.; Friedlaender, A.; Hinke, J. T.; Kim, J. H.; Kokubun, N.; Santora, J. A.; Secchi, E. R.; Seyboth, E.; Trathan, P. N. Using Seabird and Whale Distribution Models to Estimate Spatial Consumption of Krill to Inform Fishery Management. *Ecosphere* **2022**, *13* (6), No. e4083.
- (59) Cavan, E. L.; Mackay, N.; Hill, S. L.; Atkinson, A.; Belcher, A.; Visser, A. Antarctic Krill Sequester Similar Amounts of Carbon to Key Coastal Blue Carbon Habitats. *Nat. Commun.* **2024**, *15* (1), 7842.
- (60) Manno, C.; Fielding, S.; Stowasser, G.; Murphy, E. J.; Thorpe, S. E.; Tarling, G. A. Continuous Moulting by Antarctic Krill Drives Major Pulses of Carbon Export in the North Scotia Sea, Southern Ocean. *Nat. Commun.* **2020**, *11* (1), 6051.
- (61) Gambi, C.; Lo Martire, M.; Cantafaro, A. L.; Munari, C.; Rastelli, E.; Danovaro, R.; Corinaldesi, C. Effects of an Adélie Penguin Colony on Coastal Meiofaunal Assemblages of the Ross Sea (Antarctica). *Polar Biol.* **2025**, *48* (2), 55.
- (62) Monclús, L.; McCann Smith, E.; Ciesielski, T. M.; Wagner, M.; Jaspers, V. L. B. Microplastic Ingestion Induces Size-Specific Effects in Japanese Quail. *Environ. Sci. Technol.* **2022**, *56* (22), 15902–15911.
- (63) Cole, M.; Lindeque, P.; Halsband, C.; Galloway, T. S. Microplastics as Contaminants in the Marine Environment: A Review. *Mar. Pollut. Bull.* **2011**, *62* (12), 2588–2597.
- (64) Kim, Y.; Kim, H.; Jeong, M.-S.; Kim, D.; Kim, J.; Jung, J.; Seo, H.-M.; Han, H.-J.; Lee, W.-S.; Choi, C.-Y. Microplastics in Gastrointestinal Tracts of Gentoo Penguin (*Pygoscelis Papua*) Chicks on King George Island, Antarctica. *Sci. Rep.* **2023**, *13* (1), 13016.
- (65) Wilkie Johnston, L.; Bergami, E.; Rowlands, E.; Manno, C. Organic or Junk Food? Microplastic Contamination in Antarctic Krill and Salps. *R. Soc. Open Sci.* **2023**, *10* (3), 221421.
- (66) Zhu, W.; Liu, W.; Chen, Y.; Liao, K.; Yu, W.; Jin, H. Microplastics in Antarctic Krill (*Euphausia Superba*) from Antarctic Region. *Sci. Total Environ.* **2023**, *870*, 161880.
- (67) Di Benedetto, A. P. M.; Ramos, R. M. A. Marine Debris Ingestion by Coastal Dolphins: What Drives Differences between Sympatric Species? *Mar. Pollut. Bull.* **2014**, *83* (1), 298–301.
- (68) Taurozzi, D.; Scalici, M. Seabirds from the Poles: Microplastics Pollution Sentinels. *Front. Mar. Sci.* **2024**, *11*, 1343617.
- (69) Andrady, A. L. Microplastics in the Marine Environment. *Mar. Pollut. Bull.* **2011**, *62* (8), 1596–1605.
- (70) Mishra, A. K.; Singh, J.; Mishra, P. P. Microplastics in Polar Regions: An Early Warning to the World's Pristine Ecosystem. *Sci. Total Environ.* **2021**, *784*, 147149.
- (71) Cebuhar, J. D.; Negrete, J.; Rodríguez Pirani, L. S.; Picone, A. L.; Proietti, M.; Romano, R. M.; Della Védova, C. O.; Casaux, R.; Secchi, E. R.; Botta, S. Anthropogenic Debris in Three Sympatric Seal Species of the Western Antarctic Peninsula. *Sci. Total Environ.* **2024**, *922*, 171273.
- (72) Antacli, J. C.; Di Mauro, R.; Rimondino, G. N.; Alurralde, G.; Schloss, I. R.; González, G. A.; Morales, S.; Ottero, A.; Vodopivec, C. Microplastic Pollution in Waters of the Antarctic Coastal Environment of Potter Cove (25 de Mayo Island/King George Island, South Shetlands). *Sci. Total Environ.* **2024**, *915*, 170155.
- (73) Ratnarajah, L.; Nicol, S.; Bowie, A. R. Pelagic Iron Recycling in the Southern Ocean: Exploring the Contribution of Marine Animals. *Front. Mar. Sci.* **2018**, *5* (MAR), 1–9.
- (74) Durkin, C. A.; Buesseler, K. O.; Cetinić, I.; Estapa, M. L.; Kelly, R. P.; Omand, M. A Visual Tour of Carbon Export by Sinking Particles. *Glob. Biogeochem. Cycles* **2021**, *35* (10), No. e2021GB006985.
- (75) Wu, N.; Grieve, S. W. D.; Manning, A. J.; Spencer, K. L. Marine Snow as Vectors for Microplastic Transport: Multiple Aggregation Cycles Account for the Settling of Buoyant Microplastics to Deep-sea Sediments. *Limnol. Oceanogr.* **2025**, *70* (4), 899–910.
- (76) Chen, C.-S.; Le, C.; Chiu, M.-H.; Chin, W.-C. The Impact of Nanoplastics on Marine Dissolved Organic Matter Assembly. *Sci. Total Environ.* **2018**, *634*, 316–320.
- (77) Porter, A.; Lyons, B. P.; Galloway, T. S.; Lewis, C. Role of Marine Snows in Microplastic Fate and Bioavailability. *Environ. Sci. Technol.* **2018**, *52* (12), 7111–7119.
- (78) Sun, L.; Xie, Z. Relics: Penguin Population Programs. *Sci. Prog.* **2001**, *84* (1), 31–44.
- (79) Borboroglu, P.; Boersma, P. D. *Penguins: Natural History and Conservation*; JSTOR, 2013.
- (80) Herman, R.; Borowicz, A.; Lynch, M.; Trathan, P.; Hart, T.; Lynch, H. Update on the Global Abundance and Distribution of Breeding Gentoo Penguins (*Pygoscelis Papua*). *Polar Biol.* **2020**, *43* (12), 1947–1956.
- (81) OECD. *Global Plastics Outlook: Policy Scenarios to 2060*; OECD, 2022.
- (82) Zhu, L.; Zhao, S.; Bittar, T. B.; Stubbins, A.; Li, D. Photochemical Dissolution of Buoyant Microplastics to Dissolved Organic Carbon: Rates and Microbial Impacts. *J. Hazard. Mater.* **2020**, *383*, 121065.