

GCB REVIEW OPEN ACCESS

Climate Change Experienced by Life on Earth—Drivers and Mechanisms of Future Trends in Microclimates

C. Greiser^{1,2}  | E. Gril³  | I. M. D. Maclean^{4,5}  | R. Beugnon^{6,7,8,9}  | P. le Roux¹⁰  | J. Kemppinen^{11,12}  | M. W. Rhodes⁴  | J. Aalto^{13,14}  | D. M. Christiansen^{15,16}  | P. De Frenne¹⁷  | S. Haesen^{18,19}  | D. Klimes^{20,21}  | M. Kopecky^{22,23}  | J. Lenoir²⁴  | M. Luoto¹⁴  | L. M. Mercado^{25,26}  | M. Rantanen¹³  | G. Strandberg^{27,28}  | L. van den Brink²⁹  | J. Lembrechts^{9,30}  | K. Van Meerbeek^{18,19} 

Correspondence: C. Greiser (caroline.greiser@su.se)

Received: 24 November 2025 | **Revised:** 31 August 2026 | **Accepted:** 20 September 2026

Keywords: biosphere-atmosphere interactions | buffering | clouds | cold-air pooling | decoupling | evapotranspiration | surface-energy balance | vegetation-climate feedbacks

ABSTRACT

Microclimate, the climate directly experienced by organisms, is crucial for understanding ecological responses to climate change, yet its long-term dynamics and the drivers of decadal microclimate changes remain underexplored in ecology and global change biology. Microclimate changes do not necessarily track macroclimate trends. Changes in meteorological conditions (e.g., cloud cover, wind patterns) and vegetation dynamics (e.g., canopy opening, desertification in drylands, shrub encroachment in the tundra) associated with climate change can lead to locally divergent microclimate trends, thereby altering spatial variability. Understanding and quantifying how macroclimate change translates to fine-scale microclimate change experienced by organisms, populations and communities is therefore essential for predicting ecological responses to global environmental change. This review examines how macroclimate change drives microclimate changes as experienced by life on Earth—and how these micro-scale changes may, in turn, feed back into the regional and global climate system. Our primary focus is on near-surface air and soil temperature—due to their relevance to most terrestrial organisms and ecosystem processes. The review is structured around three pathways of microclimate changes: (I) direct effects of changes in meteorological conditions, (II) indirect effects via vegetation changes, and (III) feedbacks between microclimate and climate change. Using three mechanistic model simulations in real-world landscapes, we illustrate and quantify the key mechanisms that lead to divergent warming trends. We show that microclimate change emerges from interacting atmospheric, land-surface, and biophysical processes, which can differ substantially across space and time. Predicting future microclimates therefore requires accounting for these processes and their temporal dynamics. By providing an accessible synthesis of these processes, this review aims to strengthen links between ecology and atmospheric science and improve our understanding of how climate change translates into microclimatic conditions relevant to life on Earth.

1 | Introduction

Climate change impacts species worldwide (Pörtner et al. 2021). Shifts in temperatures, precipitation, and the frequency and

severity of extreme weather events are already affecting the phenology, behavior, and distribution of organisms as well as the functioning of entire ecosystems (Grimm et al. 2013; Muthukrishnan et al. 2025; Soifer et al. 2025), with even

C. Greiser, E. Gril, and I. M. D. Maclean shared first authors.

J. Lembrechts and K. Van Meerbeek shared last authors.

For affiliations refer to page 15.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2026 The Author(s). *Global Change Biology* published by John Wiley & Sons Ltd.

greater impacts projected for the coming decades (IPBES 2019). However, climate change projections are primarily based on global and regional climate models that operate at coarse spatial resolutions (typically 10–100 km), which fail to capture the microclimatic conditions experienced by organisms (Christiansen et al. 2024; Haesen, Lenoir, et al. 2023; Klimes et al. 2024; see definitions in Box 1). As a result, failing to account for the microclimatic conditions experienced by organisms remains an important limitation when predicting biological responses to climate change.

Microclimates are shaped by interactions between atmospheric conditions and the Earth's surface, with local properties such as topography, vegetation, soils and water bodies modifying the exchange of energy and moisture (see Boxes 1 and 2), creating fine-scale heterogeneous environments that can differ substantially—both spatially and temporally—from macroclimate averages, which represent broader geographic regions (Geiger 1966). In this review, we focus on microclimate change—defined as long-term (decadal or longer) changes in near-surface temperature. Importantly, microclimate changes do not always mirror macroclimate changes. Shifts in meteorological conditions directly influence microclimate warming rates, or indirectly do so by altering vegetation structure and composition (Seidl et al. 2017). In turn, changes in land surface properties can feed back on meteorological conditions through multiple biophysical processes, including vegetation-climate interactions (Duvellier et al. 2021), further influencing the extent to which microclimate change tracks macroclimatic trends (Luo et al. 2024a; Miralles et al. 2025). The decoupling between microclimate and macroclimate changes leads to spatially and temporally variable differences between microclimate and macroclimate. Using a mechanistic modeling approach, Maclean (2020) reconstructed daily maximum near-ground temperatures over approximately three decades, revealing divergent temperature trends among locations within a 2 × 2 km landscape in the UK (Figure 1). Beyond the microclimate scale, spatial heterogeneity in warming rates

also emerges at the mesoscale, for instance through variations in elevational lapse rates, coastal–inland gradients, and continentality effects. These mesoclimatic processes set the climatic context for microclimates and can lead to stronger or weaker local warming, thereby influencing the magnitude and direction of micro-macroclimate decoupling.

Despite their importance, microclimates are still often overlooked in climate change impact assessments. Yet it is these proximal conditions that determine how individual organisms, populations, and ecosystems respond to environmental change (Kemppinen et al. 2024; Kemppinen, Niittynen, le Roux, et al. 2021; Körner and Hiltbrunner 2018). Ignoring the fine-scale variability in microclimate trends risks misjudging both the vulnerability and resilience of species and ecosystems to climate change (Niittynen et al. 2018). By integrating microclimate change into ecological research, we can improve predictions of species distributions (Haesen, Lenoir, et al. 2023; Maclean and Early 2023), physiological responses (Buckley et al. 2023; Kearney and Leigh 2024), and ecosystem functioning (Beugnon et al. 2024; Virkkala et al. 2024) under climate change.

Here, we review how and why microclimate change differs from macroclimate change, and explore the drivers and feedbacks underlying these dynamics. We focus on three key pathways of microclimate change (Figure 2): (I) changing meteorological conditions; (II) physiological, structural and compositional vegetation responses to climate change; and (III) feedbacks between microclimate and climate change. While our primary focus is on microclimate temperature, we also discuss other key microclimate components—soil moisture, air humidity and radiation—as these influence many ecological processes and also interact with temperature. We deliberately exclude direct anthropogenic land-use and land-cover change to isolate the effects of climate change on microclimate, while acknowledging that these human-driven changes remain important. To better capture biologically relevant thermal conditions, we emphasize




FIGURE 1 | Change in microclimate maximum temperatures at Caerthillian Cove, Lizard Peninsula, UK (49.97° N, 5.21° W), from 1983 to 2015. Data were generated using a mechanistic microclimate modeling approach (Maclean 2020). Microclimate temperature changes were derived for each grid cell by linear regression on annual maximum air temperature (5 cm above ground). Interactions between terrain, vegetation, and weather caused local warming rates to diverge from the regional macroclimate trend (0.3°C over the ~3 decades). Three example locations illustrate strong warming (2.9°C), moderate warming (0.7°C), or even cooling (−1.3°C).




FIGURE 2 | Conceptual figure visualizing how this review is structured around three pathways of microclimate change: (I) direct effects of changes in meteorological conditions (red), (II) indirect effects via vegetation changes (green) and (III) feedbacks between microclimate and global climate change (purple, dashed arrows).

daily and seasonal extremes instead of mean temperatures, which can obscure ecologically important variability and asymmetric changes between day and night or between seasons (Helmuth et al. 2010; Cox et al. 2020; Shi and Wu 2025).

To illustrate the mechanisms that drive variation in microclimate change, we ran three different case studies of model simulations using well-validated mechanistic microclimate models (*microclimf* and *micropoint*; Maclean 2026) with air temperatures 5 cm above ground as the response variable. Further details to all case studies are provided in Boxes 3 and 4 and Supporting Information A.

Rather than providing a comprehensive treatment of atmospheric physics, this review is intended as a bridge for ecologists to the physical principles needed to understand, measure, and model microclimate change. In addition, it is intended as a bridge for atmospheric scientists and land-surface modellers to the ecological processes underlying microclimate and macroclimate change. By translating key insights from atmospheric science into an ecological context, and by highlighting that microclimate is strongly shaped by the complexity of the biosphere, we hope to foster stronger cross-disciplinary understanding and ultimately improve projections of biodiversity change and ecosystem management under rapid climate change.

2 | PART I: Changing Meteorological Conditions

In this section, we discuss how changes in temperature, lapse rates, near-surface air temperature inversions, cloud cover, precipitation, humidity, snow cover, and wind alter microclimate landscapes and cause nearby microclimates to diverge (increasing spatial heterogeneity) or converge (microclimatic homogenization) through time. In order to illustrate single mechanisms, we discuss meteorological variables one by one and present their simulated effects on microclimate independently from other variables (Box 3). In reality, however, many of these variables change in parallel (e.g., cloud cover, air humidity, rainfall and soil moisture) and realistic rates of microclimate change can only be predicted by modeling the changes in all variables in concert (Figure 1; Maclean 2020).

BOX 1 | Glossary.

Adiabatic lapse rate: The decrease in temperature of a parcel of air that rises in the atmosphere without exchanging heat or mass with the surroundings. Rising air expands as atmospheric pressure decreases and therefore cools, whereas descending air is compressed and warms.

Leaf Area Index (LAI): The total one-sided leaf area per unit ground area for broadleaf species, or projected hemi-surface leaf area for needleleaf species.

Macroclimate: Climatic conditions as measured by synoptic weather stations in locations chosen to minimise the influence of local surface features such as vegetation, soil properties, and fine-scale topography. The term refers to the general climatic conditions experienced across the broader landscape, rather than local microclimatic variation. Spatial scale: kilometres to continental/global scale.

Mesoclimate: The climate of an intermediate spatial scale, typically covering areas such as valleys, hillsides, or entire landscapes, situated between microclimate and macroclimate. It is impacted by meso-scale topographic features, land cover, distance to coastlines and local weather systems (Barry and Chorley 2009). Spatial scale: ~1 km to several tens or a few hundred km.

Microclimate: The climate in a small area near the Earth's surface that is shaped by local surface features such as vegetation, soil properties, water bodies, and fine-scale topography. The term refers to local climatic conditions that may differ substantially from the surrounding meso- or macroclimate and that often directly influence organisms and ecosystem processes (Barry and Blanken 2016; De Frenne et al. 2025). See also *proximal microclimate*. Spatial scale: ~1 cm to hundreds of metres.

Microclimate buffering: The decrease in amplitude of diurnal or seasonal temperature fluctuations compared to reference macroclimate fluctuations. While microclimate buffering leads to a lower amplitude, for example under tree canopies, microclimate amplification leads to a larger amplitude.

Proximal microclimate: The microclimate close (proximal) to an organism, population or ecosystem of interest (Klinges et al. 2024). While earlier definitions of microclimate refer to near-ground climate at high temporal and spatial resolution, the term proximal climate emphasizes the importance of biological/ecological relevance (proximity) of the climate to an organism in models linking climate and ecological processes/patterns. For instance, the proximal microclimate for an arboreal frog would be the climatic conditions in the tree canopy, whereas the proximal microclimate for an earthworm would be soil temperature and moisture conditions in the upper 10 cm of the soil.

Vapor Pressure Deficit (VPD): VPD is the difference between the *saturation vapor pressure* (i.e., water vapor pressure in fully saturated air at a given temperature) and the *actual water vapor pressure* in the air. In other words, it quantifies the potential for evaporative water loss (Jones 2014).

2.1 | Macroclimate Temperature

2.1.1 | General Increase of Air Temperature

As macroclimate air temperatures rise, surface temperatures usually also increase. Warmer surfaces emit more longwave

Microclimates are created through exchanges of energy between the ground, water, vegetation, and the air (Figure 3). Incoming solar (shortwave) radiation, that is, sunlight, and thermal (longwave) radiation are partitioned into energy that warms surfaces and the air (sensible heat), drives evaporation and other phase changes (latent heat), or is stored in the ground and vegetation. These energy exchanges determine surface and near-surface air temperatures (Chen et al. 1999; Oke 1987).

Net radiation is the sum of incoming and outgoing radiation, comprising both shortwave radiation and longwave radiation. Radiation is either absorbed or reflected by surfaces or partially transmitted through vegetation, with dense canopy reducing the amount of solar radiation that reaches the ground. Natural surfaces mainly absorb incoming shortwave radiation from the sun and re-emit energy as longwave radiation. Net radiation is typically positive during the day and negative at night but generally exhibits a smaller diurnal range beneath canopies than above them, contributing to microclimate buffering (Baldocchi and Vogel 1996; Ross 1975). Net radiation is further partitioned into sensible heat flux, latent heat flux, and heat storage change (Figure 3). Sensible heat, which results in a change of temperature of a substance, represents the energy transferred from warm surfaces to the overlying air by conduction and convection. Conduction transfers heat across the immediate surface boundary, whereas convection transports heat through air movement driven by buoyancy or wind. Latent heat is the energy absorbed or released during a phase change of water, such as evaporation, transpiration, condensation, freezing or melting, without changing temperature.

The rate of sensible and latent heat exchange depends on temperature or vapor pressure gradients and on wind speed, which enhances turbulent transport and thereby increases the exchange of heat and moisture between the surface and the atmosphere (Lewis and Randall 1923). These exchanges occur within the atmospheric boundary layer, the lowest part of the atmosphere that is directly influenced by the Earth's surface through exchanges of heat, moisture, and momentum (Oke 1987; Stull 1988). Latent heat transfer is additionally regulated by stomatal opening in response to radiation, vapor pressure deficit (VPD), temperature, CO₂ concentration, and plant water status. Wind speeds are generally lower near the ground because of surface drag and are strongly influenced by terrain and vegetation. Thus, in forest understories and other sheltered environments, sensible and latent fluxes are reduced due to lower wind speeds and suppressed turbulence, contributing to the thermal decoupling of below-canopy microclimates from above-canopy macroclimatic conditions (Baldocchi and Vogel 1996; Finnigan 2000).

These fluxes are temperature-dependent, and surface temperature can thus be estimated by assuming an equilibrium state is reached. In reality, the amount of heat stored in the ground increases during the day and decreases at night. Similarly, soils generally gain heat in spring and summer and lose heat in autumn and winter, causing subsurface temperatures to lag behind changes in air temperature. Energy storage in soil, vegetation and water forms an important component of the surface energy balance because not all absorbed energy is immediately transferred as sensible or latent heat, delaying warming and cooling over daily and seasonal timescales (Choudhury et al. 1987). Soil water content has a pronounced effect on rates of soil heat storage as wetter soils have higher thermal conductivity and specific heat capacity, and thus wet soils will typically store more heat, potentially damping temperature fluctuations near the soil surface (Campbell and Norman 2012).




FIGURE 3 | Main components of the energy exchange between the Earth's surface, vegetation, and atmosphere that shape microclimate variation across space and time (here for simplicity ignoring lateral heat transfer, such as cold-air pooling). Microclimate conditions arise from the balance between incoming shortwave (SW) radiation and outgoing longwave (LW) radiation, and from the partitioning of the net radiation into sensible heat, latent heat, and heat storage change. Net radiation is usually positive during the day (a) and negative during the night (b). The radiation balance and energy fluxes are influenced by meteorological conditions—such as clouds or air humidity—and by land surface features—such as topography, vegetation structure, and soil moisture—thereby mediating local microclimatic variability and its divergence from broader macroclimate conditions.

(Continues)

Beyond localised energy fluxes, mesoclimate-scale processes shape fine-scale climatic variation through the lateral movement and mixing of air across landscapes (Barry and Chorley 2009). Elevation affects temperature via the **adiabatic lapse rate**, whereby air cools with altitude as pressure decreases (Figure 4a; Barry 2008; Lookingbill and Urban 2003). On clear, calm nights, dense air can accumulate in topographic depressions, producing cold-air pooling, temperature inversions and frost pockets (Figure 4b). Proximity to water bodies also moderates temperature extremes, as water's high heat capacity slows heating and cooling, with inland influence depending on wind, topography, and surface roughness. Thus, both meso- and micro-scale processes generate complex spatial patterns in near-surface climate and warming rates.




FIGURE 4 | Conceptual figure of different rates of warming caused by changes in lapse rate and cold-air pooling/inversion. (a) The black line shows a normal lapse rate, which is typical for the mixed troposphere. The right grey line represents a steeper lapse rate (i.e., temperatures decline faster with height), which can occur under conditions of enhanced surface warming (e.g., due to drying). As the lapse rate steepens, more of the warming is concentrated near the surface, leading to stronger warming at low elevations. This illustrates how identical atmospheric warming can produce different surface warming rates. The opposite pattern has also been observed, where mountain tops are warming faster than valleys (Pepin et al. 2025). (b) The black line shows a typical temperature profile during an inversion. If the conditions under which inversions form become less frequent (grey line), sites in different elevations may experience different rates of warming.

radiation, which acts as a negative radiative feedback by partly counteracting further surface warming. The extent to which warming is buffered or amplified in near-ground microclimates depends on how available energy is partitioned among sensible, latent, and ground heat fluxes. Rising temperatures also increase vapor pressure deficit (VPD) and consequently potential evapotranspiration (Clapeyron 1834; Clausius 1850), although actual evapotranspiration depends on soil moisture and vegetation response (Seneviratne et al. 2010). In moist sites, evaporation and transpiration can increase with increasing temperature and VPD, and latent heat flux further slows down surface warming. However, enhanced evaporation and transpiration also accelerate soil drying. As VPD rises and stomata close to limit water loss, evapotranspiration declines and a larger share of energy is converted to sensible heat, leading to accelerated surface warming. These mechanisms thus create potentially divergent warming trajectories between wet and dry microenvironments (Seneviratne et al. 2010; Srivastava et al. 2024). For example, García-García et al. (2023) found that soil temperatures during annual heat extremes were rising faster than air temperatures (on average by 0.7 °C/decade) in two-thirds of their European measurement sites. They attributed this faster rate to declining soil moisture and associated heat-amplifying feedbacks, such as reduced latent heat transfer and the lower specific heat capacity of dry soil. Additionally, land generally warms 60% faster than oceans (Wallace and Joshi 2018) and is predicted to continue to do so (Byrne and O'Gorman 2018), causing coastal regions to warm more slowly than inland areas (Al-Ruzouq et al. 2022; IPCC 2023a). In both cases, the difference between wet and dry sites or the difference between land and water can potentially

lead to different rates of warming, increasing micro- and meso-climate variability across landscapes.

2.1.2 | Lapse Rates and Their Role in Mesoclimate Warming Patterns

The adiabatic lapse rate—the rate at which air temperature decreases with altitude—plays a critical role in determining how warming is distributed vertically through the atmosphere and how much of that warming is expressed at the surface. In the troposphere, the lower layer of the atmosphere, temperature typically declines with altitude as rising air expands and cools under lower pressure. The actual rate of decline, however, depends on the water vapor content of the air: in completely dry air, the adiabatic lapse rate approaches 9.8°C/km, while in moist air condensation releases latent heat and reduces the rate of cooling with altitude. Because water vapor content varies across regions and over time, lapse rates are not uniform (Minder et al. 2010; Rolland 2003), and this means that neighboring sites, for example, those at different elevations, can experience different rates of microclimate warming even under the same regional macroclimate (Figure 4a). For instance, Chan et al. (2024) used the laws of thermodynamics to map the adiabatic lapse rate across all mountain regions globally. By combining this global map with data on temporal rates of change in mean annual temperature during the period 1971–2020, they mapped the vertical velocities of isotherm shifts across mountain regions. Reported values of vertical climate velocities ranged from 1.8 to 10.8 m/year.

These spatial, and potentially temporal, variations in the lapse rate can either speed up or slow down surface warming, depending on the region, season, and local climatic conditions (Figure 4a). In the Mediterranean, stronger near-surface summer warming has partly been attributed to steeper lapse rates associated with drier conditions. Modeling experiments show that not accounting for lapse-rate changes reduces projected end-of-century Mediterranean summer warming by about 1°C (Brogli et al. 2021).

Changes in the lapse rate contribute to elevation-dependent climate change. Globally, mountain tops warm on average 0.21°C per century faster than nearby lowlands, even if free-air warming is spatially uniform (Pepin et al. 2025). However, regional exceptions occur. Additionally, in areas with complex topography, vertical mixing of air masses becomes more important, and altered lapse rates can influence the stability of the lower atmosphere, affecting both temperature inversions and the likelihood of cold-air pooling (see below and Figure 4b). Therefore, changes in the lapse rate are a key atmospheric mechanism contributing to spatial heterogeneity in warming rates at mesoclimate scales.

2.1.3 | Temperature Inversions and Cold-Air Pooling

Temperature inversions—where air temperature increases with altitude—are an important mechanism influencing near-ground temperatures. Surface inversions typically form during calm, clear-sky nights when strong radiative cooling of the ground cools the adjacent air, resulting in a layer of dense cold air beneath warmer air aloft. Although inversions may also arise from advective or synoptic processes, nocturnal radiative cooling dominates near-surface inversions relevant to microclimates. The strength (temperature gradient) and depth (vertical extent) of these surface inversions determine how much they insulate the ground from warmer overlying air. Climate change is altering the frequency and characteristics of these inversions. A global trend toward shallower inversions, with regional level variable changes in inversion strength has been observed between 1989 and 2019 (Zeng et al. 2022). In the Arctic, a particularly strong decline in inversion depth has been observed and projected in winter months—due to the combined effects of sea ice loss and increased cloud cover, both of which reduce surface radiative cooling (Ruman et al. 2022; Zhang, Furtado, et al. 2021). According to Ruman et al. (2022), the largest wintertime changes in inversion strength are projected to occur over the Arctic Ocean, with increases of roughly 3°C–4°C under the RCP4.5 scenario by 2070–2099. Over Arctic land areas, the projected changes are notably smaller.

More localized near-surface air temperature inversions are caused by cold-air drainage and pooling, a process in which cooler, denser air flows downslope under gravity and accumulates in surface depressions (Figure 4b). This pooling can lead to strong and frequent near-surface inversions, particularly in mountainous or forested areas (Daly et al. 2010). Cold-air pooling is therefore best understood as a specific, landscape-driven subset of surface inversions, one that is tightly coupled to local topography and land cover. For instance, under forest canopies, downslope cold-air drainage can persist during the daytime thanks to the

decoupling between canopy and sub-canopy microclimatic conditions (Pypker et al. 2007), while in treeless areas, cold-air drainage occurs typically only during the nighttime. Rupp et al. (2020) demonstrated that vertical temperature gradients within a mountain valley ranged between –10 °C/km and over 60 °C/km, with substantial diurnal and seasonal fluctuations.

The occurrence and strength of cold-air pooling—and hence its influence on microclimate—is sensitive to climate change. Decreasing cloudiness may promote more frequent and intense cold-air pooling due to enhanced night-time cooling, while increasing cloud cover can suppress it. A recent process-based analysis further suggests that greenhouse-gas-induced warming may weaken the nocturnal cold-air drainage by at least 10% under a high-emissions scenario (RCP 8.5), with the largest reductions expected in warm and humid regions (Rupp et al. 2021). Similarly, changes in wind speed and direction can either disrupt or enhance cold-air pooling events depending on local exposure and terrain configuration (Miró et al. 2018). These interactions mean that cold-air pooling-prone areas, such as forested slopes and closed basins, may experience slower rates of warming compared to adjacent open sites (Daly et al. 2010; Pastore et al. 2022). Conversely, weakening inversions due to increased cloudiness or reduced snow cover (as cold air forms over snow patches) may accelerate warming (Figure 4b). In this way, shifts in the frequency, strength, and spatial distribution of temperature inversions can contribute to increasing the heterogeneity in microclimate warming rates across the landscape.

Current evidence for different rates of climate warming due to cold-air pooling is mixed. Using 45 years of observational data from multiple meteorological stations in a forested mountain watershed, Jones et al. (2025) found little variation in multi-decadal climate trends among sites, despite frequent occurrences of cold-air pooling and temperature inversions. Hiebl and Schöner (2018), on the other hand, exploited a gridded observational dataset of daily minimum temperature across Austria over the 1961–2017 period and found that winter inversions had declined in frequency by 11%, leading to accelerated warming rates at low elevations. This illustrates the strong context dependency and therefore the need to better understand how changes in mesoclimatic conditions affect microclimatic changes over longer periods.

2.2 | Clouds

Changes in cloud cover—including timing, altitude, and extent—strongly affect local radiation balances, thereby shaping microclimate change patterns. On clear summer days with high incoming solar radiation, temperature contrasts between adjacent microclimates peak due to spatial differences in shortwave radiation (e.g., Lehtinen et al. 2026). Low-altitude clouds, like stratus and cumulus, block incoming solar radiation, thereby reducing spatial temperature contrasts near the surface. Observed and projected trends show that global average daytime cloud fraction (particularly low-level cloudiness) decreases faster than night-time cloud fraction (Luo et al. 2024b), but that there is a large variability across regions and seasons, as well as a high uncertainty in predictions due to cloud-atmosphere feedbacks (Hou et al. 2021).




FIGURE 5 | Reductions in daytime cloud cover cause nearby microclimates to diverge over time, increasing microclimate variance across space. With decreasing cloudiness, open meadows warm faster than forest interiors (a) and sun-exposed slopes warm faster than shaded slopes (b). (c) Warming trajectories of adjacent microclimates with different warming trends (yellow vs. orange triangle, meadow vs. forest interior in a and sun-exposed vs. shaded slope in b). (d) Visualizes a possible scenario where macroclimate temperature increases and cloudiness decreases over three time steps (now, +5 years, +10 years), thereby increasing microclimate contrasts over short distances (yellow vs. orange triangles).

Reductions in daytime cloud cover such as those reported over Europe, where cloud cover has declined by roughly 1.5%–2.0% between 1980–1995 and 2000–2015 (Dong et al. 2022), increase the amount of sunlight reaching the surface (Sfică et al. 2021). This accelerates warming on sun-exposed slopes compared to slowly-warming pole-facing slopes, increasing their contrast in near-ground temperature (Chesnoiu et al. 2025; Figure 5; see also Case Study 1, Box 3, Figure 6e). The effect also occurs in moderately complex terrains at low sun angles, (Guthrie 2001), or across canopy cover gradients (Grimmond et al. 2000; Johnson and Smith 2008). The magnitude of this asymmetry depends on cloud thickness, sun angle, and surface albedo (e.g., from snow, soil, or vegetation).

Clouds also play a key role at night. While daytime temperature contrasts are largely driven by incoming solar radiation, nighttime cooling is governed by the loss of longwave radiation from the surface to the atmosphere. Clouds absorb this outgoing longwave radiation and—because they are efficient emitters of thermal radiation—emit part of it back toward the surface, reducing radiative cooling, inhibiting cold-air formation and pooling in topographic depressions (Figure 5) and generally homogenizing temperatures across a landscape (Figure 6d). For example, Groot and Carlson (1996) found the difference of near-ground night temperatures between open environments and under shelterwood to be larger in calm clear-sky nights compared to nights with an overcast sky.

2.3 | Precipitation

One facet of climate change is shifts in the amount, timing, frequency, and intensity of precipitation (IPCC 2023b) and, linked to that, the amount of moisture in the soil and humidity in the air. While the impact of changes in precipitation on soil moisture may depend strongly on climate regime, topography, vegetation, and soil type, more precipitation generally leads to increases in soil moisture (Cowley et al. 2017; Litaor et al. 2008;

Sehler et al. 2019). Soil moisture is an important driver of near-ground air and soil temperatures, especially under warm conditions (Ashcroft and Gollan 2013; von Arx et al. 2013). When soils are dry, most of the incoming solar energy is converted into sensible heat, warming the soil surface and the overlying air, rather than being used for evaporation. Conversely, when soils are wet, a larger fraction of the available energy is converted into latent heat fluxes through evaporation. Soil moisture also changes how heat moves through the soil as well as how much heat a soil can store. Compared to dry soils, moderately moist soils conduct heat rapidly away from the surface, while fully saturated soils act as substantial energy stores due to water's high heat capacity. Both mechanisms effectively buffer near-surface soil temperature fluctuations, though saturated conditions provide the strongest temperature buffering at the surface (Campbell and Norman 2012). Consequently, daytime near-ground air temperatures are often lower and night-time temperatures higher over wet soils than over dry soils (Davis et al. 2019; García-García et al. 2023). These effects will, however, depend on the ecosystem's soil moisture levels since increasing precipitation will have larger effects on microclimate in drylands compared to more mesic habitats (Korell et al. 2021). Additionally, soil moisture has been shown to limit evapotranspirative cooling stronger in open ecosystems compared to forest ecosystems, which has caused daytime summer land surface temperature to increase faster in open grasslands and croplands and more slowly in forests with a difference of $-0.19^{\circ}\text{C}/\text{decade}$ (Li et al. 2025).

The timing of rainfall can also be important for microclimatic conditions. For example, increased summer rainfall generally cools soils by increasing latent heat fluxes and heat capacity (Dai et al. 1999; Mengistu et al. 2017). In contrast, winter rainfall often has smaller and more variable effects on soil temperature. In regions where soils remain unfrozen, increased winter rainfall raises soil moisture, allowing heat to be transferred and stored more efficiently in the soil, which can result in warmer winter soils. Areas with pronounced rainfall seasonality may

therefore experience stronger seasonal changes in microclimatic buffering than regions with year-round precipitation. Future changes in precipitation variability remain uncertain, adding further uncertainty to predictions of microclimate change (van der Wiel and Bintanja 2021; Zhang, Ding, et al. 2021).

While changes in rainfall are often considered, other forms of precipitation also exhibit temporal trends and influence microclimatic conditions. For example, there is a global reduction in fog (Klemm and Lin 2016) and dew (Dou et al. 2021), though with regional exceptions (Lavigne and Liu 2022). Particularly in arid systems, regular fog events decrease evapotranspiration (thus latent heat fluxes) and drought risk (Baguskas et al. 2016), while also shaping microclimate conditions by attenuating solar radiation, increasing relative humidity, and buffering thermal extremes (Sotomayor and Drezner 2019). Fog buffers minimum temperatures (Zhang et al. 2014), and a decrease in fog would likely lead to higher microclimate variability, including lower minimum temperatures.

In cold environments, changes in the depth, timing and duration of snowpacks can have large impacts on microclimates, particularly immediately above and below the snow layer. Due to snow's low thermal conductivity, it forms a strongly insulating layer, decoupling soil and surface temperature from free atmospheric temperature (Aalto et al. 2018). Under snow cover, surface temperatures cannot exceed 0°C because surplus energy input is absorbed as latent heat during melting, effectively buffering the surface against further warming. Therefore, even a thin layer of snow effectively decouples near-ground maximum temperatures from macroclimate maximum temperatures and homogenizes microclimates across a landscape (See also Case Study 1, Box 3, Figure 6k). Minimum temperatures (below freezing point) below the snow are closely linked to snowpack thickness, with greater snow depth increasing microclimatic buffering up to a threshold depth (often 15–40 cm; with more compact snow having a lower buffering capacity), beyond which additional snow has limited impact (Wilson et al. 2020; Zhang 2005). Snowpack surfaces are characterised by a higher albedo and, as a result, generally lower near-surface maximum temperatures than adjacent snow-free microsites, with this effect being strongest at high sun angles (Zhang 2005). During low sun angles, on the other hand, experiments show that snow removal reduces winter soil temperatures, as frost penetrates more easily into exposed grounds (Zhao et al. 2022; Figure 6j). During snowmelt, sites with thicker snowpack also extend the seasonal soil temperature stability, reducing near-surface frost events, and maintaining higher soil moisture. Thus, the general trend for reduced snowfall and shorter snowpack duration (see, e.g., IPCC 2023b), is likely to alter microclimates year-round.

2.4 | Wind

Considerable changes in wind speed and direction have been observed in recent decades, with global patterns showing marked heterogeneity (McVicar et al. 2012). Terrestrial systems have generally experienced wind-stilling (i.e., a decrease in mean wind speed), but trends and projections vary by region

(Zha et al. 2021) and may reflect changes in large-scale ocean–atmosphere circulation patterns, such as the North Atlantic Oscillation (Zeng et al. 2019). Wind affects the coupling between ambient and near-surface temperature or humidity by changing the atmospheric mixing close to the ground. Stronger winds enhance convection (see Box 2) as well as vertical and horizontal air mixing, leading to reduced microclimatic heterogeneity and a tighter coupling of micro- to macroclimate fluctuations (Ashcroft et al. 2009; Baker et al. 2021; Lehtinen et al. 2026; Figures 6f,g and S1g–i). However, this wind-induced homogenization is reduced in forested and topographically complex landscapes, where terrain sheltering and surface roughness limit air mixing (Renaud et al. 2011; Wever 2012). Thus, increased windiness is expected to reduce microclimate variability most strongly in flat and open landscapes. In our model scenario, increased wind speed generally cooled maximum temperatures (in agreement with, e.g., Riddell and Porter 2025), but this cooling effect was weaker under forest canopies compared to open grasslands (Figures 6g and S2g–i). In contrast, windstilling is expected to increase microclimatic extremes by reducing atmospheric mixing, leading to more frequent cold air pooling (Dobrowski 2011) and more extreme thermal minima (Geiger 1966). Under windstilling, reduced atmospheric mixing also leads to thicker leaf/surface boundary layers (i.e., the thin layer of air directly adjacent to a surface), decreasing evaporation and likely increasing near surface humidity (Jones 2014).

Shifts in wind direction (McInnes et al. 2011) may lead to a spatial redistribution of wind-sheltered and wind-exposed microsites, with microsites that become more sheltered from wind potentially experiencing, for example, more extreme temperatures, greater dew accumulation and lower evaporation (Ashcroft et al. 2009; Geiger 1966; Maclean et al. 2017). Climate models show a large regional variability in projections in annual average and seasonal wind speeds (McInnes et al. 2011), and if any unidirectional long-term trends emerge at all, they are likely very region-specific.

3 | PART II: Changes in Vegetation

Vegetation strongly influences microclimate buffering through its structural and physiological characteristics, including canopy height, Leaf Area Index (LAI) and stomatal conductance, that is, how easily water vapor passes through the stomata on leaf surfaces (De Frenne et al. 2021). Here we first consider physiological and phenological responses while community composition remains constant, before discussing structural and compositional changes.

3.1 | Functional Responses

3.1.1 | Physiology

Without changing community composition, climate change can alter vegetation physiology, thereby modifying sub-canopy microclimates. In temperature-limited systems such as boreal forests and other high-latitude or high-altitude ecosystems, moderate warming can enhance primary productivity (Piao,

Wang, et al. 2019). Rising atmospheric CO₂ concentrations may further promote plant growth (Dong et al. 2025; Forkel et al. 2016; Mndela et al. 2022; Zhan et al. 2024), with elevated-CO₂ experiments reporting above-ground biomass gains in forests of 21.4% on average (Baig et al. 2015). Nevertheless,

this “carbon fertilization effect” is often constrained by nutrient and water availability (Ellsworth et al. 2017; Peñuelas et al. 2017; Terrer et al. 2019). Elevated atmospheric CO₂ is expected to reduce stomatal conductance, limiting transpiration and increasing leaf and canopy temperatures, with a thermal

BOX 3 | Case Study 1 on effects of meteorological conditions.

Here, we present the model results for Case Study 1 showing minimum (Figure 6d,f,h,j,m) and maximum (Figure 6e,g,i,k,l) near-surface temperatures 5 cm above ground. A selection of individual meteorological variables (cloud cover, wind speed, VPD, and snow cover) were modified independently to visualize their influence on microclimate temperatures and the spatial variability of these effects across a ca. 2 × 2 km² landscape at the border of Parque Natural Posets-Maladeta in the Spanish Pyrenees (see Supporting Information A for modeling details). For each simulation, we modelled the minimum temperature of the 10 coldest days and the maximum temperature of the 10 warmest days per year by setting the focal meteorological variable to low and high values (with values respectively halved or doubled) while keeping all other variables constant. The simulations revealed substantial differences in the magnitude of microclimate responses among atmospheric drivers and between minimum and maximum temperature metrics (Figures S1, S2, and 6, Table S1). Although the outputs of a mechanistic model that integrates multiple complex and non-linear processes are not always straightforward to interpret, our objective was primarily to illustrate that changes in atmospheric conditions can produce highly heterogeneous microclimate responses across the landscape. This spatial heterogeneity is evident from the patchy temperature difference maps (Figure 6d–k), which demonstrate that identical changes in meteorological forcing do not translate into uniform microclimate temperature changes across space. As mentioned earlier, we ran the simulations in a way to isolate the effect of single atmospheric variables, which in reality do not change in isolation, sometimes complexifying model interpretation. Moreover, because the *microclimf* model does not account for mesoclimatic processes such as cold-air pooling, the spatial complexity of temperature patterns and trends in real-world landscapes is likely even greater than suggested by our simulations.

Minimum Temperature

Average changes in minimum temperature were generally below 0.2°C, except for the effect of snow (Figure 6l,m, Table S1). With low cloud cover, heterogeneity in minimum temperatures was created among others by dense vegetation, which prevented the ground from cooling down as it captured emitted thermal radiation from the ground and gradually released the heat stored during the day. Under high cloud cover, almost all upwelling emitted thermal radiation was emitted back downward irrespective of vegetation cover or terrain position (Figures 6d and S1d–f).

Higher wind speeds at night enhanced turbulent mixing, reducing the build-up of cold air near the surface and thereby raising minimum temperatures, particularly where wind speeds increased the most (Figures 6f and S1g–i).

Increasing VPD (drier air) led to slightly lower minimum temperatures (on average −0.2°C, Table S1) because the dewpoint temperature dropped, meaning the air could cool down more before condensation began. When the dewpoint was reached, condensation released latent heat, which slowed further cooling. Thus, drier air typically allows temperatures to fall a little lower overnight, yet in heterogeneous ways across the landscape (Figures 6h and S1j–l).

The most pronounced effect on near-ground minimum temperatures was caused by snow cover, which was almost by an order of magnitude larger than the effects of wind or air humidity/VPD with an average temperature change of 3°C (Figure 6j,m, Table S1). Snow cover had an insulating effect on near-ground minimum temperatures, protecting the ground from frost and therefore leading to a general warming. However, the insulating effect depends on snow depth, density, moisture content, or soil heat flux and could therefore produce slight temperature gradients across the landscape (Figures 6j and S1m–o).

Maximum Temperature

Maximum temperature effects were overall larger compared to effects on minimum temperature. Spatial heterogeneity in maximum temperatures near the ground was higher under low cloud cover as the topographic and vegetation shading effects became more pronounced. When cloud cover increased, maximum temperatures generally decreased due to reduced energy input from the sun (on average by −12.4°C, Table S1), and this cooling effect was strongest in open environments (Figures 6e and S2d–f).

Wind had a cooling effect on maximum temperatures (on average −5.6°C, Table S1) as it increased turbulent mixing and transported heat away from the surface. This cooling effect was particularly strong on the open south-east facing slopes—exposed to the mainly southerly winds in this area during the day—and less so under forest canopies which sheltered the ground from wind (Figures 6g and S2g–i).

Increasing VPD (drier air) had a cooling effect on maximum temperatures (on average −4.2°C, Table S1), as there was more evaporative cooling near the surface. The absolute effect varied throughout the landscape as a function of slope and aspect (which drive incoming energy), soil moisture, and vegetation height and density (Figures 6i and S2j–l).

When snow was present, maximum temperatures were effectively capped at around 0°C because additional energy input was used for melting the snow (latent heat of fusion) rather than further warming (Figures 6k and S2m–o). Depending on the location in the landscape, this resulted in a cooling effect ranging from −34.6°C to −4.1°C (Table S1).

(Continues)




FIGURE 6 | Model results for Case Study 1 in the Spanish Pyrenees showing the effects of changes in individual meteorological drivers on near-surface temperature (5 cm above ground). Panels d, f, h and j show changes in minimum temperature, while panels e, g, i and k show changes in maximum temperature, resulting from experimentally altering cloud cover, wind speed, air humidity (expressed as vapor pressure deficit, VPD), and snow cover, respectively. These difference maps (d–k) show the temperature change obtained by comparing simulations with low and high values of each meteorological variable (halved versus doubled values), while holding all other variables constant, providing a measure of the sensitivity of microclimate to each driver across the landscape. Panels l and m summarize the distributions of temperature changes shown in the maps for maximum and minimum temperatures, respectively (see Table S1 for summary statistics). Small triangles indicate locations used to illustrate differences in the magnitude of vegetation-mediated microclimate change. Map lines delineate study areas and do not necessarily depict accepted national boundaries (Icons from Flaticon.com).

effect of 1.3°C quantified in a mature oak canopy (Hagan Brown et al. 2025).

Climate change and the associated increase in climatic extremes can trigger rapid physiological responses, including thermal acclimation and tighter stomatal regulation (Bertolino et al. 2019; Liu et al. 2024; Wahid et al. 2007). These responses can help maintain plant function and canopy cover during heat stress, although reduced stomatal conductance and transpiration may simultaneously limit evaporative cooling and weaken microclimate buffering (Fernández et al. 2017; Wright and Francia 2024). Under severe heat or drought stress, reduced soil water availability and stomatal closure can increase canopy temperatures and weaken evaporative cooling (Grossiord et al. 2020; Slot et al. 2024). Once critical temperature thresholds are surpassed, leaf mortality and crown dieback can occur, causing a partial or complete breakdown of microclimate buffering (Doughty et al. 2023; Kopáček et al. 2020; Jump et al. 2017; von Arx et al. 2013). Nonetheless, recent studies have shown that elevated transpiration rates can persist during heatwaves even when carbon uptake ceases, under both well-watered and drought-stressed conditions (Marchin et al. 2023; Posch et al. 2024). This decoupling between transpiration and photosynthesis may allow thermal regulation during heat stress. Over longer time scales, enhanced productivity may also increase vegetation height, canopy density and LAI, linking physiological responses to the structural changes discussed below.

3.1.2 | Phenology

On interannual time scales, climate change has been shown to alter phenological cycles including leaf onset, flowering time and leaf senescence across a wide range of species and ecosystems (Badeck et al. 2004; e.g., Inouye 2022; Piao, Liu, et al. 2019). Warmer winters and springs generally advance leaf onset (Cai et al. 2025; Piao, Liu, et al. 2019), thereby reducing the rate at which spring microclimates change relative to macroclimate. In a mountainous region of Oregon (USA), years characterized by conditions resembling projected future climates—reduced snowpack and more frequent high-pressure systems—not only advanced spring phenology but also increased phenological synchrony across elevation gradients, thereby reducing microclimatic heterogeneity (Ward et al. 2018). Autumn responses are more variable: warmer autumns delay senescence, whereas drought often accelerates it (Wu et al. 2022; Zhang et al. 2025; Zohner et al. 2023). Consequently, growing-season responses to climate change vary among ecosystems. For example, substantial warming across China between 1960 and 2009 led to earlier leaf emergence by 5.4 days and delayed autumn leaf coloring by 4.6 days in four woody plant species, resulting in an overall growing-season extension of approximately 10 days (Dai et al. 2014). In our Case Study 2, we showed how earlier leaf emergence in spring increased minimum temperatures and decreased maximum temperatures, mainly through changes in radiation: increased canopy cover reduced incoming shortwave radiation during the day and reduced night-time longwave radiative heat loss (Box 4, Figure 7).

Vegetation phenology can also feed back to the climate system through biogeophysical mechanisms such as changes in

evapotranspiration, albedo and cloud formation (see section III). In a recent synthesis, Winkler et al. (2026) showed that a 10-day extension of the growing season could lead to a global surface cooling of $0.10^{\circ}\text{C} \pm 0.03^{\circ}\text{C}$ through changes in surface albedo and energy balance. However, the same study highlighted that increased spring transpiration and evaporative cooling may subsequently enhance summer soil drying, potentially offsetting some of this cooling effect through increased sensible heat fluxes.

3.2 | Structural and Compositional Changes

3.2.1 | Vegetation Structure

Structural changes in vegetation can arise both through physiological responses within existing communities and through shifts in species composition. Increased productivity under warming and elevated CO_2 may gradually produce taller and denser canopies with higher LAI, potentially increasing microclimate buffering, that is, warming night-time minimum and cooling daytime maximum temperatures. Because this relationship is non-linear, buffering changes most strongly at low LAI (Case Study 2, Box 4; Zellweger et al. 2019). Changes in canopy height and density also modify air movement and turbulent exchange, thereby influencing the transfer of heat and moisture between the vegetation and the atmosphere. In microclimate models, these effects can be represented through aerodynamic properties of the canopy.

A well-documented example of vegetation-mediated microclimate change is the “shrubification”, that is, the expansion of woody species into previously treeless ecosystems, such as tundra (Sturm et al. 2001) or savannas (Devine et al. 2017; Case Study 3, Box 4). Shrub expansion increases vegetation height, leaf area, transpiration and shading, buffering daytime temperatures while often increasing winter soil temperatures through snow insulation (Finne et al. 2023; Kemppinen, Niittynen, Virkkala, et al. 2021; Kropp et al. 2021; Mekonnen et al. 2021; Myers-Smith and Hik 2013; Wilcox et al. 2024). More generally, plant species influence microclimatic conditions through their structural characteristics, which can be partly described by traits such as canopy height, cover, and shade-casting ability (Gril, Spicher, et al. 2023; Meeussen et al. 2021). Microclimatic effects are strongest when replacement species differ markedly in stature or phenology, for example, when shifts occur between grasslands to shrublands or from evergreen to deciduous dominance. For instance, in low-growing shrublands, non-native trees lowered near-surface air temperatures by increasing shading, whereas dense invasions raised near-surface temperatures by suppressing native ground cover and exposing more bare ground (Garcia and Clusella-Trullas 2019). Furthermore, forest disturbances such as droughts, pest insects, fires and storms reduce canopy buffering and cause rapid increases in daytime summer sub-canopy temperature (e.g., Greiser et al. 2025). They can effectively erase decades of thermal decoupling, exposing the understory to thermal conditions equivalent to several decades of macroclimate warming (Haesen et al. 2026). Increasing structural diversity can have the opposite effect. In a large-scale tree diversity experiment in China, near-surface temperatures were up to 4.4°C lower in 24-species mixtures than in monocultures through greater canopy density and structural complexity (Schnabel et al. 2025). Likewise, increased species diversity can enhance canopy density and rooting depth diversity, leading

Case Study 2: Long-term shifts in leaf phenology

To assess how variation in leaf phenology influences microclimate buffering, we simulated near-surface air temperature beneath a forest canopy in a mixed, predominantly deciduous forest in New Hampshire, USA. To isolate the effects of phenology, the observed mean monthly values of the leaf area index (LAI) in April between 1982 and 2019 were used as an input to model forest understory monthly minimum and maximum temperature, all else—including macroclimate—being held constant (see Supporting Information A for modeling details). Between 1982 and 2019, the study forest observed an average increase in April LAI of 0.104 per decade from 0.679 to 1.054 (Figure 7a), raising modelled minimum temperature from -7.1°C to -6.9°C (Figure 7b) and reducing maximum temperature from 16.7°C to 16.5°C (Figure 7c), with strongest changes at low LAI (Figure 7e,f). Earlier leaf emergence increased buffering by altering both shortwave and longwave radiation, reducing daytime warming and night-time cooling.

Case Study 3: Shrub and tree encroachment

We modeled the effect of increased tree and shrub cover on minimum and maximum near-ground air temperature in a $1 \times 1 \text{ km}^2$ grassland area in the Central Anatolian Steppe of Niğde Province, Western Turkey. Invasion of shrubs and trees was modeled by superimposing structural vegetation data, namely vegetation height (Figure 8a,b) and LAI, from a nearby steppe-woodland site onto a digital terrain model of the grassland site. Minimum and maximum temperatures (at 5 cm above ground) refer to the average of daily minimum and maximum temperatures of the years' 10 coldest or hottest days, respectively (Figure 8e,f,h,i).

The modelled change in vegetation cover generally led to warmer near-ground minimum temperatures (on average by 0.2°C) and cooler maximum temperatures (on average by 11.9°C , Table S2). However, microclimate change varied across the study area as a function of vegetation height and leaf area index (Figure 8g,j). For example, we observed changes of maximum temperature ranging from -16.6°C (cooling) under tall dense shrubs and trees to $+6.0^{\circ}\text{C}$ (warming) in open patches where grass height was modelled to become shorter (Table S2, Figure 8d,g). Minimum temperature changes varied less than maximum temperatures and ranged from -2.5°C under invading shrubs and trees to 1.5°C in patches with shorter grass (Table S2, Figure 8d,j). Similar to Case Study 2, these temperature changes mainly resulted from changes in the radiation balance at day and night.




FIGURE 7 | The influence of changes in leaf phenology on microclimate change in a mixed forest in New Hampshire, USA (Case Study 2). Observed mean monthly values of the leaf area index (LAI) in April between 1982 and 2019 (a) were used as an input to model forest understory minimum (b, e) and maximum temperature (c, f), all else—including macroclimate—being held constant. Map lines delineate study areas and do not necessarily depict accepted national boundaries.

(Continues)

BOX 4 | (Continued)




FIGURE 8 | The effect of increased tree and shrub cover on minimum and maximum near-ground air temperature in a grassland area in the Central Anatolian Steppe of Niğde Province, Western Turkey (Case Study 3). (e, f, h, i) Minimum and maximum temperatures refer to average of daily minimum and maximum temperatures of the years' 10 coldest or hottest days, respectively. (g, j) Temperature difference between the grassland and the woodland scenario, negative (blue) values mean temperatures are getting cooler. The difference (microclimate change) maps are summarized in boxplots (d). Both the average magnitude and the spread of microclimate change are larger for maximum than for minimum temperatures (d, see also Table S2 for summary statistics). The small triangles mark example locations with different magnitudes of microclimate change caused by different trends in vegetation cover. Map lines delineate study areas and do not necessarily depict accepted national boundaries.

to higher evapotranspiration rates and stronger microclimate buffering (Huang et al. 2024; Schnabel et al. 2025; Zhang et al. 2022).

3.2.2 | Surface Albedo

Species favored by climate change can differ in albedo, thereby altering the amount of solar radiation absorbed at the surface. For instance, in boreal forests, birch exhibits a higher stand albedo (0.22 ± 0.01) than Scots pine (0.14 ± 0.02) and Norway spruce (0.11 ± 0.03 , Lukeš et al. 2013). Arctic and alpine greening has emerged as a major trend at high latitudes and elevations, driven by warming and a lengthened growing season (Choler et al. 2021; Roland et al. 2024; Rumpf et al. 2022). In these regions, open land is typically snow-covered in winter or lichen-dominated in summer, both characterized by high albedo. However, shrub-dominated communities can have half the albedo values of the lichen *Cladonia stellaris* in mountainous Norway (Aartsma et al. 2020), thereby enhancing microclimate warming through increased absorption of solar

radiation (Mallen-Cooper et al. 2021). Comparing satellite-derived land surface temperature trends between forests and open lands in Europe, Li et al. (2025) demonstrated that decreases in winter snow cover decreased the albedo differences and thereby the temperature differences between forested and open habitats, with open grasslands warming $-0.14^\circ\text{C}/\text{decade}$ faster than forests. Conversely, in warmer regions, rising temperatures and declining precipitation may promote desertification, where reduced vegetation cover increases surface albedo (IPCC 2022).

3.2.3 | Transpiration

Community turnover also alters transpiration through changes in species composition and functional traits. Many species replacing current species assemblages possess morphological and hydraulic traits that enhance their ability to cope with new climatic conditions (Kramp et al. 2022; Kühn et al. 2021). In some alpine meadows, colonizing lowland species exhibit

larger leaves (Debouk et al. 2015; Schuchardt et al. 2023), which increase transpiration rates, elevate air humidity, and reduce temperatures through latent heat exchange—though only in ecosystems where water availability is not limiting. In regions facing more frequent droughts, however, “winner” species are likely to be characterized by short stature, sparse canopies, small and thick leaves, and tighter stomatal regulation. The increasing dominance of such water-conserving species may weaken vegetation-mediated cooling (Michalet et al. 2024; Richter et al. 2022). Deep-rooted drought-adapted species may better maintain canopy cover during prolonged drought (Kühn et al. 2021).

4 | PART III: Vegetation and Microclimate Feedbacks

The processes described in the previous sections not only determine how climate change alters microclimates but can also feed back to influence regional and global climate. At the global scale, climate feedbacks regulate the rate of warming driven by greenhouse gas emissions, whereas at local and regional scales, feedbacks arise through changes in surface energy exchange, hydrology, and vegetation that can influence the atmosphere. We therefore distinguish between global climate-system feedbacks and local and regional land-atmosphere feedbacks that originate at the land surface and are driven by land surface features including vegetation.

At global scales, the primary stabilizing mechanism of the climate system is the Planck feedback, whereby warming surfaces emit more thermal radiation (IPCC 2023a). This negative radiative feedback limits global warming, while local energy exchange processes (Box 2) determine how global warming is expressed in near-surface microclimates. In contrast, warming increases atmospheric water vapor, strengthening the greenhouse effect through a positive feedback. Together with the closely coupled lapse-rate feedback, this contributes approximately $+1.3 \text{ W/m}^2/\text{°C}$ to the global climate feedback (IPCC 2023a). Clouds strongly influence Earth's radiation balance by reflecting incoming solar radiation while also trapping outgoing longwave radiation. The net cloud feedback remains uncertain and varies among cloud types, regions, and seasons (Sherwood et al. 2020). Changes in surface albedo provide another major feedback, with the loss of highly reflective surfaces such as snow and ice, or the expansion of darker vegetation, reinforcing warming (Bright et al. 2014; Evans et al. 2017; Jiao et al. 2017; Pearson et al. 2013; Pecl et al. 2017).

At local and regional scales, land–atmosphere exchanges provide an important pathway by which microclimates feed back to the atmosphere. Vegetation strongly regulates the exchange of energy and water between the land surface and atmosphere (De Frenne et al. 2021; Miralles et al. 2025; Winkler et al. 2026). Vegetation cools leaves and surrounding air through transpiration while increasing atmospheric water vapor, thereby potentially promoting cloud formation (Duveiller et al. 2021). Conversely, vegetation loss caused by drought, repeated heat stress or disturbance can accelerate

surface warming (IPCC 2023a). Besides regulating radiation and evapotranspiration, vegetation structure also alters the aerodynamic roughness of the land surface. Taller and structurally more complex vegetation generally enhances turbulent exchanges of heat, moisture and momentum between the land and atmosphere, influencing boundary-layer development, cloud formation and regional climate (Miralles et al. 2025). Vegetation can further influence regional cloud formation and precipitation through emissions of BVOCs and atmospheric pollen, which contribute to atmospheric aerosols (Dada et al. 2023; Duveiller et al. 2021; Greenberg et al. 2004; Kretzschmar et al. 2024; Rap et al. 2018). Longer-term vegetation–climate feedbacks arise through shifts in vegetation structure and tree-line position, which alter surface albedo, evapotranspiration and carbon storage. Trees and shrubs can facilitate their own persistence and expansion by locally elevating night-time temperatures (D'Odorico et al. 2013), reinforcing these broader climate feedbacks. Deforestation in the Amazon has increased low-altitude cloud cover, producing top-of-atmosphere cooling of up to $6.8 \pm 0.6 \text{ W/m}^2$ in areas of extensive forest loss (Dror and Feingold 2026), illustrating that the local climatic effects of deforestation and its top-of-atmosphere radiative effects do not necessarily act in the same direction.

Beyond these biophysical feedbacks, microclimates also influence climate through biogeochemical feedbacks associated with the carbon cycle (Arora et al. 2020; Song et al. 2019). While these processes are typically represented in coarse-resolution Earth system models, many are influenced by microclimate. One major feedback mechanism is the release of long-stored CO_2 and methane from thawing permafrost, enhancing atmospheric greenhouse gas concentrations and adding an estimated $0.29^\circ\text{C} \pm 0.21^\circ\text{C}$ of warming under RCP8.5 and 0.05°C – 0.15°C under RCP4.5 (Schoor et al. 2022). Similarly, climate change may induce forest dieback, releasing CO_2 into the atmosphere and reinforcing warming (Munday et al. 2025). Conversely, higher atmospheric CO_2 concentrations can temporarily enhance plant growth and carbon uptake (Norby 2025), although this fertilization effect is constrained over time by nutrient limitation, drought stress and shifts toward less carbon-dense vegetation (Peñuelas et al. 2017; Terrer et al. 2019). Likewise, climate-driven transitions from forests and shrublands to more open ecosystems reduce carbon storage and reinforce atmospheric CO_2 increases (Rosan et al. 2019; Venter et al. 2018).

Together, these examples illustrate how local microclimatic processes can influence regional meteorology directly through energy and water exchange, and potentially global climate through vegetation-mediated changes in energy, water, and carbon cycling. While the strongest and most consistent feedbacks operate through surface energy exchange, hydrological processes, albedo change, and carbon storage, many remain context dependent and poorly quantified. Key research priorities include quantifying the strength and timing of these feedbacks, integrating microclimatic processes into biosphere–atmosphere models, and resolving physiological limits, adaptation, time lags, non-linear responses, and regime shifts in vegetation–climate feedbacks (Winkler et al. 2026).

5 | Conclusions and Outlook

This review highlights three overarching conclusions. First, microclimate warming does not necessarily track macroclimate warming, and nearby locations may experience substantially different warming rates, leading to increased or decreased heterogeneity of microclimate temperatures (Figure 1). Second, these divergent warming trajectories and shifts in spatial heterogeneity arise from the combined effects of changing meteorological conditions, vegetation dynamics, and land-atmosphere feedbacks rather than from a single dominant driver (Figure 2). Third, predicting future microclimates therefore requires moving beyond static microclimate-macroclimate offsets and instead accounting for the physical and ecological processes governing microclimate change. Ultimately, this microclimatic perspective may improve predictions of ecological responses and help anticipate climate impacts at scales relevant to organisms and ecosystems.

Although this review focuses largely on trends in mean microclimate change, such averages may obscure ecologically critical shifts in the frequency, intensity, and duration of extreme events—often the primary triggers of biological responses like sudden range shifts (Soifer et al. 2025) or ecosystem tipping-points (Lenton 2013). Importantly, microclimate changes can be spatially heterogeneous, and therefore what constitutes an extreme event at a regional scale (e.g., heat waves or droughts) may be locally amplified or buffered.

Microclimate ecology has recently focused on increasing the spatial resolution of climate datasets, but we argue that it is at least equally important to understand the processes driving their temporal dynamics and how these processes manifest across landscapes and ecosystems. A major unresolved challenge is quantifying turbulent transport of heat and moisture in structurally complex vegetation. This limits our ability to predict how effectively vegetation buffers or amplifies warming and how much structural complexity must be represented in climate models. These uncertainties propagate into our understanding of vegetation-microclimate feedbacks, because vegetation responses themselves depend on fine-scale spatial and temporal variation in microclimate that is often poorly resolved, particularly under novel future climate conditions.

Resolving these uncertainties requires long-term observations linking vegetation change and microclimate across heterogeneous landscapes. The growing availability of long-term microclimate records and the increasing length of these time series (Lembrechts et al. 2020) hold great value for capturing fine-scale spatiotemporal patterns (Haesen et al. 2026). Microclimate observation can also strategically prioritize data collection that supports the development, refinement, and validation of mechanistic models—such as those we used in this paper—building on available open-source approaches (Buckley et al. 2023; Kearney and Porter 2017; Maclean 2026). Grounded in physical principles, mechanistic models explicitly represent underlying processes and can extrapolate beyond observed conditions, making them valuable tools for forecasting and hindcasting (Malmborg et al. 2024). However, these models require joint measurements of microclimate and vegetation for

validation, which remain rare. Nevertheless, the microclimate community has made impressive progress in producing high-resolution gridded products of microclimate temperature at local (Gril, Laslier, et al. 2023; Niittynen et al. 2024), regional (Brüna et al. 2026; Greiser et al. 2018), continental (Haesen, Lembrechts, et al. 2023), and global scales (Lembrechts et al. 2022). While these datasets offer broad spatial coverage, many represent static microclimate landscapes despite substantial temporal variability and uncertainty. Researchers should therefore critically evaluate their temporal representativeness and develop gridded products that better integrate spatial and temporal dynamics (Klinges et al. 2024).

Many of the physical principles governing microclimates have been established through decades of work in micrometeorology, boundary-layer climatology, and land-atmosphere science (Bonan 2016, 2019; Oke 1987). However, these advances remain only partly integrated into ecological research (Klinges et al. 2026), while the representation of vegetation, and the complexity of the biosphere, in many atmospheric and land-surface models is still largely based on a few broad plant functional types that inadequately capture species composition, vegetation dynamics, disturbance, and fine-scale variation in vegetation structure. Progress will require closer integration among ecology, plant ecophysiology, and atmospheric sciences. Ecologists increasingly need to move beyond treating microclimates as static offsets from macroclimate and engage with turbulent exchange, canopy coupling, and land-atmosphere interactions, while also seeking collaborations with atmospheric scientists when developing or applying microclimate models and interpreting field observations. Atmospheric scientists, in turn, need to better represent vegetation dynamics and biologically relevant microclimatic heterogeneity (Kempainen et al. 2024). Plant ecophysiology provides a key bridge because vegetation both modifies and responds to microclimate through transpiration, phenology, growth, and stress responses. While many of the underlying physical principles are well established, integrating them with ecological and ecophysiological understanding remains a major interdisciplinary challenge. We intend this review to serve as a bridge between ecology and atmospheric science by providing an accessible synthesis of the physical processes governing microclimate change while highlighting the need for closer collaboration between these disciplines. Much of the foundational knowledge needed for this integration has existed for decades (Geiger 1966)—what is now needed is the interdisciplinary will to apply it.

Affiliations

¹Department of Physical Geography and Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden | ²Department of Forest Ecology and Management, Swedish University of Agricultural Sciences, Umeå, Sweden | ³CNRS, Chrono-Environnement (UMR 6249), Université Marie et Louis Pasteur, Besançon, France | ⁴Environment and Sustainability Institute, University of Exeter, Penryn Campus, Penryn, UK | ⁵Centre for Ecology and Evolution, University of Exeter, Penryn Campus, Penryn, UK | ⁶German Center for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany | ⁷Institute of Biology, Leipzig University, Leipzig, Germany | ⁸CNRS, EPHE, IRD, CEF, Univ Montpellier, Montpellier Cedex 5, France | ⁹Ecology and

Biodiversity, Department of Biology, Utrecht University, Utrecht, the Netherlands | ¹⁰Department of Plant and Soil Sciences, University of Pretoria, Pretoria, South Africa | ¹¹Botany and Mycology Unit, Finnish Museum of Natural History, University of Helsinki, Helsinki, Finland | ¹²Department of Organismal and Evolutionary Biology, Faculty of Biological and Environmental Sciences, University of Helsinki, Helsinki, Finland | ¹³Climate Impacts and Adaptation, Finnish Meteorological Institute, Helsinki, Finland | ¹⁴Department of Geosciences and Geography, University of Helsinki, Helsinki, Finland | ¹⁵Department of Plant and Environmental Sciences, University of Copenhagen, Frederiksberg C, Denmark | ¹⁶Globe Institute, University of Copenhagen, Copenhagen K, Denmark | ¹⁷Forest & Nature Lab, Gontrode, Belgium | ¹⁸Department of Earth and Environmental Sciences, KU Leuven, Leuven, Belgium | ¹⁹KU Leuven Plant Institute, KU Leuven, Leuven, Belgium | ²⁰School of the Environment, Yale University, New Haven, Connecticut, USA | ²¹Department of Ecology, Evolution, & Natural Resources, Rutgers University, New Brunswick, New Jersey, USA | ²²Institute of Botany of the Czech Academy of Sciences, Průhonice, Czech Republic | ²³Department of Forest Ecology, Faculty of Forestry and Wood Sciences, Czech University of Life Sciences Prague, Praha-Suchbát, Czech Republic | ²⁴UMR CNRS 7058 “Ecologie et Dynamique des Systèmes Anthropisés” (EDYSAN), Université de Picardie Jules Verne, Amiens, France | ²⁵Faculty of Environment, Science and Economy, University of Exeter, Exeter, UK | ²⁶UK Centre for Ecology & Hydrology, Wallingford, UK | ²⁷Rosby Centre, Swedish Meteorological and Hydrological Institute (SMHI), Norrköping, Sweden | ²⁸Department of Thematic Studies—Environmental Change, Linköping University, Linköping, Sweden | ²⁹Department of Ecosystem Management, Climate and Biodiversity, Institute of Botany, BOKU University, Vienna, Austria | ³⁰Plants and Ecosystems (PLECO), Department of Biology, University of Antwerp, Wilrijk, Belgium

Author Contributions

C. Greiser: conceptualization, supervision, visualization, writing – original draft, writing – review and editing. **E. Gril:** conceptualization, supervision, visualization, writing – original draft, writing – review and editing. **I. M. D. Maclean:** conceptualization, data curation, formal analysis, supervision, visualization, writing – original draft, writing – review and editing. **R. Beugnon:** conceptualization, visualization, writing – original draft, writing – review and editing. **P. le Roux:** conceptualization, writing – original draft, writing – review and editing. **J. Kemppinen:** conceptualization, writing – original draft, writing – review and editing. **M. W. Rhodes:** conceptualization, data curation, formal analysis, visualization, writing – original draft, writing – review and editing. **J. Aalto:** conceptualization, writing – original draft, writing – review and editing. **D. M. Christiansen:** conceptualization, writing – original draft, writing – review and editing. **P. De Frenne:** conceptualization, writing – original draft, writing – review and editing. **S. Haesen:** conceptualization, writing – original draft, writing – review and editing. **D. Klínges:** conceptualization, writing – original draft, writing – review and editing. **M. Kopecký:** conceptualization, writing – original draft, writing – review and editing. **J. Lenoir:** conceptualization, writing – original draft, writing – review and editing. **M. Luoto:** conceptualization, writing – original draft, writing – review and editing. **M. Rantanen:** conceptualization, writing – original draft, writing – review and editing. **G. Strandberg:** conceptualization, writing – original draft, writing – review and editing. **L. van den Brink:** conceptualization, writing – original draft, writing – review and editing. **L. M. Mercado:** conceptualization, writing – original draft, writing – review and editing. **J. Lembrechts:** conceptualization, funding acquisition, supervision, writing – original draft, writing – review and editing. **K. Van Meerbeek:** conceptualization, funding acquisition, supervision, visualization, writing – original draft, writing – review and editing.

Acknowledgments

We all thank the SoilTemp Scientific Research Network (W004024) of the Research Foundation Flanders (FWO). C.G. acknowledges

funding from the Swedish Research Council FORMAS (project grant 2021-01993). J.L. acknowledges funding from the Agence Nationale de la Recherche (ANR), under the framework of the collaborative research program funding scheme (PRC Grant no. ANR-21-CE32-0012-03: MaCCMic project) and the France 2030 program (PEPR FORESTT Grant no. ANR-24-PEFO-0003: MONITOR project). J.L. also acknowledges the Région Hauts-de-France, the Ministère de l'Enseignement Supérieur et de la Recherche and the European Fund for Regional Economic Development for their financial support (CPER ECRIN). I.M. acknowledges funding from the UK Natural Environment Research Council (NERC) projects NE/W006618/1 and NE/V000772/1. P.D.F. received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (ERC Consolidator Grant CanopyChange 101124948). J.K. acknowledges funding from the Research Council of Finland (grant no. 353218; 370245). R.B. was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation)—FZT 118, 202548816; Ei862/29-1. M.K. was supported by the Czech Science Foundation (project GACR 23-06614S) and the Czech Academy of Sciences (project RVO 67985939). L.M.M. acknowledges funding from the UK Natural Environment Research Council (NERC) projects NE/X001172/1 and NE/W004895/1. M.R. was supported by the Research Council of Finland (decision no 342890). D.M.C. acknowledges funding from Novo Nordisk Foundation (NNF22OC0078703).

Funding

This work was supported by Novo Nordisk Fonden (NNF22OC0078703), Deutsche Forschungsgemeinschaft (202548816 and Ei862/29-1), Agence Nationale de la Recherche (ANR-21-CE32-0012-03, ANR-24-PEFO-0003), Fonds Wetenschappelijk Onderzoek (W004024), Natural Environment Research Council (NE/V000772/1, NE/W004895/1, NE/W006618/1, NE/X001172/1), Research Council of Finland (342890, 353218, 370245), Grantová Agentura České Republiky (GACR 23-06614S), European Fund for Regional Economic Development (CPER ECRIN), Région Bourgogne Franche-Comté (ATTRACT), France 2030 (A2ForBois), Région Hauts-de-France (CPER ECRIN), Horizon Europe European Research Council (101124948), Akademie Věd České Republiky (RVO 67985939), Svenska Forskningsrådet Formas (2021-01993), and Ministère de l'Enseignement supérieur et de la Recherche (CPER ECRIN).

Conflicts of Interest

The authors declare no conflicts of interest.

Data Availability Statement

The environmental data used in the illustrative modeling scenarios are publicly available from the sources described in the [Supporting Information](#). Simulations were run using the R packages *microclimf* v2.0.1 (<https://doi.org/10.5281/zenodo.18274614>) and *micropoint* (<https://github.com/ilyamaclean/micropoint>), using Git commit c9d95d852fb576313165792e0ffda5c70e546361).

References

- Aalto, J., D. Scherrer, J. Lenoir, A. Guisan, and M. Luoto. 2018. “Biogeophysical Controls on Soil-Atmosphere Thermal Differences: Implications on Warming Arctic Ecosystems.” *Environmental Research Letters* 13, no. 7: 074003. <https://doi.org/10.1088/1748-9326/aac83e>.
- Aartsma, P., J. Asplund, A. Odland, S. Reinhardt, and H. Renssen. 2020. “Surface Albedo of Alpine Lichen Heaths and Shrub Vegetation.” *Arctic, Antarctic, and Alpine Research* 52, no. 1: 312–322. <https://doi.org/10.1080/15230430.2020.1778890>.
- Al-Ruzouq, R., A. Shanableh, M. A. Khalil, et al. 2022. “Spatial and Temporal Inversion of Land Surface Temperature Along Coastal Cities

- in Arid Regions." *Remote Sensing* 14, no. 8: 1893. <https://doi.org/10.3390/rs14081893>.
- Arora, V. K., A. Katavouta, R. G. Williams, et al. 2020. "Carbon-Concentration and Carbon-Climate Feedbacks in CMIP6 Models and Their Comparison to CMIP5 Models." *Biogeosciences* 17, no. 16: 4173–4222. <https://doi.org/10.5194/bg-17-4173-2020>.
- Ashcroft, M. B., L. A. Chisholm, and K. O. French. 2009. "Climate Change at the Landscape Scale: Predicting Fine-Grained Spatial Heterogeneity in Warming and Potential Refugia for Vegetation." *Global Change Biology* 15, no. 3: 656–667. <https://doi.org/10.1111/j.1365-2486.2008.01762.x>.
- Ashcroft, M. B., and J. R. Gollan. 2013. "Moisture, Thermal Inertia, and the Spatial Distributions of Near-Surface Soil and Air Temperatures: Understanding Factors That Promote Microrefugia." *Agricultural and Forest Meteorology* 176: 77–89. <https://doi.org/10.1016/j.agrformet.2013.03.008>.
- Badeck, F.-W., A. Bondeau, K. Böttcher, et al. 2004. "Responses of Spring Phenology to Climate Change." *New Phytologist* 162, no. 2: 295–309. <https://doi.org/10.1111/j.1469-8137.2004.01059.x>.
- Baguskas, S. A., C. J. Still, D. T. Fischer, C. M. D'Antonio, and J. Y. King. 2016. "Coastal Fog During Summer Drought Improves the Water Status of Sapling Trees More Than Adult Trees in a California Pine Forest." *Oecologia* 181, no. 1: 137–148. <https://doi.org/10.1007/s00442-016-3556-y>.
- Baig, S., B. E. Medlyn, L. M. Mercado, and S. Zaehle. 2015. "Does the Growth Response of Woody Plants to Elevated CO₂ Increase With Temperature? A Model-Oriented Meta-Analysis." *Global Change Biology* 21, no. 12: 4303–4319. <https://doi.org/10.1111/gcb.12962>.
- Baker, T. P., M. T. Moroni, M. A. Hunt, D. Worledge, and D. S. Mendham. 2021. "Temporal, Environmental and Spatial Changes in the Effect of Windbreaks on Pasture Microclimate." *Agricultural and Forest Meteorology* 297: 108265. <https://doi.org/10.1016/j.agrformet.2020.108265>.
- Baldocchi, D. D., and C. A. Vogel. 1996. "Energy and CO₂ Flux Densities Above and Below a Temperate Broad-Leaved Forest and a Boreal Pine Forest." *Tree Physiology* 16, no. 1–2: 5–16. <https://doi.org/10.1093/treephys/16.1-2.5>.
- Barry, R. G. 2008. *Mountain Weather and Climate*. 3rd ed. Cambridge University Press. <https://doi.org/10.1017/CBO9780511754753>.
- Barry, R. G., and P. D. Blanken. 2016. *Microclimate and Local Climate*. Cambridge University Press. <https://doi.org/10.1017/cbo9781316535981>.
- Barry, R. G., and R. G. Chorley. 2009. *Atmosphere, Weather and Climate*. 9th ed. Routledge. <https://doi.org/10.4324/9780203871027>.
- Bertolino, L. T., R. S. Caine, and J. E. Gray. 2019. "Impact of Stomatal Density and Morphology on Water-Use Efficiency in a Changing World." *Frontiers in Plant Science* 10: 427588. <https://doi.org/10.3389/fpls.2019.00225>.
- Beugnon, R., N. Le Guyader, A. Milcu, et al. 2024. "Microclimate Modulation: An Overlooked Mechanism Influencing the Impact of Plant Diversity on Ecosystem Functioning." *Global Change Biology* 30, no. 3: e17214. <https://doi.org/10.1111/gcb.17214>.
- Bonan, G. 2016. *Ecological Climatology*. 3rd ed. Cambridge University Press.
- Bonan, G. 2019. *Climate Change and Terrestrial Ecosystem Modeling*. Cambridge University Press.
- Bright, R. M., C. Antón-Fernández, R. Astrup, F. Cherubini, M. Kvælevåg, and A. H. Strømman. 2014. "Climate Change Implications of Shifting Forest Management Strategy in a Boreal Forest Ecosystem of Norway." *Global Change Biology* 20, no. 2: 607–621. <https://doi.org/10.1111/gcb.12451>.
- Brogli, R., S. Lund Sørland, N. Kröner, and C. Schär. 2021. "Future Summer Warming Pattern Under Climate Change Is Affected by Lapse-Rate Changes." *Weather and Climate Dynamics* 2, no. 4: 1093–1110. <https://doi.org/10.5194/wcd-2-1093-2021>.
- Brůna, J., M. Macek, M. Man, et al. 2026. "High-Resolution Microclimatic Grids for the Bohemian Forest Ecosystem Based on in Situ Measurements." *Scientific Data* 13, no. 1: 246. <https://doi.org/10.1038/s41597-026-06566-z>.
- Buckley, L. B., B. A. Briones Ortiz, I. Caruso, et al. 2023. "TrenchR: An R Package for Modular and Accessible Microclimate and Biophysical Ecology." *PLOS Climate* 2, no. 8: e0000139. <https://doi.org/10.1371/journal.pclm.0000139>.
- Byrne, M. P., and P. A. O'Gorman. 2018. "Trends in Continental Temperature and Humidity Directly Linked to Ocean Warming." *Proceedings of the National Academy of Sciences of the United States of America* 115, no. 19: 4863–4868. <https://doi.org/10.1073/pnas.1722312115>.
- Cai, Y., K. Sawada, and K. Makoto. 2025. "How Microclimate Influences the Spring Phenological Responses to Decreased Snow Cover in Four Tree Species Seedlings in a Boreal Forest." *Scientific Reports* 15, no. 1: 10742. <https://doi.org/10.1038/s41598-025-94523-y>.
- Campbell, G. S., and J. Norman. 2012. *An Introduction to Environmental Biophysics*. Springer Science & Business Media. <https://doi.org/10.1007/978-1-4612-1626-1>.
- Chan, W.-P., J. Lenoir, G.-S. Mai, H.-C. Kuo, I.-C. Chen, and S.-F. Shen. 2024. "Climate Velocities and Species Tracking in Global Mountain Regions." *Nature* 629, no. 8010: 114–120. <https://doi.org/10.1038/s41586-024-07264-9>.
- Chen, J., S. C. Saunders, T. R. Crow, et al. 1999. "Microclimate in Forest Ecosystem and Landscape Ecology: Variations in Local Climate Can Be Used to Monitor and Compare the Effects of Different Management Regimes." *Bioscience* 49, no. 4: 288–297. <https://doi.org/10.2307/1313612>.
- Chesnoiu, G., I. Chiapello, N. Ferlay, P. Nabat, M. Mallet, and V. Riffault. 2025. "Regional Modeling of Surface Solar Radiation, Aerosol, and Cloud Cover Spatial Variability and Projections Over Northern France and Benelux." *Atmospheric Chemistry and Physics* 25, no. 2: 1307–1331. <https://doi.org/10.5194/acp-25-1307-2025>.
- Choler, P., A. Bayle, B. Z. Carlson, C. Randin, G. Filippa, and E. Cremonese. 2021. "The Tempo of Greening in the European Alps: Spatial Variations on a Common Theme." *Global Change Biology* 27, no. 21: 5614–5628. <https://doi.org/10.1111/gcb.15820>.
- Choudhury, B., S. Idso, and R. Reginato. 1987. "Analysis of an Empirical Model for Soil Heat Flux Under a Growing Wheat Crop for Estimating Evaporation by an Infrared-Temperature Based Energy Balance Equation." *Agricultural and Forest Meteorology* 39, no. 4: 283–297. [https://doi.org/10.1016/0168-1923\(87\)90021-9](https://doi.org/10.1016/0168-1923(87)90021-9).
- Christiansen, D. M., G. Römer, J. P. Dahlgren, et al. 2024. "High-Resolution Data Are Necessary to Understand the Effects of Climate on Plant Population Dynamics of a Forest Herb." *Ecology* 105, no. 1: e4191. <https://doi.org/10.1002/ecy.4191>.
- Clapeyron, E. 1834. "Mémoire Sur la Puissance Motrice de la Chaleur." *Journal de l'Ecole Polytechnique*: 153–190. <http://catalogue.bnf.fr/ark:/12148/cb40936143n>.
- Clausius, R. 1850. "Über die Bewegende Kraft der Wärme und die Gesetze, Welche Sich Daraus Für Die Wärmelehre Selbst Ableiten Lassen." *Annalen der Physik* 155, no. 3: 368–397. <https://doi.org/10.1002/andp.18501550306>.
- Cowley, G. S., J. D. Niemann, T. R. Green, M. S. Seyfried, A. S. Jones, and P. J. Grazaitis. 2017. "Impacts of Precipitation and Potential Evapotranspiration Patterns on Downscaling Soil Moisture in Regions With Large Topographic Relief." *Water Resources Research* 53, no. 2: 1553–1574. <https://doi.org/10.1002/2016WR019907>.
- Cox, D. T. C., I. M. D. Maclean, A. S. Gardner, and K. J. Gaston. 2020. "Global Variation in Diurnal Asymmetry in Temperature, Cloud Cover,

- Specific Humidity and Precipitation and Its Association With Leaf Area Index." *Global Change Biology* 26, no. 12: 7099–7111. <https://doi.org/10.1111/gcb.15336>.
- Dada, L., D. Stolzenburg, M. Simon, et al. 2023. "Role of Sesquiterpenes in Biogenic New Particle Formation." *Science Advances* 9, no. 36: eadi5297. <https://doi.org/10.1126/sciadv.adi5297>.
- Dai, A., K. E. Trenberth, and T. R. Karl. 1999. "Effects of Clouds, Soil Moisture, Precipitation, and Water Vapor on Diurnal Temperature Range." *Journal of Climate* 12, no. 8: 2451–2473. [https://doi.org/10.1175/1520-0442\(1999\)012<2451:EOCSMP>2.0.CO;2](https://doi.org/10.1175/1520-0442(1999)012<2451:EOCSMP>2.0.CO;2).
- Dai, J., H. Wang, and Q. Ge. 2014. "The Spatial Pattern of Leaf Phenology and Its Response to Climate Change in China." *International Journal of Biometeorology* 58, no. 4: 521–528. <https://doi.org/10.1007/s00484-013-0679-2>.
- Daly, C., D. R. Conklin, and M. H. Unsworth. 2010. "Local Atmospheric Decoupling in Complex Topography Alters Climate Change Impacts." *International Journal of Climatology* 30, no. 12: 1857–1864. <https://doi.org/10.1002/joc.2007>.
- Davis, K. T., S. Z. Dobrowski, Z. A. Holden, P. E. Higuera, and J. T. Abatzoglou. 2019. "Microclimatic Buffering in Forests of the Future: The Role of Local Water Balance." *Ecography* 42, no. 1: 1–11. <https://doi.org/10.1111/ecog.03836>.
- De Frenne, P., R. Beugnon, D. Klimes, et al. 2025. "Ten Practical Guidelines for Microclimate Research in Terrestrial Ecosystems." *Methods in Ecology and Evolution* 16, no. 2: 269–294. <https://doi.org/10.1111/2041-210x.14476>.
- De Frenne, P., J. Lenoir, M. Luoto, et al. 2021. "Forest Microclimates and Climate Change: Importance, Drivers and Future Research Agenda." *Global Change Biology* 27, no. 11: 2279–2297. <https://doi.org/10.1111/gcb.15569>.
- Debouk, H., F. de Bello, and M.-T. Sebastià. 2015. "Functional Trait Changes, Productivity Shifts and Vegetation Stability in Mountain Grasslands During a Short-Term Warming." *PLoS One* 10, no. 10: e0141899. <https://doi.org/10.1371/journal.pone.0141899>.
- Devine, A. P., R. A. McDonald, T. Quaife, and I. M. D. Maclean. 2017. "Determinants of Woody Encroachment and Cover in African Savannas." *Oecologia* 183, no. 4: 939–951. <https://doi.org/10.1007/s00442-017-3807-6>.
- Dobrowski, S. Z. 2011. "A Climatic Basis for Microrefugia: The Influence of Terrain on Climate." *Global Change Biology* 17, no. 2: 1022–1035. <https://doi.org/10.1111/j.1365-2486.2010.02263.x>.
- D'Odorico, P., Y. He, S. Collins, S. F. J. De Wekker, V. Engel, and J. D. Fuentes. 2013. "Vegetation-Microclimate Feedbacks in Woodland-Grassland Ecotones." *Global Ecology and Biogeography: A Journal of Macroecology* 22, no. 4: 364–379. <https://doi.org/10.1111/geb.12000>.
- Dong, B., R. T. Sutton, and L. J. Wilcox. 2022. "Decadal Trends in Surface Solar Radiation and Cloud Cover Over the North Atlantic Sector During the Last Four Decades: Drivers and Physical Processes." *Climate Dynamics* 60, no. 7: 2533–2546. <https://doi.org/10.1007/s00382-022-06438-3>.
- Dong, X., H. Lin, F. Wang, S. Shi, J. Ma, and X. He. 2025. "Effects of Increasing CO₂ Concentration on Crop Growth and Soil Ammonia-Oxidizing Microorganisms in a Fababean (*Vicia faba* L.) and Wheat (*Triticum aestivum* Yunmai) Intercropping System." *Plants* 14, no. 4: 516. <https://doi.org/10.3390/plants14040516>.
- Dou, Y., J. Quan, X. Jia, Q. Wang, and Y. Liu. 2021. "Near-Surface Warming Reduces Dew Frequency in China." *Geophysical Research Letters* 48, no. 7. <https://doi.org/10.1029/2020gl091923>.
- Doughty, C. E., J. M. Keany, B. C. Wiebe, et al. 2023. "Tropical Forests Are Approaching Critical Temperature Thresholds." *Nature* 621, no. 7977: 105–111. <https://doi.org/10.1038/s41586-023-06391-z>.
- Dror, T., and G. Feingold. 2026. "Amazon Forest Loss: An All-Sky Biophysical Top-of-Atmosphere Cooling Feedback." *Science* 392, no. 6796: 429–432. <https://doi.org/10.1126/science.adz8296>.
- Duveiller, G., F. Filippini, A. Ceglar, J. Bojanowski, R. Alkama, and A. Cescatti. 2021. "Revealing the Widespread Potential of Forests to Increase Low Level Cloud Cover." *Nature Communications* 12, no. 1: 4337. <https://doi.org/10.1038/s41467-021-24551-5>.
- Ellsworth, D. S., I. C. Anderson, K. Y. Crous, et al. 2017. "Elevated CO₂ Does Not Increase Eucalypt Forest Productivity on a Low-Phosphorus Soil." *Nature Climate Change* 7, no. 4: 279–282. <https://doi.org/10.1038/nclimate3235>.
- Evans, J. P., X. Meng, and M. F. McCabe. 2017. "Land Surface Albedo and Vegetation Feedbacks Enhanced the Millennium Drought in South-East Australia." *Hydrology and Earth System Sciences* 21, no. 1: 409–422. <https://doi.org/10.5194/hess-21-409-2017>.
- Fernández, V., H. A. Bahamonde, J. Javier Peguero-Pina, et al. 2017. "Physico-Chemical Properties of Plant Cuticles and Their Functional and Ecological Significance." *Journal of Experimental Botany* 68, no. 19: 5293–5306. <https://doi.org/10.1093/jxb/erx302>.
- Finne, E. A., J. W. Bjerke, R. Erlandsson, H. Tømmervik, F. Stordal, and L. M. Tallaksen. 2023. "Variation in Albedo and Other Vegetation Characteristics in Non-Forested Northern Ecosystems: The Role of Lichens and Mosses." *Environmental Research Letters* 18, no. 7: 074038. <https://doi.org/10.1088/1748-9326/ace06d>.
- Finnigan, J. 2000. "Turbulence in Plant Canopies." *Annual Review of Fluid Mechanics* 32, no. 1: 519–571. <https://doi.org/10.1146/annurev.fluid.32.1.519>.
- Forkel, M., N. Carvalhais, C. Rödenbeck, et al. 2016. "Enhanced Seasonal CO₂ Exchange Caused by Amplified Plant Productivity in Northern Ecosystems." *Science* 351, no. 6274: 696–699. <https://doi.org/10.1126/science.aac4971>.
- García, R. A., and S. Clusella-Trullas. 2019. "Thermal Landscape Change as a Driver of Ectotherm Responses to Plant Invasions." *Proceedings of the Royal Society B: Biological Sciences* 286: 20191020. <https://doi.org/10.1098/rspb.2019.1020>.
- García-García, A., F. J. Cuesta-Valero, D. G. Miralles, et al. 2023. "Soil Heat Extremes Can Outpace Air Temperature Extremes." *Nature Climate Change* 13, no. 11: 1237–1241. <https://doi.org/10.1038/s41558-023-01812-3>.
- Geiger, R. 1966. *The Climate Near the Ground*. Harvard University Press. https://books.google.fr/books/about/Climate_Near_the_Ground.html?id=e0zYzwEACAAJ.
- Greenberg, J. P., A. B. Guenther, G. Pétron, et al. 2004. "Biogenic VOC Emissions From Forested Amazonian Landscapes." *Global Change Biology* 10, no. 5: 651–662. <https://doi.org/10.1111/j.1365-2486.2004.00758.x>.
- Greiser, C., L. Huo, M. Ghaly, et al. 2025. "Bark Beetles as Microclimate Engineers—Thermal Characteristics of Infested Spruce Trees at the Canopy Surface and Below the Canopy." *Agricultural and Forest Meteorology* 375: 110796. <https://doi.org/10.1016/j.agrformet.2025.110796>.
- Greiser, C., E. Meineri, M. Luoto, J. Ehrlén, and K. Hylander. 2018. "Monthly Microclimate Models in a Managed Boreal Forest Landscape." *Agricultural and Forest Meteorology* 250–251: 147–158. <https://doi.org/10.1016/j.agrformet.2017.12.252>.
- Gril, E., M. Laslier, E. Gallet-Moron, et al. 2023. "Using Airborne LiDAR to Map Forest Microclimate Temperature Buffering or Amplification." *Remote Sensing of Environment* 298: 113820. <https://doi.org/10.1016/j.rse.2023.113820>.
- Gril, E., F. Spicher, C. Greiser, et al. 2023. "Slope and Equilibrium: A Parsimonious and Flexible Approach to Model Microclimate." *Methods in Ecology and Evolution* 14, no. 3: 885–897. <https://doi.org/10.1111/2041-210x.14048>.

- Grimm, N. B., F. S. Chapin III, B. Bierwagen, et al. 2013. "The Impacts of Climate Change on Ecosystem Structure and Function." *Frontiers in Ecology and the Environment* 11, no. 9: 474–482. <https://doi.org/10.1890/120282>.
- Grimmond, C. S. B., S. M. Robeson, and J. T. Schoof. 2000. "Spatial Variability of micro-Climatic Conditions Within a Mid-Latitude Deciduous Forest." *Climate Research* 15: 137–149. <https://doi.org/10.3354/cr015137>.
- Groot, A., and D. W. Carlson. 1996. "Influence of Shelter on Night Temperatures, Frost Damage, and Bud Break of White Spruce Seedlings." *Canadian Journal of Forest Research* 26, no. 9: 1531–1538. <https://doi.org/10.1139/x26-172>.
- Grossiord, C., T. N. Buckley, L. A. Cernusak, et al. 2020. "Plant Responses to Rising Vapor Pressure Deficit." *New Phytologist* 226, no. 6: 1550–1566. <https://doi.org/10.1111/nph.16485>.
- Guthrie, D. R. 2001. "Origin and Causes of the Mammoth Steppe: A Story of Cloud Cover, Woolly Mammal Tooth Pits, Buckles, and Inside-Out Beringia." *Quaternary Science Reviews* 20, no. 1–3: 549–574. [https://doi.org/10.1016/s0277-3791\(00\)00099-8](https://doi.org/10.1016/s0277-3791(00)00099-8).
- Haesen, S., J. J. Lembrechts, P. De Frenne, et al. 2023. "ForestClim-Bioclimatic Variables for Microclimate Temperatures of European Forests." *Global Change Biology* 29, no. 11: 2886–2892. <https://doi.org/10.1111/gcb.16678>.
- Haesen, S., J. Lenoir, E. Gril, et al. 2023. "Microclimate Reveals the True Thermal Niche of Forest Plant Species." *Ecology Letters* 26, no. 12: 2043–2055. <https://doi.org/10.1111/ele.14312>.
- Haesen, S., C. Senf, R. Seidl, et al. 2026. "Undisturbed Forest Canopies Slow Understory Warming by One-Third Across Europe." *Research Square*. <https://doi.org/10.21203/rs.3.rs-8811214/v1>.
- Hagan Brown, W., E. Gloor, R. Fyfe, et al. 2025. "Elevated CO₂ Increases the Canopy Temperature of Mature *Quercus robur* (Pedunculate Oak)." *Global Change Biology* 31, no. 11: e70565. <https://doi.org/10.1111/gcb.70565>.
- Helmuth, B., B. R. Broitman, L. Yamane, et al. 2010. "Organismal Climatology: Analyzing Environmental Variability at Scales Relevant to Physiological Stress." *Journal of Experimental Biology* 213, no. 6: 995–1003. <https://doi.org/10.1242/jeb.038463>.
- Hiebl, J., and W. Schöner. 2018. "Temperature Inversions in Austria in a Warming Climate—Changes in Space and Time." *Meteorologische Zeitschrift* 27, no. 4: 309–323.
- Hou, X., M. Wild, D. S. Folini, S. Kazadzis, and J. Wohland. 2021. "Climate Change Impacts on Solar Power Generation and Its Spatial Variability in Europe Based on CMIP6." *Earth System Dynamics* 12, no. 4: 1099–1113. <https://doi.org/10.5194/esd-2021-57>.
- Huang, Y., G. Stein, O. Kolle, et al. 2024. "Enhanced Stability of Grassland Soil Temperature by Plant Diversity." *Nature Geoscience* 17, no. 1: 44–50. <https://doi.org/10.1038/s41561-023-01338-5>.
- Inouye, D. W. 2022. "Climate Change and Phenology." *Wiley Interdisciplinary Reviews: Climate Change* 13, no. 3. <https://doi.org/10.1002/wcc.764>.
- IPBES. 2019. *Summary for Policymakers of the Global Assessment Report on Biodiversity and Ecosystem Services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services*, edited by S. Diaz, J. Settele, E. S. Brondizio, et al. IPBES secretariat, Bonn.
- IPCC. 2022. *Climate Change and Land: IPCC Special Report on Climate Change, Desertification, Land Degradation, Sustainable Land Management, Food Security, and Greenhouse Gas Fluxes in Terrestrial Ecosystems*. Cambridge University Press. <https://doi.org/10.1017/9781009157988>.
- IPCC. 2023a. *Climate Change 2021—The Physical Science Basis: Working Group I Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press. <https://doi.org/10.1017/9781009157896>.
- IPCC. 2023b. *Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Core Writing Team, H. Lee, and J. Romero. <https://doi.org/10.59327/IPCC/AR6-9789291691647.001>.
- Jiao, T., C. A. Williams, B. Ghimire, J. Masek, F. Gao, and C. Schaaf. 2017. "Global Climate Forcing From Albedo Change Caused by Large-Scale Deforestation and Reforestation: Quantification and Attribution of Geographic Variation." *Climatic Change* 142, no. 3: 463–476. <https://doi.org/10.1007/s10584-017-1962-8>.
- Johnson, D. M., and W. K. Smith. 2008. "Cloud Immersion Alters Microclimate, Photosynthesis and Water Relations in Rhododendron Catawbiense and *Abies fraseri* Seedlings in the Southern Appalachian Mountains, USA." *Tree Physiology* 28, no. 3: 385–392. <https://doi.org/10.1093/treephys/28.3.385>.
- Jones, H. G. 2014. *Plants and Microclimate*. 3rd ed. Cambridge University Press. <https://doi.org/10.1017/cbo9780511845727>.
- Jones, J. A., C. Daly, M. Schulze, and C. J. Still. 2025. "Microclimate Refugia Are Transient in Stable Old Forests, Pacific Northwest, USA." *AGU Advances* 6, no. 3. <https://doi.org/10.1029/2024av001492>.
- Jump, A. S., P. Ruiz-Benito, S. Greenwood, et al. 2017. "Structural Overshoot of Tree Growth With Climate Variability and the Global Spectrum of Drought-Induced Forest Dieback." *Global Change Biology* 23, no. 9: 3742–3757. <https://doi.org/10.1111/gcb.13636>.
- Kearney, M. R., and A. Leigh. 2024. "Fast, Accurate and Accessible Calculations of Leaf Temperature and Its Physiological Consequences With NicheMapR." *Methods in Ecology and Evolution* 15, no. 9: 1516–1531. <https://doi.org/10.1111/2041-210x.14373>.
- Kearney, M. R., and W. P. Porter. 2017. "NicheMapR—An R Package for Biophysical Modelling: The Microclimate Model." *Ecography* 40, no. 5: 664–674. <https://doi.org/10.1111/ecog.02360>.
- Kemppinen, J., J. J. Lembrechts, K. Van Meerbeek, et al. 2024. "Microclimate, an Important Part of Ecology and Biogeography." *Global Ecology and Biogeography: A Journal of Macroecology* 33, no. 6. <https://doi.org/10.1111/gcb.13834>.
- Kemppinen, J., P. Niittynen, P. C. le Roux, et al. 2021. "Consistent Trait-Environment Relationships Within and Across Tundra Plant Communities." *Nature Ecology & Evolution* 5, no. 4: 458–467. <https://doi.org/10.1038/s41559-021-01396-1>.
- Kemppinen, J., P. Niittynen, A.-M. Virkkala, et al. 2021. "Dwarf Shrubs Impact Tundra Soils: Drier, Colder, and Less Organic Carbon." *Ecosystems* 24, no. 6: 1378–1392. <https://doi.org/10.1007/s10021-020-00589-2>.
- Klemm, O., and N. Lin. 2016. "What Causes Observed Fog Trends: Air Quality or Climate Change?" *Aerosol and Air Quality Research* 16, no. 5: 1131–1142. <https://doi.org/10.4209/aaqr.2015.05.0353>.
- Klinges, D. H., J. A. Baecher, J. J. Lembrechts, et al. 2024. "Proximal Microclimate: Moving Beyond Spatiotemporal Resolution Improves Ecological Predictions." *Global Ecology and Biogeography: A Journal of Macroecology* 33, no. 9: e13884. <https://doi.org/10.1111/gcb.13884>.
- Klinges, D. H., M. M. Muñoz, S. F. Domínguez-Guerrero, I. M. D. Maclean, M. R. Kearney, and D. K. Skelly. 2026. "Matching Climate to Biological Scales." *Trends in Ecology & Evolution* 41, no. 4: 329–343. <https://doi.org/10.1016/j.tree.2025.11.015>.
- Kopáček, J., R. Bače, J. Hejzlar, et al. 2020. "Changes in Microclimate and Hydrology in an Unmanaged Mountain Forest Catchment After Insect-Induced Tree Dieback." *Science of the Total Environment* 720: 137518. <https://doi.org/10.1016/j.scitotenv.2020.137518>.

- Korell, L., H. Auge, J. M. Chase, W. S. Harpole, and T. M. Knight. 2021. "Responses of Plant Diversity to Precipitation Change Are Strongest at Local Spatial Scales and in Drylands." *Nature Communications* 12, no. 1: 2489. <https://doi.org/10.1038/s41467-021-22766-0>.
- Körner, C., and E. Hiltbrunner. 2018. "The 90 Ways to Describe Plant Temperature." *Perspectives in Plant Ecology, Evolution and Systematics* 30: 16–21. <https://doi.org/10.1016/j.ppees.2017.04.004>.
- Kramp, R. E., P. Liancourt, M. M. Herberich, et al. 2022. "Functional Traits and Their Plasticity Shift From Tolerant to Avoidant Under Extreme Drought." *Ecology* 103, no. 12: e3826. <https://doi.org/10.1002/ecy.3826>.
- Kretzschmar, J., M. Pöhlker, F. Stratmann, H. Wex, C. Wirth, and J. Quaas. 2024. "From Trees to Rain: Enhancement of Cloud Glaciation and Precipitation by Pollen." *Environmental Research Letters* 19, no. 10: 104052. <https://doi.org/10.1088/1748-9326/ad747a>.
- Kropp, H., M. M. Loranty, S. M. Natali, et al. 2021. "Shallow Soils Are Warmer Under Trees and Tall Shrubs Across Arctic and Boreal Ecosystems." *Environmental Research Letters* 16, no. 1: 015001. <https://doi.org/10.1088/1748-9326/abc994>.
- Kühn, N., C. Tovar, J. Carretero, V. Vandvik, B. J. Enquist, and K. J. Willis. 2021. "Globally Important Plant Functional Traits for Coping With Climate Change." *Frontiers of Biogeography* 13, no. 4: e53774. <https://doi.org/10.21425/f5fbg53774>.
- Lavigne, T., and C. Liu. 2022. "Validity of Global Fog-Day Trends Indicated by the Global Surface Summary of the Day (GSOD) Data Set." *Journal of Geophysical Research-Atmospheres* 127, no. 10. <https://doi.org/10.1029/2021jd035881>.
- Lehtinen, J., J. Aalto, P. Niittynen, and M. Luoto. 2026. "Synoptic Atmospheric Conditions Drive Spatio-Temporal Microclimatic Variability in High-Latitude Landscapes." *Climate Dynamics* 64, no. 4: 201. <https://doi.org/10.1007/s00382-026-08170-8>.
- Lembrechts, J. J., J. Aalto, M. B. Ashcroft, et al. 2020. "SoilTemp: A Global Database of Near-Surface Temperature." *Global Change Biology* 26, no. 11: 6616–6629. <https://doi.org/10.1111/gcb.15123>.
- Lembrechts, J. J., J. van den Hoogen, J. Aalto, et al. 2022. "Global Maps of Soil Temperature." *Global Change Biology* 28, no. 9: 3110–3144. <https://doi.org/10.1111/gcb.16060>.
- Lenton, T. M. 2013. "Environmental Tipping Points." *Annual Review of Environment and Resources* 38, no. 1: 1–29. <https://doi.org/10.1146/annurev-environ-102511-084654>.
- Lewis, G. N., and M. Randall. 1923. *Thermodynamics and the Free Energy of Chemical Substances*. McGraw-Hill. <https://archive.org/details/thermodynamics00lewi/>.
- Li, Y., J. Ge, H. Wu, et al. 2025. "Amplified Local Cooling Effect of Forestation in Warming Europe." *Nature Communications* 16, no. 1: 8412. <https://doi.org/10.1038/s41467-025-63556-2>.
- Litaor, M. I., M. Williams, and T. R. Seastedt. 2008. "Topographic Controls on Snow Distribution, Soil Moisture, and Species Diversity of Herbaceous Alpine Vegetation, Niwot Ridge, Colorado." *Journal of Geophysical Research: Biogeosciences* 113, no. G2. <https://doi.org/10.1029/2007JG000419>.
- Liu, J., Y. Ryu, X. Luo, et al. 2024. "Evidence for Widespread Thermal Acclimation of Canopy Photosynthesis." *Nature Plants* 10, no. 12: 1919–1927. <https://doi.org/10.1038/s41477-024-01846-1>.
- Lookingbill, T. R., and D. L. Urban. 2003. "Spatial Estimation of Air Temperature Differences for Landscape-Scale Studies in Montane Environments." *Agricultural and Forest Meteorology* 114, no. 3–4: 141–151. [https://doi.org/10.1016/s0168-1923\(02\)00196-x](https://doi.org/10.1016/s0168-1923(02)00196-x).
- Lukeš, P., P. Stenberg, and M. Rautiainen. 2013. "Relationship Between Forest Density and Albedo in the Boreal Zone." *Ecological Modelling* 261–262: 74–79. <https://doi.org/10.1016/j.ecolmodel.2013.04.009>.
- Luo, H., J. Quaas, and Y. Han. 2024a. "Decreased Cloud Cover Partially Offsets the Cooling Effects of Surface Albedo Change due to Deforestation." *Nature Communications* 15, no. 1: 7345. <https://doi.org/10.1038/s41467-024-51783-y>.
- Luo, H., J. Quaas, and Y. Han. 2024b. "Diurnally Asymmetric Cloud Cover Trends Amplify Greenhouse Warming." *Science Advances* 10, no. 25: eado5179. <https://doi.org/10.1126/sciadv.ado5179>.
- Maclean, I. M. D. 2020. "Predicting Future Climate at High Spatial and Temporal Resolution." *Global Change Biology* 26, no. 2: 1003–1011. <https://doi.org/10.1111/gcb.14876>.
- Maclean, I. M. D. 2026. "Microclimf: Fast Modelling of Microclimate Across Real Landscapes in R." *Methods in Ecology and Evolution* 17, no. 4: 1112–1123. <https://doi.org/10.1111/2041-210x.70253>.
- Maclean, I. M. D., and R. Early. 2023. "Macroclimate Data Overestimate Range Shifts of Plants in Response to Climate Change." *Nature Climate Change* 13, no. 5: 484–490. <https://doi.org/10.1038/s41558-023-01650-3>.
- Maclean, I. M. D., A. J. Suggitt, R. J. Wilson, J. P. Duffy, and J. J. Bennie. 2017. "Fine-Scale Climate Change: Modelling Spatial Variation in Biologically Meaningful Rates of Warming." *Global Change Biology* 23, no. 1: 256–268. <https://doi.org/10.1111/gcb.13343>.
- Mallen-Cooper, M., B. J. Graae, and W. K. Cornwell. 2021. "Lichens Buffer Tundra Microclimate More Than the Expanding Shrub *Betula nana*." *Annals of Botany* 128, no. 4: 407–418. <https://doi.org/10.1093/aob/mcab041>.
- Malmberg, C. A., A. M. Willson, L. M. Bradley, et al. 2024. "Defining Model Complexity: An Ecological Perspective." *Meteorological Applications* 31, no. 3: e2202. <https://doi.org/10.1002/met.2202>.
- Marchin, R. M., B. E. Medlyn, M. G. Tjoelker, and D. S. Ellsworth. 2023. "Decoupling Between Stomatal Conductance and Photosynthesis Occurs Under Extreme Heat in Broadleaf Tree Species Regardless of Water Access." *Global Change Biology* 29, no. 22: 6319–6335. <https://doi.org/10.1111/gcb.16929>.
- McInnes, K. L., T. A. Erwin, and J. M. Bathols. 2011. "Global Climate Model Projected Changes in 10 m Wind Speed and Direction due to Anthropogenic Climate Change." *Atmospheric Science Letters* 12, no. 4: 325–333. <https://doi.org/10.1002/asl.341>.
- McVicar, T. R., M. L. Roderick, R. J. Donohue, et al. 2012. "Global Review and Synthesis of Trends in Observed Terrestrial Near-Surface Wind Speeds: Implications for Evaporation." *Journal of Hydrology* 416–417: 182–205. <https://doi.org/10.1016/j.jhydrol.2011.10.024>.
- Meeussen, C., S. Govaert, T. Vanneste, et al. 2021. "Microclimatic Edge-to-Interior Gradients of European Deciduous Forests." *Agricultural and Forest Meteorology* 311: 108699. <https://doi.org/10.1016/j.agrformet.2021.108699>.
- Mekonnen, Z. A., W. J. Riley, L. T. Berner, et al. 2021. "Arctic Tundra Shrubification: A Review of Mechanisms and Impacts on Ecosystem Carbon Balance." *Environmental Research Letters* 16, no. 5: 053001. <https://doi.org/10.1088/1748-9326/abf28b>.
- Mengistu, A. G., L. D. van Rensburg, and S. S. W. Mavimbela. 2017. "The Effect of Soil Water and Temperature on Thermal Properties of Two Soils Developed From Aeolian Sands in South Africa." *Catena* 158: 184–193. <https://doi.org/10.1016/j.catena.2017.07.001>.
- Michalet, R., C. Carcaillet, F. Delerue, J.-C. Domec, and J. Lenoir. 2024. "Assisted Migration in a Warmer and Drier Climate: Less Climate Buffering Capacity, Less Facilitation and More Fires at Temperate Latitudes?" *Oikos* 2024, no. 8: e10248. <https://doi.org/10.1111/oik.10248>.
- Minder, J. R., P. W. Mote, and J. D. Lundquist. 2010. "Surface Temperature Lapse Rates Over Complex Terrain: Lessons From the Cascade Mountains." *Journal of Geophysical Research* 115, no. D14. <https://doi.org/10.1029/2009jd013493>.

- Miralles, D. G., J. Vilà-Guerau de Arellano, T. R. McVicar, and M. D. Mahecha. 2025. "Vegetation–Climate Feedbacks Across Scales." *Annals of the New York Academy of Sciences* 1544, no. 1: 27–41. <https://doi.org/10.1111/nyas.15286>.
- Miró, J. R., J. C. Peña, N. Pepin, A. Sairouni, and M. Aran. 2018. "Key Features of Cold-Air Pool Episodes in the Northeast of the Iberian Peninsula (Cerdanya, Eastern Pyrenees)." *International Journal of Climatology* 38, no. 3: 1105–1115. <https://doi.org/10.1002/joc.5236>.
- Mndela, M., J. T. Tjelele, I. C. Madakadze, et al. 2022. "A Global Meta-Analysis of Woody Plant Responses to Elevated CO₂: Implications on Biomass, Growth, Leaf N Content, Photosynthesis and Water Relations." *Ecological Processes* 11, no. 1. <https://doi.org/10.1186/s13717-022-00397-7>.
- Munday, G., C. D. Jones, N. J. Steinert, et al. 2025. "Risks of Unavoidable Impacts on Forests at 1.5°C With and Without Overshoot." *Nature Climate Change* 15, no. 6: 650–655. <https://doi.org/10.1038/s41558-025-02327-9>.
- Muthukrishnan, R., T. M. Smiley, P. O. Title, A. M. Fudickar, A. E. Jahn, and J. A. Lau. 2025. "Chasing the Niche: Escaping Climate Change Threats in Place, Time, and Space." *Global Change Biology* 31, no. 4: e70167. <https://doi.org/10.1111/gcb.70167>.
- Myers-Smith, I. H., and D. S. Hik. 2013. "Shrub Canopies Influence Soil Temperatures but Not Nutrient Dynamics: An Experimental Test of Tundra Snow–Shrub Interactions." *Ecology and Evolution* 3, no. 11: 3683–3700. <https://doi.org/10.1002/ece3.710>.
- Niittynen, P., R. K. Heikkinen, and M. Luoto. 2018. "Snow Cover Is a Neglected Driver of Arctic Biodiversity Loss." *Nature Climate Change* 8, no. 11: 997–1001. <https://doi.org/10.1038/s41558-018-0311-x>.
- Niittynen, P., H. Salminen, P. Peña-Aguilera, et al. 2024. "A Gridded Microclimate Dataset From a Sub-Arctic Biodiversity Hotspot in Finland." *bioRxiv: the preprint server for biology*. <https://doi.org/10.1101/2024.03.30.587419>.
- Norby, R. J. 2025. "Forest Productivity Response to Elevated CO₂ in Free-Air CO₂ Enrichment Experiments: The 23 Percent Solution, Revisited." *New Phytologist* 246, no. 5: 1952–1959. <https://doi.org/10.1111/nph.70162>.
- Oke, T. R. 1987. *Boundary Layer Climates*. 2nd ed. <https://doi.org/10.4324/9780203407219>.
- Pastore, M. A., A. W. D'Amato, J. R. Foster, and E. Carol Adair. 2022. "Cold-Air Pools as Microrefugia for Ecosystem Functions in the Face of Climate Change." *Ecology* 103, no. 8: e3717. <https://doi.org/10.1002/ecy.3717>.
- Pearson, R. G., S. J. Phillips, M. M. Loranty, et al. 2013. "Shifts in Arctic Vegetation and Associated Feedbacks Under Climate Change." *Nature Climate Change* 3, no. 7: 673–677. <https://doi.org/10.1038/nclimate1858>.
- Pecl, G. T., M. B. Araújo, J. D. Bell, et al. 2017. "Biodiversity Redistribution Under Climate Change: Impacts on Ecosystems and Human Well-Being." *Science* 355, no. 6332: eaai9214. <https://doi.org/10.1126/science.aai9214>.
- Peñuelas, J., P. Ciais, J. G. Canadell, et al. 2017. "Shifting From a Fertilization-Dominated to a Warming-Dominated Period." *Nature Ecology & Evolution* 1, no. 10: 1438–1445. <https://doi.org/10.1038/s41558-017-0274-8>.
- Pepin, N., M. Apple, J. Knowles, et al. 2025. "Elevation-Dependent Climate Change in Mountain Environments." *Nature Reviews Earth & Environment* 6, no. 12: 772–788. <https://doi.org/10.1038/s43017-025-00740-4>.
- Piao, S., Q. Liu, A. Chen, et al. 2019. "Plant Phenology and Global Climate Change: Current Progresses and Challenges." *Global Change Biology* 25, no. 6: 1922–1940. <https://doi.org/10.1111/gcb.14619>.
- Piao, S., X. Wang, T. Park, et al. 2019. "Characteristics, Drivers and Feedbacks of Global Greening." *Nature Reviews Earth & Environment* 1, no. 1: 14–27. <https://doi.org/10.1038/s43017-019-0001-x>.
- Pörtner, H.-O., R. J. Scholes, J. Agard, et al. 2021. "Scientific Outcome of the IPBES-IPCC Co-Sponsored Workshop on Biodiversity and Climate Change." *Zenodo*. <https://doi.org/10.5281/ZENODO.5101125>.
- Posch, B. C., S. E. Bush, D. F. Koepke, et al. 2024. "Intensive Leaf Cooling Promotes Tree Survival During a Record Heatwave." *Proceedings of the National Academy of Sciences of the United States of America* 121, no. 43: e2408583121. <https://doi.org/10.1073/pnas.2408583121>.
- Pypker, T. G., M. H. Unsworth, B. Lamb, et al. 2007. "Cold Air Drainage in a Forested Valley: Investigating the Feasibility of Monitoring Ecosystem Metabolism." *Agricultural and Forest Meteorology* 145, no. 3–4: 149–166. <https://doi.org/10.1016/j.agrformet.2007.04.016>.
- Rap, A., C. E. Scott, C. L. Reddington, et al. 2018. "Enhanced Global Primary Production by Biogenic Aerosol via Diffuse Radiation Fertilization." *Nature Geoscience* 11, no. 9: 640–644. <https://doi.org/10.1038/s41561-018-0208-3>.
- Renaud, V., J. L. Innes, M. Dobberty, and M. Rebetez. 2011. "Comparison Between Open-Site and Below-Canopy Climatic Conditions in Switzerland for Different Types of Forests Over 10 Years (1998–2007)." *Theoretical and Applied Climatology* 105, no. 1–2: 119–127. <https://doi.org/10.1007/s00704-010-0361-0>.
- Richter, R., H. Ballasus, R. A. Engelmann, C. Zielhofer, A. Sanaei, and C. Wirth. 2022. "Tree Species Matter for Forest Microclimate Regulation During the Drought Year 2018: Disentangling Environmental Drivers and Biotic Drivers." *Scientific Reports* 12, no. 1: 17559. <https://doi.org/10.1038/s41598-022-22582-6>.
- Riddell, E. A., and C. K. Porter. 2025. "The Wind Niche: The Thermal and Hydric Effects of Wind Speed on Terrestrial Organisms." *Integrative and Comparative Biology* 65, no. 4: 1024–1035. <https://doi.org/10.1093/icb/icafo25>.
- Roland, T. P., O. T. Bartlett, D. J. Charman, et al. 2024. "Sustained Greening of the Antarctic Peninsula Observed From Satellites." *Nature Geoscience* 17, no. 11: 1121–1126. <https://doi.org/10.1038/s41561-024-01564-5>.
- Rolland, C. 2003. "Spatial and Seasonal Variations of Air Temperature Lapse Rates in Alpine Regions." *Journal of Climate* 16, no. 7: 1032–1046. [https://doi.org/10.1175/1520-0442\(2003\)016<1032:sasvoa>2.0.co;2](https://doi.org/10.1175/1520-0442(2003)016<1032:sasvoa>2.0.co;2).
- Rosan, T. M., L. E. O. Aragão, I. Oliveras, et al. 2019. "Extensive 21st-Century Woody Encroachment in South America's Savanna." *Geophysical Research Letters* 46, no. 12: 6594–6603. <https://doi.org/10.1029/2019GL082327>.
- Ross, J. 1975. "Radiative Transfer in Plant Communities." In *Vegetation and the Atmosphere*, edited by J. L. Monteith, vol. 1, 13–55. Academic Press. <https://cir.nii.ac.jp/crid/1571417124422508288>.
- Ruman, C. J., A. H. Monahan, and L. Sushama. 2022. "Climatology of Arctic Temperature Inversions in Current and Future Climates." *Theoretical and Applied Climatology* 150, no. 1–2: 121–134. <https://doi.org/10.1007/s00704-022-04147-9>.
- Rumpf, S. B., M. Gravey, O. Brönnimann, et al. 2022. "From White to Green: Snow Cover Loss and Increased Vegetation Productivity in the European Alps." *Science* 376, no. 6597: 1119–1122. <https://doi.org/10.1126/science.abn6697>.
- Rupp, D. E., S. L. Shafer, C. Daly, J. A. Jones, and S. J. K. Frey. 2020. "Temperature Gradients and Inversions in a Forested Cascade Range Basin: Synoptic- to Local-Scale Controls." *Journal of Geophysical Research: Atmospheres* 125, no. 23: e2020JD032686. <https://doi.org/10.1029/2020JD032686>.
- Rupp, D. E., S. L. Shafer, C. Daly, J. A. Jones, and C. W. Higgins. 2021. "Influence of Anthropogenic Greenhouse Gases on the Propensity for

- Nocturnal Cold-Air Drainage." *Theoretical and Applied Climatology* 146: 231–241. <https://doi.org/10.1007/s00704-021-03712-y>.
- Schnabel, F., R. Beugnon, B. Yang, et al. 2025. "Tree Diversity Increases Forest Temperature Buffering via Enhancing Canopy Density and Structural Diversity." *Ecology Letters* 28, no. 3: e70096. <https://doi.org/10.1111/ele.70096>.
- Schuchardt, M. A., B. J. Berauer, A. L. Duc, et al. 2023. "Increases in Functional Diversity of Mountain Plant Communities Is Mainly Driven by Species Turnover Under Climate Change." *Oikos* 2023, no. 11: e09922. <https://doi.org/10.1111/oik.09922>.
- Schuur, E. A. G., B. W. Abbott, R. Commene, et al. 2022. "Permafrost and Climate Change: Carbon Cycle Feedbacks From the Warming Arctic." *Annual Review of Environment and Resources* 47: 343–371. <https://doi.org/10.1146/annurev-environ-012220-011847>.
- Sehler, R., J. Li, J. T. Reager, and H. Ye. 2019. "Investigating Relationship Between Soil Moisture and Precipitation Globally Using Remote Sensing Observations." *Journal of Contemporary Water Research & Education* 168, no. 1: 106–118. <https://doi.org/10.1111/j.1936-704X.2019.03324.x>.
- Seidl, R., D. Thom, M. Kautz, et al. 2017. "Forest Disturbances Under Climate Change." *Nature Climate Change* 7, no. 6: 395–402. <https://doi.org/10.1038/nclimate3303>.
- Seneviratne, S. I., T. Corti, E. L. Davin, et al. 2010. "Investigating Soil Moisture–Climate Interactions in a Changing Climate: A Review." *Earth-Science Reviews* 99, no. 3–4: 125–161. <https://doi.org/10.1016/j.earscirev.2010.02.004>.
- Sfică, L., C. Beck, A.-I. Nita, M. Voiculescu, M.-V. Birsan, and A. Philipp. 2021. "Cloud Cover Changes Driven by Atmospheric Circulation in Europe During the Last Decades." *International Journal of Climatology: A Journal of the Royal Meteorological Society* 41, no. S1. <https://doi.org/10.1002/joc.6841>.
- Sherwood, S. C., M. J. Webb, J. D. Annan, et al. 2020. "An Assessment of Earth's Climate Sensitivity Using Multiple Lines of Evidence." *Reviews of Geophysics* 58, no. 4: e2019RG000678. <https://doi.org/10.1029/2019RG000678>.
- Shi, Y., and G. Wu. 2025. "Differences in Temperature Variation Between Winter and Summer Across China in Recent Decades." *International Journal of Climatology: A Journal of the Royal Meteorological Society* 45, no. 8: e8828. <https://doi.org/10.1002/joc.8828>.
- Slot, M., S. W. Rifai, C. E. Eze, and K. Winter. 2024. "The Stomatal Response to Vapor Pressure Deficit Drives the Apparent Temperature Response of Photosynthesis in Tropical Forests." *New Phytologist* 244, no. 4: 1238–1249. <https://doi.org/10.1111/nph.19806>.
- Soifer, L. G., J. L. Lockwood, J. J. Lembrechts, et al. 2025. "Extreme Events Drive Rapid and Dynamic Range Fluctuations." *Trends in Ecology & Evolution* 40, no. 9: 862–873. <https://doi.org/10.1016/j.tree.2025.06.009>.
- Song, J., S. Wan, S. Piao, et al. 2019. "A Meta-Analysis of 1,119 Manipulative Experiments on Terrestrial Carbon-Cycling Responses to Global Change." *Nature Ecology & Evolution* 3, no. 9: 1309–1320. <https://doi.org/10.1038/s41559-019-0958-3>.
- Sotomayor, D. A., and T. D. Drezner. 2019. "Dominant Plants Alter the Microclimate Along a Fog Gradient in the Atacama Desert." *Plant Ecology* 220, no. 4: 417–432. <https://doi.org/10.1007/s11258-019-00924-1>.
- Srivastava, A. K., M. Wehner, C. Bonfils, P. A. Ullrich, and M. Risser. 2024. "Local Hydroclimate Drives Differential Warming Rates Between Regular Summer Days and Extreme Hot Days in the Northern Hemisphere." *Weather and Climate Extremes* 45: 100709. <https://doi.org/10.1016/j.wace.2024.100709>.
- Stull, R. B. 1988. *An Introduction to Boundary Layer Meteorology*. Atmospheric and Oceanographic Sciences Library. Vol. 13. Kluwer Academic Publishers.
- Sturm, M., C. Racine, and K. Tape. 2001. "Increasing Shrub Abundance in the Arctic." *Nature* 411, no. 6837: 546–547. <https://doi.org/10.1038/35079180>.
- Terrer, C., R. B. Jackson, I. C. Prentice, et al. 2019. "Nitrogen and Phosphorus Constrain the CO₂ Fertilization of Global Plant Biomass." *Nature Climate Change* 9, no. 9: 684–689. <https://doi.org/10.1038/s41558-019-0545-2>.
- van der Wiel, K., and R. Bintanja. 2021. "Contribution of Climatic Changes in Mean and Variability to Monthly Temperature and Precipitation Extremes." *Communications Earth & Environment* 2, no. 1: 1–11. <https://doi.org/10.1038/s43247-020-00077-4>.
- Venter, Z. S., M. D. Cramer, and H.-J. Hawkins. 2018. "Drivers of Woody Plant Encroachment Over Africa." *Nature Communications* 9, no. 1: 1–7. <https://doi.org/10.1038/s41467-018-04616-8>.
- Virkkala, A.-M., P. Niittynen, J. Kemppinen, et al. 2024. "High-Resolution Spatial Patterns and Drivers of Terrestrial Ecosystem Carbon Dioxide, Methane, and Nitrous Oxide Fluxes in the Tundra." *Biogeosciences* 21, no. 2: 335–355. <https://doi.org/10.5194/bg-21-335-2024>.
- von Arx, G., E. Graf Pannatier, A. Thimonier, and M. Rebetez. 2013. "Microclimate in Forests With Varying Leaf Area Index and Soil Moisture: Potential Implications for Seedling Establishment in a Changing Climate." *Journal of Ecology* 101, no. 5: 1201–1213. <https://doi.org/10.1111/1365-2745.12121>.
- Wahid, A., S. Gelani, M. Ashraf, and M. Foolad. 2007. "Heat Tolerance in Plants: An Overview." *Environmental and Experimental Botany* 61, no. 3: 199–223. <https://doi.org/10.1016/j.envexpbot.2007.05.011>.
- Wallace, C. J., and M. Joshi. 2018. "Comparison of Land–Ocean Warming Ratios in Updated Observed Records and CMIP5 Climate Models." *Environmental Research Letters* 13, no. 11: 114011. <https://doi.org/10.1088/1748-9326/aae46f>.
- Ward, S. E., M. Schulze, and B. Roy. 2018. "A Long-Term Perspective on Microclimate and Spring Plant Phenology in the Western Cascades." *Ecosphere* 9, no. 10: e02451. <https://doi.org/10.1002/ecs2.2451>.
- Wever, N. 2012. "Quantifying Trends in Surface Roughness and the Effect on Surface Wind Speed Observations." *Journal of Geophysical Research* 117, no. D11. <https://doi.org/10.1029/2011jd017118>.
- Wilcox, E. J., K. E. Bennett, and J. Boike. 2024. "Bridging Gaps in Permafrost-Shrub Understanding." *PLOS Climate* 3, no. 3: e0000360. <https://doi.org/10.1371/journal.pclm.0000360>.
- Wilson, G., M. Green, J. Brown, et al. 2020. "Snowpack Affects Soil Microclimate Throughout the Year." *Climatic Change* 163, no. 2: 705–722. <https://doi.org/10.1007/s10584-020-02943-8>.
- Winkler, A. J., J. Kranz, A. Graf, et al. 2026. "The Phenology-Climate Feedback Loop: Current Understanding and Future Directions." *ESS Open Archive*. <https://doi.org/10.22541/essoar.177092640.00260655/v1>.
- Wright, A. J., and R. M. Francia. 2024. "Plant Traits, Microclimate Temperature and Humidity: A Research Agenda for Advancing Nature-Based Solutions to a Warming and Drying Climate." *Journal of Ecology* 112, no. 11: 2462–2470. <https://doi.org/10.1111/1365-2745.14313>.
- Wu, C., J. Peng, P. Ciais, et al. 2022. "Increased Drought Effects on the Phenology of Autumn Leaf Senescence." *Nature Climate Change* 12, no. 10: 943–949. <https://doi.org/10.1038/s41558-022-01464-9>.
- Zellweger, F., D. Coomes, J. Lenoir, et al. 2019. "Seasonal Drivers of Understorey Temperature Buffering in Temperate Deciduous Forests Across Europe." *Global Ecology and Biogeography: A Journal of Macroecology* 28, no. 12: 1774–1786. <https://doi.org/10.1111/geb.12991>.
- Zeng, H., P. Tian, M. Zhang, X. Cao, J. Liang, and L. Zhang. 2022. "Rapid Change in Surface-Based Temperature Inversions Across the World During the Last Three Decades." *Journal of Applied Meteorology and Climatology* 61, no. 2: 175–184. <https://doi.org/10.1175/jamc-d-21-0093.1>.

- Zeng, Z., A. D. Ziegler, T. Searchinger, et al. 2019. "A Reversal in Global Terrestrial Stilling and Its Implications for Wind Energy Production." *Nature Climate Change* 9, no. 12: 979–985. <https://doi.org/10.1038/s41558-019-0622-6>.
- Zha, J., C. Shen, Z. Li, et al. 2021. "Projected Changes in Global Terrestrial Near-Surface Wind Speed in 1.5°C–4.0°C Global Warming Levels." *Environmental Research Letters* 16, no. 11: 114016. <https://doi.org/10.1088/1748-9326/ac2fdd>.
- Zhan, C., R. Orth, H. Yang, et al. 2024. "Estimating the CO₂ Fertilization Effect on Extratropical Forest Productivity From Flux-Tower Observations." *Journal of Geophysical Research: Biogeosciences* 129, no. 6. <https://doi.org/10.1029/2023jg007910>.
- Zhang, L., M. Ding, T. Dou, Y. Huang, J. Lv, and C. Xiao. 2021. "The Shallowing Surface Temperature Inversions in the Arctic." *Journal of Climate* 34, no. 10: 4159–4168. <https://doi.org/10.1175/jcli-d-20-0621.1>.
- Zhang, S., D. Landuyt, K. Verheyen, and P. De Frenne. 2022. "Tree Species Mixing Can Amplify Microclimate Offsets in Young Forest Plantations." *Journal of Applied Ecology* 59, no. 6: 1428–1439. <https://doi.org/10.1111/1365-2664.14158>.
- Zhang, T. 2005. "Influence of the Seasonal Snow Cover on the Ground Thermal Regime: An Overview." *Reviews of Geophysics* 43, no. 4. <https://doi.org/10.1029/2004rg000157>.
- Zhang, W., K. Furtado, P. Wu, et al. 2021. "Increasing Precipitation Variability on Daily-to-Multiyear Time Scales in a Warmer World." *Science Advances* 7, no. 31. <https://doi.org/10.1126/sciadv.abf8021>.
- Zhang, X., X. Wang, C. M. Zohner, et al. 2025. "Declining Precipitation Frequency May Drive Earlier Leaf Senescence by Intensifying Drought Stress and Enhancing Drought Acclimation." *Nature Communications* 16, no. 1: 910. <https://doi.org/10.1038/s41467-025-56159-4>.
- Zhang, Y.-J., N. M. Holbrook, and K.-F. Cao. 2014. "Seasonal Dynamics in Photosynthesis of Woody Plants at the Northern Limit of Asian Tropics: Potential Role of Fog in Maintaining Tropical Rainforests and Agriculture in Southwest China." *Tree Physiology* 34, no. 10: 1069–1078. <https://doi.org/10.1093/treephys/tpu083>.
- Zhao, Z., P. De Frenne, J. Peñuelas, et al. 2022. "Effects of Snow Cover-Induced Microclimate Warming on Soil Physicochemical and Biotic Properties." *Geoderma* 423: 115983. <https://doi.org/10.1016/j.geoderma.2022.115983>.
- Zohner, C. M., L. Mirzagholi, S. S. Renner, et al. 2023. "Effect of Climate Warming on the Timing of Autumn Leaf Senescence Reverses After the Summer Solstice." *Science* 381, no. 6653: eadf5098. <https://doi.org/10.1126/science.adf5098>.

Supporting Information

Additional supporting information can be found online in the Supporting Information section. **Figure S1:** Model results of Case Study 1 for minimum near-surface temperature (5 cm above ground), where single meteorological variables (clouds, wind speed, air humidity, and snow cover) were modified in isolation to visualize their effect on microclimate warming and how it varies across a ca. 2 × 2 km² landscape on the border of Parque Natural Posets Maladeta in the Spanish Pyrenees (See [Supporting Information](#) for more modeling details and an interpretation of observed patterns). Left and middle columns show the minimum temperature of the 10 coldest days per year when setting the respective meteorological variables at low and high values, respectively, and keeping all other variables constant. Difference maps (right column) show the differences between high and low simulated values of the meteorological variables, that is, the effect of going from low to high values (Icons from Flaticon.com). **Figure S2:** Model results of Case Study 1 for maximum near-surface temperature (5 cm above ground), where single meteorological variables (clouds, wind speed, air humidity, and snow cover) were modified in isolation to visualize their effect on microclimate warming and how it varies across a ca.

2 × 2 km² landscape on the border of Parque Natural Posets Maladeta in the Spanish Pyrenees (See [Supporting Information](#) for more modeling details and an interpretation of observed patterns). Left and middle columns show the maximum temperature of the 10 warmest days per year when setting the respective meteorological variables at low and high values, respectively, and keeping all other variables constant. Difference maps (right column) show the differences between high and low simulated values of the meteorological variables, that is, the effect of going from low to high values (Icons from Flaticon.com). **Table S1:** Summary statistics of the microclimate change induced by changing atmospheric variables (cloudiness, wind speed, VPD, and snow cover). IQR, interquartile range; min/max, absolute minimum/maximum; q1/q3, first/third quartile; SD, standard deviation. **Table S2:** Summary statistics of the microclimate change induced by changing vegetation cover from grassland to shrub/tree dominated woodland. IQR, interquartile range; min/max, absolute minimum/maximum; q1/q3, first/third quartile; SD, standard deviation.